

UNIVERSAL
LIBRARY

OU_170638

UNIVERSAL
LIBRARY

170638

OSMANIA UNIVERSITY LIBRARY

Call No. 633.51 / D455 Accession No. 21479
Author Deshpandeey and Munsakarney
Title Spotted boll - worms of Cotton. 1928

This book should be returned on or before the date last marked below.

Scientific Monograph No. 10.

The Imperial Council of Agricultural Research

**THE SPOTTED BOLL-WORMS OF COTTON (*EARIAS
FABIA* STOLL. AND *EARIAS INSULANA* BOISD.) IN
SOUTH GUJARAT, BOMBAY PRESIDENCY.**

**(Final report on investigations financed by the Indian
Central Cotton Committee, 1923 to 1931)**

BY

B. P. DESHPANDE, M.Ag.,

and

N. T. NADKARNY, B.Ag.



DELHI · MANAGER OF PUBLICATIONS

1936

Price Rs. 5-14 or 9s. 6d.

List of Agents from whom Government of India Publications are available.

ENGLAND.

THE HIGH COMMISSIONER FOR INDIA, INDIA HOUSE, ALDWYCH, LONDON, W. C. 2.

PALESTINE.

Stelmatsky, Jerusalem.

INDIA.

(a) PROVINCIAL GOVERNMENT BOOK DEPOTS.

MADRAS :—Superintendent, Government Press, Mount Road, Madras.

BOMBAY :—Superintendent, Government Printing and Stationery, Queen's Road, Bombay.

SIND :—Manager, Sind Government Book Depot, Karachi (Sadai).

UNITED PROVINCES OF AGRA AND OUDH :—Superintendent of Government Press, United Provinces of Agra and Oudh, Allahabad.

PUNJAB :—Superintendent, Government Printing, Punjab, Lahore.

BURMA :—Superintendent, Government Printing, Burma, Rangoon.

CENTRAL PROVINCES AND BERRAR :—Superintendent, Government Printing, Central Provinces, Nagpur.

ASSAM :—Superintendent, Assam Secretariat Press, Shillong.

BIHAR AND ORISSA :—Superintendent, Government Printing, Bihar and Orissa, P. O. Hulzbarbagh, Patna.

NORTH-WEST FRONTIER PROVINCE :—Manager, Government Printing and Stationery, Peshawar.

(b) PRIVATE BOOK-SELLERS.

Acro Stores, Karachi City *

Albert Library, Dacca

Banerjee & Bros, Ranchi, Messrs. G.

Bantulya & Co., Ltd., Station Road, Ajmer

Bengal Flying Club, Dum Dum Cant *

Bhawanani & Sons, New Delhi

Book Company, Calcutta.

Booklover's Resort, Talukad, Trivandrum, South India

Burma Book Club, Ltd., Rangoon.

Butterworth & Co. (India), Ltd., Calcutta

Calcutta Book Agency, 16-1, Shama Chatterjee Street, Calcutta

Chatterjee & Co., 3, Bacharam Chatterjee Lane, Calcutta

Chatterjee, Chatterjee & Co., Ltd., 11, College Square, Calcutta

City Book Co., Madras.

City Book House, Weston Road, Cawnpore.

Commercial Book Co., Lahore

Das Gupta & Co., 51/3, College Street, Calcutta

Deccan Bookstall, Poonia 4

Delhi and U. P. Flying Club, Ltd., Delhi.

English Book Depot, Patna

English Book Depot, Taj Road, Agra, and Sadul Raj, Jhansi

English Book Depot, Bank Road, Ambala Cantonment

and Kasauli

English Bookstall, Karachi

Fakir Chand Marwari, Peshawar Cantonment

Fono Book Agency, Simla

Gaya Prasad & Sons, Agra

Graha Mandir, Cuttack

Higginsbothams, Madras

Hindu Library, 137-F, Balaram De Street, Calcutta

Hyderabad Book Depot, Chaderghat, Hyderabad

(Deccan)

Imperial Book Depot and Press, near Jamia Masjid

(Machhliwadan), Delhi

Indian Army Book Depot, Dayalbagh, Agra

Indian Army Book Depot, Jullundur City and Daryaganj, Delhi

Indian Book Shop, Benares City

Indian School Supply Depot, Central Avenue, South, P. O.

Dharamtala, Calcutta

Insurance Publicity Co., Ltd., Lahore

International Book Service, Poonia 4

Jaina & Bros, Mori Gate, Delhi, and Connaught Place, New

Delhi, Messrs. J. M.

James Murray & Co., 12, Govt. Place, Calcutta (for

Meteorological publications only)

Kali Charan & Co., Municipal Market, Calcutta.

Kamala Book Depot, 15, College Square, Calcutta

Kamala Book Stores, Bankipore, Patna

Karnataka Publishing House, Bangalore City

Kende & Co., Karachi

Krishnaswami & Co., Teppakulam P. O., Trichinopoly

Fort, Messrs. S.

Lahiri & Co., Calcutta, Messrs. S. K.

Law Printing House, 11, Mount Road, Madras

Law Publishing Co., Mylapore, Madras

Lawrence and Mayo, Ltd., Bombay (for Meteorological

publications only)

Local Self-Govt. Institute, Bombay.

London Book Co. (India), Arab Road, Peshawar, Murree,

Nowshera and Rawalpindi

London Book Depot, B. I. Bazar, Bareilly, U. P.

Malhotra & Co., Post Box No. 94, Lahore, Messrs. U. P.

Modern Book Depot, Bazar Road, Sialkot Cantonment.

Mohammad Ghouseh Shah, Rajkot

Motilal Banarsi Das, Oriental Book Depot, Sadmitha

Street, Lahore.

Nandkishore & Bros*, Chowk, Benares City

Natelson & Co., Publishers, George Town, Madras

Messrs. G. A.

New Book Co., "Kilab Mahal", 192, Hornby Road,

Bombay.

Newman & Co., Ltd., Calcutta, Messrs. W.

North India Christian Tract and Book Society, 18, Chive

Road, Allahabad.

Oriental Book Supplying Agency, 15, Shukrawar, Poonia

City.

Oxford Book and Stationery Company, Delhi, Lahore,

Simla, Meerut and Calcutta.

Parish & Co., Baroda, Messrs. B.

Pioneer Book Supply Co., 20, Shih Narayan Das Lane,

Calcutta, and 21B, Cloth Market, Delhi

Popular Book Depot, Grant Road, Bombay

Punjab Religious Book Society, Lahore

Rachmath Prasad & Sons, Patna City

Ram Krishna Bros., Opposite Bishrambag, Poonia City.

Ram Narain Lal, Katra, Allahabad

Rama Krishna & Sons, Booksellers, Anarkali, Lahore

Ramesh Book Depot, Stationary Mart, Kashmiri Gate,

Delhi

Ray & Sons, 43, K. & I. Edwardes Road, Rawalpindi,

Murree and Peshawar, Messrs. J.

Ray Chowdhury & Co., 119, Ashutosh Mukherjee Road,

Bhowanipore, Calcutta.

Rochoose & Sons, Madras.

Roy Chowdhury & Co., 11, College Square, Calcutta,

Messrs. N. M.

Sampson William & Co., 127-B, The Mall, Cawnpore

Sargar & Sons, 15, College Square, Calcutta, Messrs.

M. C.

Sarkar & Co., Ltd., 18, Shama Charan De Street and

8/2, Hastings Street, Calcutta, Messrs. P. C.

Scientific Publishing Co., 9, Talola Lane, Calcutta.

Seshachalam & Co., Masulipatam, Messrs. M.

Sharma & Co., P. O. Chaulanganj, Cuttack

Shri Shankar Karantaka Pustaka Bhandara, Ma'nauddi,

Dharwar.

S. P. Bookstall, 21, Badliwar, Poonia.

Standard Book Depot, Lahore, Dalhousie and Delhi

Standard Bookstall, Karachi.

Standard Bookstall, Quetta.

Standard Law Book Society, 5, Hastings Street, Calcutta

Standard Literature Company, Ltd., Calcutta.

Students' Popular Depot, Kachari Road, Lahore.

Surat and District Trading Society, Surat.

Taraporewalla Sons & Co., Bombay, Messrs. D. B.

Thacker & Co., Ltd., Bombay

Thacker, Spink & Co., Ltd., Calcutta and Simla

Tripathi & Co., Booksellers, Princess Street, Kailadevi

Road, Bombay, Messrs. N. M.

Union Stores, Indore City.

University Book Agency, Kachari Road, Lahore

Upper India Publishing House, Ltd., Literature Palace,

Annundadnan Park, Lucknow.

Varadachary & Co., Madras, Messrs. P.

Venkatasubban, A., Law Bookseller, Vellore.

Wheeler & Co., Allahabad, Calcutta and Bombay,

Messrs. A. H.

Young Man & Co., Ajmer and Egerton Road, Delhi,

* Agents for publications on Aviation only.

PREFACE

The investigations on the Spotted Boll-worms described in this Monograph were carried out at Surat from September 1923 till March 1931, by a group of workers appointed by the Bombay Department of Agriculture. Mr. Ramrao S. Kasargode was in charge of them during the first five years, after which he had to retire prematurely on account of ill health and consequently they were conducted by me during the remaining four years. My thanks are due to my colleagues, Messrs. S. G. Kadkol, N. T. Nadkarny, B. J. Thakar, M. S. Patel and B. K. Desai for their sincere co-operation. I am greatly indebted to Dr. W. Burns for the guidance and encouragement which I received from him during the period of the last four years of this work. Throughout the period from 1923 to 1931, these investigations were financed by the Indian Central Cotton Committee.

As a result of these investigations, it was found that the 'clean-up' measures, described on p. 91, offered the most promising means of controlling the Spotted Boll-worms. A scheme was, therefore, started in the Broach District in April, 1931, in order to determine the efficiency of these measures and this is still in progress. This scheme also is financed by the Indian Central Cotton Committee.

BROACH,

B. P. DESHPANDE.

31st July 1933.

CONTENTS

| | PAGE |
|--|------|
| Preface | i |
| Contents | ii |
| List of illustrations | iii |
| Introduction | 1 |
| Summary | 3 |
| Life-history and habits of the Spotted Boll-worms | 6 |
| Damage to the cotton crop by the Spotted Boll-worms | 19 |
| Abundance of the Spotted Boll-worms in the cotton fields at Surat | 26 |
| Behaviour of the cotton plant when it is grown free from Boll-worms and estimation of the improvement in the yield of <i>kapas</i> | 32 |
| Study of control measures | 43 |
| (1) Removal of attacked shoots | 43 |
| (2) Trap crops | 45 |
| (3) Parasites of the Spotted Boll-worms | 49 |
| (4) Deterrents | 62 |
| (5) Attractants | 63 |
| (6) Soil-mulch as a control measure | 65 |
| (7) Insecticides | 72 |
| (8) Clean-up measures | 91 |
| APPENDIX I. | |
| Tables I to L | 101 |
| APPENDIX II. | |
| Spinning Test Reports | 150 |
| APPENDIX III. | |
| Description of the multiple temperature incubator | 155 |
| APPENDIX IV. | |
| Spotted Boll-worms in Khandesh | 157 |
| APPENDIX V. | |
| <i>Phycita infusella</i> Meyr. (cotton shoot-roller) at Surat | 171 |
| APPENDIX VI. | |
| Pink Boll-worm (<i>Platyedra gossypiella</i> Saund.) in South Gujarat | 180 |
| APPENDIX VII. | |
| Pink Boll-worm (<i>Platyedra gossypiella</i> Saund.) in Khandesh | 196 |
| APPENDIX VIII. | |
| Capacity of the cotton plant to replace its forms shed due to boll-worm injury | 202 |
| References | 208 |

LIST OF ILLUSTRATIONS

| | | PAGE. |
|--------------------|---|-------|
| Plate I, Fig. 1. | <i>Earias</i> pupae in the soil | 12 |
| | 2. A cotton shoot damaged by a caterpillar of the Spotted Boll-worms | 12 |
| " II | Showing periodical flower-bud formation, total bud-shedding and the bud-shedding due to boll-worms from cotton plants at Surat during three seasons | 22 |
| " III | Showing periodical flower opening, boll-shedding and the boll-shedding due to boll-worms from cotton plants at Surat during three seasons | 23 |
| " IV | Showing the distribution of the population of <i>Earias</i> larvae on different parts of 25 plants during 6 consecutive seasons at Surat | 26 |
| " V | Showing incidence of larval population of <i>Earias</i> larvae on 25 cotton plants and weekly rainfall during 6 consecutive seasons at Surat | 30 |
| " VI | Showing mean night temperatures and the periodical population of <i>Earias</i> larvae, during 4 consecutive seasons at Surat | 31 |
| " VII, Fig. 1. | Plant cage with wire gauze sides and top | 32 |
| " 2. | Cotton plant grown free from boll-worms in a plant house with sides and top of wire gauze | 32 |
| " VIII, Fig. 1. | A cotton plant grown free from boll-worm by night-caging. (Note the tin trench round the stem of the plant) | 33 |
| " 2. | Plant curtains used for night-caging | 33 |
| " IX, Fig. 1 | Right—Cotton plant protected from boll-worm by night-caging; Left—Unprotected plant | 34 |
| " 2. | Diagrammatic sketch of a cotton plant grown free from boll-worm by night-caging | 34 |
| " X | Showing weekly performance of caged and uncaged plants during 3 seasons at Surat | 35 |
| " XI | Showing weekly flower bud-formation, shedding of flower-buds, opening of flowers and shedding of bolls from caged and uncaged plants at Surat during 1926-27 | 35 |
| " XII | Showing periodical flower opening and relative success of bolls from dusted and undusted plants at Surat during 1926-27 | 74 |
| " XIII | Early opening of bolls in plot (left) dusted with calcium arsenate | 75 |
| " XIV | Showing weekly flower-opening and relative success of bolls in cotton plots dusted and undusted with insecticides | 77 |
| " XV | Showing weekly flower-opening, shedding of bolls and bolls shed due to boll-worms in a cotton plot dusted with calcium arsenate during the period of shoot attack, and also in the control plot | 84 |
| " XVI, Fig. 1 | Removing the cotton plants with a light pick axe | 96 |
| " 2. | A shoot sprouting from a stump of cotton plant | 96 |
| " XVII, Fig. 1. | A shoot sprouting from a side root of a stump of cotton plant | 97 |
| " 2. | A cotton plant puller | 97 |
| Plates XVIII & XIX | Showing sources of plant food of the Spotted Boll-worms during the period between two cotton-growing seasons at Surat | 98 |
| Plate XX | A cotton plant puller in action | 100 |
| " XXI | Plan and Section of a multiple temperature incubator | 155 |
| " XXII | Showing weekly rainfall and periodical population of <i>Earias</i> larvae on cotton and <i>bhendi</i> plants at Jalgaon during 1929 and 1930 | 159 |
| " XXIII | Showing the weekly flower-opening, shedding of bolls and relative success of bolls from weekly flowers on cotton plants at Jalgaon | 162 |
| " XXIV | Showing sources of food for the Spotted Boll-worms during the period between two cotton-growing seasons in Khandesh | 170 |
| " XXV | Showing weekly flowering, proportion of flowers pruned and the number of successful bolls from weekly flowers in the flower-pruning experiments of 1926-27 | 204 |

THE SPOTTED BOLL-WORMS (*EARIAS FABIA* STOLL. AND *EARIAS INSULANA* BOISD.) IN SOUTH GUJARAT

BY

B. P. DESHPANDE, M.Ag.,

AND

N. T. NADKARNY, B.Ag.

(Received for publication on 31st March 1933)

INTRODUCTION

Earias is a widely-distributed genus, occurring all over the world, North and South America [Wardle 1929]. Its occurrence as a pest on cotton has particularly been reported from India, Egypt, West Africa, South Africa, Uganda, Portuguese East Africa, South Rhodesia and Fiji.

Boll-worms were known to affect the cotton crop in India prior to 1903, but they came into prominence in 1905 and 1906 [Fletcher and Misra 1921].

During the year 1905, the cotton crop in the Punjab was threatened with a failure due to the attack of some caterpillars and green fly (aphis), [Renouf 1907]. The Director of Agriculture of that Province communicated the facts to the Entomologist to the Government of India, whose observations showed that the failure was largely due to the ravages of the Spotted Boll-worms (especially *Earias insulana* Boisd.). He attributed the unusual abundance of this pest during that season to the rigorous cold of the previous winter, which he thought had destroyed the natural controls, viz., the parasites of these worms [Lefroy 1906]. The anxiety about the possible recurrence of this pest in the succeeding season was keenly felt and hence the measures mentioned below, as recommended by him, were advocated.

- (1) Cutting and burning of all cotton and *bhendi* (*Hibiscus esculentus*) stalks during the winter.
- (2) The use of *bhendi* as a trap crop.
- (3) The re-introduction of the parasite, *Rhyssalus lefroyi*.

The cotton crop in the Punjab during 1906 turned out to be a normal one and therefore Lefroy [1907] inferred that his faith in the re-introduction of the parasites was amply justified.

“From 1906 until 1911, the varieties of cotton grown at Pusa were mostly utilised for the despatch of the parasites to the Punjab, whenever required. It was in the year 1912 that regular sowings of cotton were made and details of observations

started regarding the boll-worms, *E. fabia* and *E. insulana* as well as *Platyedra gossypiella*." As a result of these investigations, Fletcher and Misra [1921] presented a few of their conclusions in their bulletin on cotton boll-worms. Besides the re-introduction of parasites, the control measure which appealed to them most, was the removal and burning of the cotton plants, which are attacked in their shoots by the boll-worms, before the final thinning of the seedlings is done. A larger seed rate was recommended to enable this to be done effectively, and without detriment to the interest of the cultivators.

During 1913, Madan Mohan Lal prepared a preliminary report on cotton boll-worms in the Punjab and ascertained that the early summer rains and the parasite *Rhogas lefroyi* effectively checked the prevalence of the worms. If, however, these natural agencies were unfavourable, he suggested the destruction of the early brood of boll-worms in the cotton buds and flowers; and the introduction of the parasite *Rhogas lefroyi*, by parasite boxes.

In the Entomological Meeting at Pusa during 1921, Javeri reported satisfactory results with the use of *bhendi* (*Hibiscus esculentus*) as a trap crop for controlling the boll-worms.

At the next Entomological Meeting at Pusa, Husain and Mathur [1923] presented very useful information about the parasites of the Spotted Boll-worms.

Later on Patel [1924] attributed an enormous loss of early flower-buds and bolls to these worms at Surat, and argued that a tall open habit of the bracts increased the resistance to boll-worms, probably because the parasite (*Microbracon lefroyi* D. and G.) could more easily attack the worms in the flower-buds and bolls of such strains.

This is a short history of the efforts made at intervals by different workers in India for studying this pest and for finding out remedial measures. Ultimately the position with regard to this pest was reviewed by the Indian Central Cotton Committee, Bombay [Burt, 1922] and investigations on the Spotted Boll-worms were started at Surat in September 1923, for ascertaining the loss caused by this pest to the cotton crop in this tract, and for studying control measures. These investigations, which were conducted at Surat from 1923 till 1931, form the subject matter of this publication.

The work was divided into three parts *viz.*, (1) detailed study of life-history and habits of the Spotted Boll-worms, (2) estimation of the damage caused by this pest to the cotton crop, together with the study of the possibilities of improvement in the crop if the pest can be eliminated, and (3) study of control measures.

During the progress of these investigations it was possible to make a few observations on two more pests of cotton, *viz.*, the cotton shoot-roller (*Phycita infusella* Meyr.) and the Pink Boll-worm (*Platyedra gossypiella* Saund.). These observations are included in the Appendices.

Before proceeding further, it is essential to mention that although the Spotted Boll-worms appear on the cotton crop every year in Gujarat, there does not exist any serious complaint against this pest from the cultivators as it is difficult for them to appreciate the damage caused by these insects, due to the peculiar method of feeding of the caterpillars. In the first place, cultivators do not mind the destruction of the cotton shoots, because even otherwise they consider the pruning of the shoots essential for the development of the side branches. Secondly, the larvae feeding in the flower-buds do not attract the attention of an untrained eye, and the shedding of forms caused by these worms passes off as natural shedding which must occur; because they find that out of the large number of flower-buds produced by the cotton plants, only a very few develop into mature bolls, and the remainder drop away as flower-buds or young bolls. In the normal course, therefore, all this imperceptible damage does not alarm the cultivators, and it is often very difficult to convince them of the loss caused by this pest to their cotton crop. It will not be out of place to quote a few remarks made by Lefroy [1906] in this connection while writing about the Bombay locusts.

“The immense swarms and the extraordinary movements of the Bombay locusts give it an enormously exaggerated importance; it fills the imagination and appeals to the human mind. The damage done by the locusts is probably far less than that done yearly by two pests of cotton (*Earias fabia* and *Earias insulana*) or by the moth borer in cane and sorghum (*Chilo simplex*). In point of numbers the locusts are far below the probable numbers of these pests in any one year; in point of destruction and actual loss of crop, the ordinary pests are far ahead of the Bombay locust in its worst years.”

SUMMARY

Attention was prominently drawn to the Spotted Boll-worms, for the first time, during the year 1905, due to the failure of the cotton crop in the Punjab. Since then, introduction of parasites, destruction of the damaged plants in the early part of the season and the use of *bhendi* (*Hibiscus esculentus*) as a trap crop, were some of the measures which were advocated for the control of this pest. The position with regard to these worms was reviewed by the Indian Central Cotton Committee during 1922, and at their instance, further investigations on the Spotted boll-worms were started at Surat in 1923. This work continued till 1931 and the present publication embodies the results of these investigations.

The Spotted Boll-worms are active throughout the year, and the period of their life varies between 22 and 35 days. Oviposition takes place only during the night and the moths were observed to lay, on an average, 432 eggs. The larvae invariably go to the soil for pupation in this locality. Both *Earias fabia* and *Earias insulana* are present but it is the former which predominates on cotton, almost throughout the year.

A vigorous growth of the cotton crop commences soon after the close of the monsoon, and the Spotted Boll-worms appear in the cotton fields as soon as the plants are about 9 inches in height. In the beginning, vegetative shoots are destroyed in large numbers, usually during the months of September and October. The larvae turn their attention to the flower-buds as soon as these begin to develop, and the damaged buds drop from the plants. Observations during three years from 1925 to 1928, showed that 34 to 51 per cent of the buds were damaged by these insects out of the total number of buds which dropped from the plants. A part of the larval population diverts its attention to the bolls as soon as they begin to appear. Young bolls drop if they are attacked by these worms, but the older ones continue to stick to the plants, and yield a reduced quantity of damaged *kapas*. During the three years mentioned above, 20 to 69 per cent of the shed bolls were found damaged by these caterpillars.

It was noticed that the rapid multiplication of the pest during the early part of the cotton-growing season was considerably checked, if heavy showers of rain were received during that period. The larval population of *Earias* on cotton was comparatively very high during November and December, and normally it declined rapidly at the end of December or the beginning of January. This decline was partly due to the cold weather and partly due to the activities of parasites.

A method, which has been described as the night-caging method was devised for growing normal cotton plants free from boll-worms for experimental purposes. Trials of growing more than 100 plants by this method, during each of the three seasons, from 1928-29 to 1930-31, showed that the protected plants yielded 81, 54 and 7 per cent more *kapas* than the respective control plants. During the last season the increase was markedly low, because the caged plants suffered heavily from a storm which occurred at the end of October, when these plants had a very much large number of forms on them than the control plants. On the caged plants the bolls opened about 4 to 6 weeks earlier than on the control plants, because the earliest growing forms on these plants were saved from the Spotted Boll-worms.

Measures which were studied for ascertaining the possibilities of controlling this pest consisted of the removal of the attacked shoots; *bhendi* (*Hibiscus esculentus*) as a trap crop; attractants and deterrents for the moths; parasites; soil-mulch; insecticides; and the prevention of the carry-over of this pest from one season to another.

The removal of attacked shoots was not found to be very useful because only a small fraction of the total population of the *Earias* larvae could be destroyed by collecting the shoots which showed the damage prominently.

The use of *bhendi* as a trap crop also appeared to be of questionable utility, because it was found that these insects multiplied more rapidly on *bhendi* than on cotton, and that the disadvantages of encouraging the cultivation of this crop in the midst of cotton seemed to outweigh the advantages.

Cynogas dust and flowers of sulphur were tried as deterrents for the moths, but both of them were found to be injurious to the cotton plants.

Several substances were tried for attracting the *Earias* moths. Cotton-seed cake and sesamum cake, exposed in the cotton fields after mixing with large quantities of water, showed a certain amount of attraction both for the Spotted and Pink Boll-worm moths. The attraction, however, was not sufficiently strong for using these baits as control measures.

All the known important parasites of *Earias* were found to be present at Surat, and there was, therefore, no possibility of importing any of them from outside. A detailed study of the life and habits of the most important parasite, viz. *Microbracon lefroyi* D. and G., showed that it was not practicable to increase its utility by artificial means.

Preliminary trials indicated that the presence of fine soil-mulch in the cotton fields prevented the larvae from going deep into the soil for pupation, and thus exposed them to the heat of the sun and to their predators. It was also observed that if any of the pupae happened to be buried below a layer of an inch or two of pulverised soil, most of the moths emerging from them could not make their way out of the soil. Further critical observations, however, showed that it was impracticable to maintain the required type of soil mulch in the fields, on a large scale, for its successful use as a control measure.

It was ascertained that spraying or dusting the cotton plants with lead arsenate or Paris Green was not useful in controlling this pest. Later on, it was found that calcium arsenate and sodium silico-fluoride were very effective in reducing the population of *Earias* larvae in the cotton fields, because the larvae were killed in large numbers on the dusted plants (especially the tiny ones, which wandered about before commencing to feed after emerging from the eggs). Serious aphid infestation, however, developed on the cotton plants dusted with these insecticides, a few weeks after the use of these poisons, and it proved to be a great obstacle in the way of using these substances for combating the boll-worms. It appeared that the winged aphid were attracted to the plants dusted with these white poisons, and they could multiply there very rapidly because it seemed that the insecticides afforded them immunity from their enemies. Blackening these insecticides, before using, did not serve entirely to get over this trouble. It is considered that insecticides like calcium arsenate and sodium silico-fluoride could be successfully used against these insects if a cheap and efficient method could be discovered for preventing the subsequent trouble, viz., aphid infestation.

The cotton crop in this locality is harvested in the month of March or April, and the next crop is normally sown after the third week of June, according to the condition of the monsoon. The Spotted Boll-worms are active throughout the year and they get their supply of food, between the two cotton-growing seasons, from the fresh shoots sprouting from the standing plants of cotton and their stumps; from the

bhendi plants growing during that period ; and from the stray malvaceous weeds, if any of them happen to be in green condition at that time. These observations show that it will be possible to prevent the carry-over of this pest from one season to another, by cutting off its supply of food during the period between the two cotton-growing seasons. This can be achieved if all the cotton plants are uprooted immediately after the harvest of *kapas* ; if the cultivation of *bhendi* is prevented during this period ; and if stray malvaceous weeds found in green condition during summer are destroyed.

A small implement [Plate XVII, fig. 2 and Plate XX] has been devised by us for uprooting the cotton plants, and trials are now in progress for judging the efficiency of the above mentioned clean-up measures in a selected area of about 480 square miles in Gujarat.

LIFE-HISTORY AND HABITS OF THE SPOTTED BOLL-WORMS

The Spotted Boll-worms which damage the cotton crop in Gujarat consist of two species, viz., *Earias fabia* Stoll., and *Earias insulana* Boisd. Both these are meant when the term Spotted Boll-worms is used in this publication, except where one or the other is specifically mentioned.

LIFE OF THE MOTHS

The Spotted Boll-worm moths emerge from their cocoons after sunset till mid-night. Only very rare instances of their emergence after mid-night have been noted, whereas they do not emerge during the day time.

TABLE I

Time of emergence of the Spotted Boll-worm moths

| Time | NUMBER OF MOTHS EMERGED | |
|---|-------------------------|------------|
| | 1st case | 2nd case |
| Between 6 P.M. and 8 P.M. | 16 | 8 |
| Between 8 P.M. and 10 P.M. | 27 | 157 |
| Between 10 P.M. and mid-night | 5 | 48 |
| Between mid-night and 6 A.M. | Nil | 1 |
| TOTAL EMERGENCE . | 48 | 214 |

Copulation of the moths takes place in the early hours of the morning. A few of the moths copulate on the night of their emergence, only a few hours after they come out of their cocoons. In most of the cases, however, mating takes place on the 2nd or the 3rd night.

TABLE II

Day of mating after emergence

(Total cases observed = 80 pairs)

| — | On the night of emergence | On 2nd night | On 3rd night | On 4th night | On 5th night | On 6th night |
|---------------------------------------|---------------------------|--------------|--------------|--------------|--------------|--------------|
| Number of pairs of moths which mated. | 10 | 38 | 18 | 7 | 4 | 3 |

The mating generally extends from 1 to 3 hours. The moths usually separate soon after day-break, but during November and December, a few pairs can often be found resting on the leaves of the cotton plants till 10 A.M.

A good proportion of the moths begins to lay eggs on the first night after their mating, but most of the remaining moths commence oviposition before the fourth night. In a few cases, however, it was noticed that oviposition was delayed for about a week.

TABLE III

Number of moths which commenced oviposition on consecutive nights, after mating

| — | 1st night | 2nd night | 3rd night | 4th night | 5th night | 6th night | 7th night | 8th night | Total moths observed |
|---|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|----------------------|
| Number of moths which commenced oviposition.] | 34 | 20 | 10 | 2 | 3 | 1 | .. | 1 | 71 |

For various observations the moths were kept in the laboratory in cylindrical glass jars 5 inches in height and $3\frac{1}{2}$ inches in diameter, and the jars were covered with pieces of thin cloth. Moths were fed on sugar solution or diluted honey, which was applied with brushes to the cloth covers.

In captivity, the average life of a moth was found to vary from about 8 to 22 days. A few moths were, however, observed to live for over 30 days, but out of the large number observed, there was only one which lived for 34 days. (App. I, Table I).

TIME OF OVIPOSITION

Casual observations, while handling the rearing cages of moths, gave an impression that the moths laid their eggs during the night-time only. It was surmised that the absence of light may be one of the requirements for oviposition, and therefore a few pairs of moths were kept in perfectly dark chambers during the day time. This, however, did not induce them to lay eggs by day. Definite observations were therefore undertaken to ascertain the exact time of their oviposition.

Moths which had emerged in the laboratory were selected and eleven pairs of female and male moths were arranged. One pair of moths was allotted for each hour of the night, commencing from 8 P.M. The eggs laid by each pair, before and after the allotted hour, were recorded. Thus the first pair was used to note the number of eggs laid by a female moth from 6 P.M. to 8 P.M. and during the rest of the night after 8 P.M. The second pair showed the number of eggs laid before 9 P.M. and after 9 P.M. The other pairs were similarly allotted for each of the remaining hours of the night.

TABLE IV

Number of eggs laid by each female during the period of 5 days of observation

| Pair No. | Total No. of eggs laid | Eggs laid before the allotted hour | | | Eggs laid after the allotted hour | | |
|----------|------------------------|------------------------------------|-----------|-------|-----------------------------------|-----------|--|
| 1 | 390 | 230 before | 8 P.M. | . . . | 160 after | 8 P.M. | |
| 2 | 253 | 154 „ | 9 P.M. | . . . | 99 „ | 9 P.M. | |
| 3 | 351 | 274 „ | 10 P.M. | . . . | 77 „ | 10 P.M. | |
| 4 | 383 | 359 „ | 11 P.M. | . . . | 24 „ | 11 P.M. | |
| 5 | 397 | 374 „ | mid-night | . . . | 23 „ | mid-night | |
| 6 | 406 | 384 „ | 1 A.M. | . . . | 22 „ | 1 A.M. | |
| 7 | 392 | 368 „ | 2 A.M. | . . . | 24 „ | 2 A.M. | |
| 8 | 320 | 306 „ | 3 A.M. | . . . | 14 „ | 3 A.M. | |
| 9 | 268 | 250 „ | 4 A.M. | . . . | 18 „ | 4 A.M. | |
| 10 | 398 | 388 „ | 5 A.M. | . . . | 10 „ | 5 A.M. | |
| 11 | 434 | 434 „ | 6 A.M. | . . . | Nil „ | 6 A.M. | |

(For details of daily oviposition *vide* App. I, Table II).

These trials showed very clearly the proportion of eggs laid before and after the various hours of the night ; that the oviposition was restricted to night-time only ; and that 94 per cent of the eggs were laid between 6 p.m. and mid-night and the remaining 6 per cent were spread over the other six hours of the night.

These observations were later useful for devising a method of growing experimental cotton plants free from the Spotted Boll-worms. The details of this method are discussed separately (Page 33).

EGG-LAYING CAPACITY

In calculating the rate of increase of the Spotted Boll-worms, Lefroy [1906, 2] had assumed 60 as the normal number of eggs laid by a single moth. At Surat, however, it was found that the capacity of oviposition of these moths was considerably higher. Trials during the months of August and September of 1924, showed that a moth could lay as many as 432 eggs on an average (App. I, Table III). It was also noticed that the largest number of eggs is laid during the first few days of oviposition. The above trials showed that 84 per cent of the eggs were laid during the first seven days of the period of egg-laying. The highest number of eggs laid by one moth in a single night was found to be 162, and the maximum number of eggs oviposited by a moth during all its life was 697.

Egg-laying capacity is not constant throughout the year. It declines considerably during the cold months of December and January as can be seen from the following results :—

- (1) An average of 341 eggs per moth was recorded from 14 pairs which were under observation in November 1930.
- (2) This average went down to 298 eggs per moth in December 1930 when another lot of 14 pairs was under observation.
- (3) An average of only 185 eggs per moth was recorded in January 1931 when 8 pairs were kept under observation. (App. I, Table 1).

It was further noted that the food on which the *Earias* larvae are reared has a great influence on the capacity of the moths for oviposition. During September 1929, one set of larvae was fed exclusively on tender shoots of cotton plants ; and *bhendi* pods (*Hibiscus esculentus*) alone were offered to the second set of larvae. The moths emerging from both the sets were kept separately and the eggs laid by them were recorded. In the first case (where the larvae were fed on cotton shoots), the moths laid, on an average, 88 eggs, whereas, in the second case (where the larvae were fed on *bhendi*) the moths could lay 399 eggs each. This trial was repeated during February 1930, when 3 sets of larvae were reared exclusively on flower-buds of cotton, bolls and *bhendi* pods respectively ; and the moths which developed from them were kept separately for oviposition.

TABLE V

Eggs laid by moths in the above three cases

| | No. of moths observed | Average No. of eggs laid |
|---|--------------------------|-----------------------------|
| 1. Moths reared from larvae fed on flower-buds . . . | 9 | 196 |
| 2. Moths reared from larvae fed on bolls . . . | 12 | 342 |
| 3. Moths reared from larvae fed on <i>bhendi</i> pods . . . | 12 | 451 |

WHERE THE EGGS ARE LAID

The eggs are laid singly and cannot be said to be properly fixed to the plants. They are entangled in the hairy parts of the bushes. The moths prefer hairy objects to non-hairy ones for depositing their eggs. When hairy and non-hairy stems of plants are offered to them simultaneously, it is always the hairy ones which have the largest number of eggs on them. In one of the trials *bhendi* pods (*Hibiscus esculentus*) and the fruits of *Abutilon* (*Abutilon indicum*) were simultaneously placed in the cages of the moths for oviposition. (The fruits of *Abutilon* are very hairy and give a velvety appearance). The result was that 92.4 per cent of the total eggs were deposited in the hairy surface of the *Abutilon* capsules and the remaining 7.6 per cent only on the *bhendi* pods.

TABLE VI

Number of eggs laid on the fruits of bhendi and Abutilon

1924.

| Pair No. | Period of recording observations | No. of eggs on <i>Abutilon</i> capsules | No. of eggs on <i>bhendi</i> pods |
|-------------|--|---|---|
| 1 | 27th to 29th August | 54 | 5 |
| 2 | 27th to 31st August | 94 | 1 |
| 3 | 27th August to 6th September | 468 | 33 |
| 4 | 27th August to 3rd September | 287 | 21 |
| 5 | 27th August to 2nd September | 241 | 32 |
| 6 | 27th to 31st August | 164 | 15 |
| | TOTAL . | 1,308 | 107 |

It was further observed in the laboratory that in the absence of any other material the eggs were thrust in the meshes of the cloth covering of the glass cages ; but whenever coarse felt was inserted (for feeding, by soaking pieces with sugar solution) the muslin cloth was abandoned and the eggs were laid deep in the hairy meshes of the felt.

The same tendency has been noticed in the cotton fields, where hairy portions, such as tender shoots, stalks of flower-buds and bolls, and the petioles of leaves are selected for oviposition. The results of the examination of five plants weekly during November and December 1926, and January 1927 for position of eggs on the cotton plant are tabulated below.

TABLE VII

Positions on the cotton plants where the eggs are laid by the Spotted Boll-worms

(20 plants examined in each case)

| Positions | November | December | January | Total |
|---|----------|----------|---------|-------|
| Vegetative shoots | 41 | 41 | 9 | 91 |
| Internodes of branches | 12 | 13 | 5 | 30 |
| Leaf petioles | 5 | 15 | 6 | 26 |
| Stalks of flower-buds | 11 | 18 | 13 | 42 |
| Stalks of bolls | .. | 5 | 10 | 15 |
| Body of flower-buds and bolls | .. | 2 | 1 | 3 |
| TOTAL ON 20 PLANTS | 69 | 94 | 44 | 207 |

INCUBATION PERIOD OF EGGS

The incubation period of the eggs of the Spotted Boll-worms consists of four days almost throughout the year except during the cold months of December and January, when it occupies about 5 to 7 days.

Just before the tiny caterpillar emerges from the egg, a little movement of a black spot can be observed inside the shell, and within a few minutes the larva comes out by knocking off the micropyle, or sometimes by making an irregular hole on the side. The empty white shell is left behind, and many can be found sticking to the stalks of the flower-buds and bolls on the cotton plants for many days.

LIFE OF THE CATERPILLARS

The term boll-worms is partly a misnomer because the larvae of the Spotted Boll-worms do not restrict themselves to bolls only but feed on the tender vegetative shoots and the flower-buds of cotton plants for a considerable period of the season.

It is an important fact to remember that the tiny larvae, as they emerge from eggs, do not begin to feed directly on the nearest available food. They wander about for a few hours before they select their food and begin to bore into it. The rest of the feeding is done from within, either in tender shoots, flower-buds or bolls. The larva does not stay in any one of these forms during all the period of its life and generally it leaves the form before all the available food in it is exhausted. It feeds for a short period in one part of the plant and then it migrates to another. It is this habit which is responsible for the destruction of many more forms than would actually be required for nourishing the caterpillars. It is very common to find a large number of flower-buds and bolls which have been left by the insects after destroying only small portions of these forms.

The larval period was determined by rearing about a dozen larvae every month. Each caterpillar, as soon as it emerged out of the egg, was kept in a separate glass tube for observation. When these larvae pupated, the pupae were further kept under observation for noting the pupal period.

December, January and a part of February are generally the coldest months in this tract, and during this part of the season the larval life extends to about 14 to 16 days. It diminishes to only 9 days during the month of October. During the remaining part of the year it varies between 10 and 12 days. (Table IX, Page 13.)

The kind of food available for the larvae also has some influence on the length of their life. Observations during October 1924, showed, that larvae which were fed on *bhendi* pods could pupate within 9 days, whereas those which were given only tender vegetative shoots of cotton required 14 days before they could pupate. Similarly, further trials during November and December 1924, showed, that the larvae which were fed on *bhendi* pods pupated after feeding for 11 days in November and 14 days in December, whereas the larvae which were fed only on flower-buds of cotton, required 13 and 15 days respectively during the above two months.

PUPAL PERIOD

The caterpillars when they are fully fed descend the plant and invariably go to the soil for pupation in this tract. Very rare cases are observed when the larvae pupate on the plants, either in the monsoon when the soil is wet or during the hottest part of the summer when the heat of the surface soil is unbearable.

The larvae generally commence their search for a suitable place for pupation between 9 A.M. and mid-day. They move through the cracks and crevices until they find a suitable place for spinning their silken cocoons. These are generally attached to the clods or are sometimes found in small crevices (Pl. I, Fig. 1). In cases where the pupae are found in finely pulverised surface soil, small particles of soil are attached all over the cocoon.

In normal field conditions the pupae can be traced to a maximum depth of about 10 to 12 inches below the surface of the soil,



FIG. 1. *Earias* pupae in the soil.




FIG. 2. A cotton shoot damaged by a caterpillar of the spotted boll-worms.

TABLE VIII

Depth at which pupae are found in normal field plots

| | DEPTH IN INCHES | | | | | | | | | | | Total | |
|-------------------------------------|-----------------|------|------|------|------|------|------|------|------|-------|--------|-------|--------|
| | 0-1. | 1-2. | 2-3. | 3-4. | 4-5. | 5-6. | 6-7. | 7-8. | 8-9. | 9-10. | 10-11. | | 11-12. |
| No. of pupae traced during 1928-29. | 4 | 10 | 5 | 2 | 1 | 1 | 1 | 1 | . | . | 1 | . | 20 |
| No. of pupae traced during 1929-30. | 8 | 5 | 9 | 7 | .. | .. | 4 | 5 | .. | 2 | .. | 1 | 41 |

The larvae make their way to these depths through clods and crevices for pupation, and the moths, when they emerge from the pupae, have, therefore, to escape through similar interstices between clods and the loose particles of the soil.

The pupal period varies between 8 and 14 days. The minimum period is observed during April to June and the maximum period during January. (Table IX)

TOTAL PERIOD OF PRE-IMAGINAL LIFE OF THE SPOTTED BOLL-WORMS.

The period of each stage of life of these insects has already been mentioned and the total length of their pre-imaginal life during the different months of the year, from the time the eggs are laid till the emergence of the moths, is summarised in the following table from the information which has already been presented.

TABLE IX

Life of Spotted Boll-worms

| Month. | Incubation period in days | Larval period in days | Pupal period in days | Total life in days |
|-----------------------------|---------------------------|-----------------------|----------------------|--------------------|
| December 1923 | 5 | 14 | 13 | 32 |
| January 1924 | 5 | 16 | 14 | 35 |
| February 1924 | 5 | 11 | 10 | 26 |
| April 1924 | 4 | 10 | 10 | 24 |
| 1st fortnight April 1924 | 4 | 12 | 8 | 24 |
| 2nd fortnight May 1924 | 4 | 11 | 8 | 23 |
| June 1924 | 4 | 11 | 8 | 23 |
| July 1924 | 4 | 10 | 10 | 24 |
| August 1924 | 4 | 10 | 9 | 23 |
| September 1924 | 4 | 10 | 8 | 22 |
| October 1924 | 4 | 9 | 9 | 22 |
| November 1924 | 4 | 12 | 12 | 28 |
| 1st fortnight November 1924 | 4 | 11 | 13 | 28 |
| 2nd fortnight December 1924 | 5 | 14 | 13 | 32 |

It is seen that the shortest period of pre-imaginal life is 22 days in October, the longest period being 35 days during January.

DISTRIBUTION OF THE SPECIES OF SPOTTED BOLL-WORMS

Earias insulana Boisd. and *Earias fabia* Stoll. are the two species of Spotted Boll-worms which damage the cotton crop in Gujarat. Both *insulana* and *fabia* are present throughout the year in the cotton fields but it is *fabia* which always predominates. The proportion of *insulana* is generally the highest during the month of November.

TABLE X

Proportion of Earias fabia and Earias insulana from the larvae collected from the cotton fields at Surat

| Month | <i>Earias fabia</i> | <i>Earias insulana</i> | Total | Percentage of <i>fabia</i> | Percentage of <i>insulana</i> |
|--------------------------|---------------------|------------------------|-------|----------------------------|-------------------------------|
| June 1928 | 44 | 28 | 72 | 61 | 39 |
| July 1928 | 12 | 2 | 14 | 86 | 14 |
| August 1928 | 24 | 4 | 28 | 86 | 14 |
| September 1928 | 28 | 4 | 32 | 88 | 12 |
| October 1928 | 30 | 14 | 44 | 68 | 32 |
| November 1928 | 30 | 30 | 60 | 50 | 50 |
| December 1928 | 68 | 30 | 98 | 69 | 31 |
| January 1929 | 25 | 8 | 33 | 76 | 24 |
| February 1929 | 31 | 7 | 38 | 82 | 18 |
| March 1929 | 28 | 4 | 32 | 88 | 12 |
| April 1929 | 36 | 13 | 49 | 73 | 27 |
| May 1929 | 56 | 18 | 74 | 76 | 24 |

The predominance of *Earias fabia* can also be seen from the classification of moths which emerged in the laboratory during 1923-24 from the larvae collected from the cotton fields.

TABLE XI

| Period of observation | Total moths emerged | <i>E. fabia</i> | <i>E. insulana</i> | Percentage of <i>fabia</i> | Percentage of <i>insulana</i> |
|---|---------------------|-----------------|--------------------|----------------------------|-------------------------------|
| 19th November 1923 to 1st January 1924 | 130 | 63 | 67 | 48 | 52 |
| 17th December 1923 to 30th January 1924 | 142 | 72 | 70 | 51 | 49 |
| 24th January 1924 to 27th February 1924 | 67 | 39 | 28 | 58 | 42 |
| 23rd February 1924 to 19th March 1924 | 155 | 106 | 49 | 68 | 32 |
| 14th March 1924 to 5th May 1924 . . . | 173 | 138 | 35 | 80 | 20 |
| 10th April 1924 to 15th May 1924 . . . | 244 | 184 | 60 | 75 | 25 |
| 10th May 1924 to 25th June 1924 . . . | 390 | 265 | 125 | 68 | 32 |

The *bhendi* plants (*Hibiscus esculentus*) are also attacked by the Spotted Boll-worms, but it is *E. fabia* which is mainly responsible for this damage because the proportion of *insulana* found on *bhendi* is very low.

TABLE XII

Larvae collected from Bhendi

| Months | <i>Earias fabia</i> | <i>Earias insulana</i> | Total | Percentage <i>fabia</i> | Percentage <i>insulana</i> |
|--------------------|---------------------|------------------------|-------|-------------------------|----------------------------|
| April 1928 | 286 | 8 | 294 | 97 | 3 |
| May 1928 | 69 | 8 | 77 | 90 | 10 |
| June 1928 | 33 | 1 | 34 | 97 | 3 |

On *Abutilon* species a good proportion of *insulana* was always noticed.

ARE THE SPOTTED BOLL-WORMS ACTIVE THROUGHOUT THE YEAR ?

There are many insects which remain inactive in one stage of life or another for a part of the year. This period often happens to be the cold season, or it coincides with the period of scarcity of food supply for the particular insect. The study of this factor in the life-history of the Spotted Boll-worms was considered to be an important point and attempts were therefore made to ascertain if these insects had, in this tract, any such period of hibernation or aestivation.

Experience at Surat, while carrying out various trials, showed that these insects could be collected in the fields and could also be bred in the laboratory throughout the year. All the four stages of this pest could be easily found in the cotton fields even during severe winter, or during the summer months. The question had, however, to be examined of the possibility of their aestivating as pupae in the field-soil during summer, after the *kupas* was harvested. Observations were, therefore, undertaken to clear this issue. A large number of full-grown larvae were liberated in small selected areas in a cotton field during the summer months, with a view to find out if these worms would pupate in the soil and remain in aestivation till the commencement of the next season, or would emerge as moths after the normal pupal period. The details of these trials are given below :—

A small area (6 ft. by 6 ft.) was selected in the cotton fields and 1,076 full-grown larvae were liberated in this small space in the last week of March 1928. Before liberating these larvae, zinc sheets were buried around this block of soil to a depth of 3 feet, so that the larvae may not escape from the area under observation through cracks in the soils. The emergence of moths was noted from this area by placing a cloth cage over it.

It was found that 135 moths emerged from this area between the 10th and 15th day, after the larvae were liberated and no emergence was noticed after this period. At the beginning of June, the enclosed soil was excavated and was carefully examined for the cocoons. 497 cocoons were recovered but none of them contained a live pupa.

It is seen that though we liberated 1,076 larvae, we could trace only 497 cocoons after a period of about two months. Out of the remaining 579 larvae, a few must have failed to pupate and the rest must have been completely destroyed after they had pupated in the soil. It is further interesting to note that even out of these 497 cocoons, we could secure only 135 moths, and the remaining pupae were damaged by predators before the moths could emerge.

In addition to this trial, two zinc cylinders, two and a half feet in diameter and open at both ends, were buried in the field soil to a depth of 3 ft., and 200 full-grown larvae were liberated in each of them during the last week of March. The emergence from each of them was recorded by caging the enclosed area. It was found that 75 and 84 moths emerged from the area enclosed by the two cylinders respectively within 11 days after the larvae were liberated. The soil from the first cylinder was examined after 4 weeks and that from the second was examined after 8 weeks from the commencement of the experiment. In both the cases, no live pupa could be traced at the time of soil examination.

Later on, in the last week of May, another group of 200 larvae was liberated in a small space enclosed in a cylinder as mentioned above. Within 10 days, 111 moths emerged from this area and no moth was found after that period. The soil was examined in the last week of June and again no live pupa could be secured.

The observations indicated that there was no possibility of the Spotted Boll-worms aestivating as pupae in the soils, in this tract, during the summer months which intervene between the old and the new crop of cotton.

So far as the cold season is concerned, it was found that it prolonged the life of these worms, and the total period of life extended to about 35 days during the cold weather but no hibernation as such was observed in this locality.

The effect of continuous cold on the pupal period was studied under artificial conditions in one of the experiments. Nine sets of 10 fresh pupae in each case were prepared and they were placed in 9 different compartments of a multiple temperature incubator (Appendix III) where the temperatures ranged from 58° to 125° F.

It will be seen from the next table that the period of pupation was considerably prolonged at lower temperatures, and it covered as many as 24 days in the case of the pupae, which were exposed to a temperature varying between 66° and 70° F. In the adjoining compartment, the temperature varied between 58° and 64° F. and the result was that the pupae from this lot did not hatch out for 30 days. They were removed from there, after this period, and were kept at the room temperature, where the moths emerged normally after four days.

TABLE XIII

Emergence of moths from pupae exposed to varying temperatures

| Days after | Number of moths that emerged from different chambers with varying ranges of temperature. (In Fahrenheit.) | | | | | | | | | | | |
|-------------------|---|-------------------|-------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|------------------------|
| | 122 to 125. | 110 to 112. | 100 to 102. | 96 to 98. | 90 to 93. | 89 to 91. | 84 to 86. | 80 to 83. | 79 to 81. | 73 to 75. | 66 to 70. | 58 to 61. |
| 7 | All pupae dead and dried up. | | | .. | .. | 3 | .. | . | .. | .. | .. | .. |
| 8 | .. | .. | .. | 3 | 7 | 1 | 3 | .. | .. | .. | .. | .. |
| 9 | .. | .. | .. | 4 | 2 | 5 | 4 | 4 | .. | .. | .. | .. |
| 10 | .. | .. | .. | .. | .. | 1 | 2 | 5 | .. | .. | .. | .. |
| 11 | .. | .. | .. | .. | .. | .. | .. | 1 | 3 | .. | .. | .. |
| 12 | .. | .. | .. | .. | .. | .. | .. | . | 3 | .. | .. | .. |
| 13 | .. | .. | .. | .. | .. | .. | .. | . | 2 | 1 | .. | .. |
| 14 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 5 | .. | .. |
| 15 | .. | .. | .. | .. | .. | .. | .. | . | .. | 2 | .. | .. |
| 16 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | .. | .. |
| 17 | .. | .. | .. | .. | .. | .. | . | .. | .. | . | . | .. |
| 18 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 19 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 20 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 21 | .. | .. | .. | .. | .. | .. | .. | . | .. | .. | .. | .. |
| 22 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | 2 | .. |
| 23 | .. | .. | . | . | .. | .. | .. | . | .. | .. | 4 | .. |
| 24 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | .. |
| 25 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | . | .. |
| 26 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 27 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | . | .. |
| TOTAL EMERGENCE . | .. | .. | .. | 7 | 9 | 10 | 9 | 10 | 8 | 9 | 10 | Nd (all live pupae) |

The above information will be of considerable interest for tracts like Sind and the Punjab because it indicates the possibilities of the Spotted Boll-worms passing a very long period, in their pupal stage, during the months of severe cold of these provinces.

Moths which emerged in the laboratory during the months of June, July and August were kept in glass jars and were fed regularly on honey in order to find out if the Spotted Boll-worms would be able to pass the period between the two cotton growing seasons in the adult stage.

TABLE XIV

Life of moths kept under observation during June, July and August 1928

| Month | No. of moths under observation | Average life in days | Maximum life in days |
|------------------|--------------------------------|----------------------|----------------------|
| June | 41 | 8 | 25 |
| July | 89 | 9.5 | 22 |
| August | 51 | 8.6 | 24 |

It was found that the average life of these moths was not more than 10 days and that the maximum period for which a moth lived did not exceed 25 days in any case.

All the observations, therefore, indicated that the Spotted Boll-worms were active throughout the year in this tract and that no evidence was available of their passing any time of the year in hibernation or aestivation.

FLIGHT RANGE OF MOTHS

Knowledge of the flight range of these moths would be of considerable value for estimating the rate at which fresh infection of this pest would spread. Accurate determination of flight range would probably be possible only in tracts which are free from Spotted Boll-worms. Such a suitable tract was not, however, available for us, and therefore only a few attempts were made to study the question of flight dispersal within narrow limits.

The moths when disturbed during the day time, only flit about a little and settle on plants within a few yards. During the night, however, they travel more extensively.

During one of the weeks of March 1928, 1,875 moths were coloured red and were liberated in the evening in the midst of a big cotton area. Next morning there was no trace of a single moth, near about the spot where they were let out. During the subsequent two or three days, a careful search was made for locating these moths

in the surrounding areas, with a radius of about 500 yards. Only five coloured moths could be secured after this search and the maximum distance at which one of these moths was caught was 300 yards away from the spot where they were originally liberated. It is, therefore, concluded that the moths must have spread out very rapidly beyond the distance of 500 yards where it was not possible to carry on a thorough search.

DAMAGE TO THE COTTON CROP BY THE SPOTTED BOLL-WORMS

SEASON AND PLANT GROWTH

Before describing the damage caused by the Spotted Boll-worms to the cotton crop, it is desirable to give a brief general account of the environmental conditions of this crop in this tract. (A detailed account of these conditions is given by Maganlal L. Patel [1924] in his memoirs on Gujarat cottons.)

In South Gujarat, the monsoon generally breaks out some time during the latter part of June, and the rainy season continues almost till the end of October. The average rainfall at Surat is 40 inches, and a large part of it is received during the months of July and August. Cotton is sown soon after the first few showers, at the end of June or early in July. The cultivator generally drills his cotton, but dibbling the seed at regular distances is being adopted by some of the cultivators. Desai and Naik [1925] have reported successful results of growing the crop on ridges, which help it to stand the water-logging of the soils during the monsoon at Surat.

In a normal season, cotton seedlings, about four to six inches in height, are seen all over the cotton fields in the month of July, but they do not make any rapid progress in growth for nearly 6 to 8 weeks in the clayey soils of this tract, due to the excessive moisture in the soil and the continuously overcast skies. With the advent of drier and more open conditions, some time during the months of September or October, vigorous growth commences and is followed by rapid formation and development of the flower-buds. Flowers begin to appear in November, and boll development continues during the months of December and January.

The opening of the boll takes place during the months of March and April, and consequently it is during these months that the pickings of *kapas* are made.

Most of the observations at Surat were on plants of the strain called 1027 A. L. F., which is being extensively cultivated in this tract, south of the Nurbuda. The seed was dibbled on small ridges which were prepared before the beginning of the monsoon.

DAMAGE TO THE VEGETATIVE SHOOTS

As soon as the vigorous growth of the cotton crop commences, after the cessation of the monsoon, the Spotted Boll-worms make their appearance in the cotton fields.

They either destroy the tiny buds of the growing shoots or bore into the succulent internodes about 3 or 4 inches below the top of the shoot. When a larva bores into an internode, the portion of the shoot above the bore generally withers and dries up. The appearance of these withered shoots is so characteristic that they can easily be observed from a distance (Pl. I, Fig. 2). In a few cases, however, when the injury is not very severe, the shoot does not dry and the injury caused by the bore is healed up. In the case of the shoots which survive the attack, the scars are visible almost throughout the season. When only the growing vegetative bud is devoured by these worms, the injury is not easily perceptible. Whether the growing bud is destroyed or the shoot is bored into, the ultimate effect, in most cases, is that the further elongation of the branch is stopped and fresh growth is stimulated at the lower nodes.

It must be mentioned here that when the above damage is in progress, the shoot-roller (*Phycita infusella* Meyr.) is also active on the shoots of the cotton plants (Appendix V, Tables II to V). These larvae bind together the top leaves of the branches and feed on the tender leaves from within. The bunch of tied leaves withers and dries up, and consequently in these cases also the further growth of the branches is prevented.

During the year 1925-26, two hundred plants were kept under observation from the beginning of the cotton-growing season for recording the injury caused by the Spotted Boll-worms and the shoot-rollers. Withering of the shoot completely, or merely the destruction of the growing bud, were both reckoned as injury. Ultimately full records for 181 plants were available. It was found that the shoots of the main axis of all these plants were damaged by these insects, 38 per cent of them being destroyed by the Spotted Boll-worms and remainder by the shoot-rollers.

Similar observations were recorded on a group of 200 plants during the year 1926-27 and the progress of attack by the Spotted Boll-worms on the shoots of the main stems can be seen from the next table.

TABLE XV

| Week ending | Average height of the plants in cms. | No. of shoots damaged by boll-worms |
|-------------------------------|--------------------------------------|-------------------------------------|
| 25th September 1926 | 19 | 13 |
| 2nd October 1926 | 23 | 18 |
| 9th October 1926 | 28 | 11 |
| 16th October 1926 | 35 | 48 |
| 23rd October 1926 | 40 | 18 |
| 30th October 1926 | 44 | 9 |
| 6th November 1926 | 47 | 10 |
| 13th November 1926 | 49 | 1 |
| TOTAL | | 128 |

It is seen that the attack on the shoots commenced when the plants were about 19 cms. in height and it continued till the average height of the plants was 49 cms. The maximum number of damaged shoots was in the week ending 16th October, when the average height of the plants was 35 cms.

The boll-worms do not restrict themselves to the shoots of the main stems, but the succulent tops of the side branches are also bored and the growing vegetative buds are destroyed, in several positions on the plants. By the middle of November, however, the internodes of the shoots begin to be woody and are no longer inviting to the worms. The withering of the shoots therefore disappears, but the injury to the growing tips does continue thereafter. A few instances of tiny larvae feeding on the vegetative buds can be traced even during the months of December and January. (Appendix I, Table IX.) An idea of the incidence of the larval population on the vegetative shoots during 6 successive seasons can be obtained from Appendix I, Tables VIII to XIII. It is observed that during most of these years, the boll-worms appeared on the cotton plants some time during the month of September, and a few continued to feed on the shoots at least till the end of November. This continuous mutilation of the growing points, during the period when the main scaffolding of the plant is being built up, results in an irregular growth of the branches, and ultimately the plant appears to be very much more bushy than it would otherwise do.

DAMAGE TO FLOWER-BUDS

With the vigorous growth of the plants, the formation of the flower-buds also commences, and as the developing buds begin to increase in numbers, the Spotted Boll-worms gradually divert their attention from the shoots to the flower-buds. The larvae get into the flower-buds by making a hole and feed on the anthers and ovaries from within. Frequently the buds are abandoned by these larvae after causing only slight injury. It must, however, be remembered that the entrance hole bored by the tiniest larvae invariably causes the shedding of even the biggest bud. Damage by the Spotted Boll-worms is therefore one of the important factors which are responsible for the shedding of flower-buds. Careful observations were maintained on a small number of plants during 3 cotton growing seasons, commencing from 1925-26, for studying this damage to the flower-buds. Weekly counts were maintained from these plants, recording the periodical flower-bud formation, flower opening and the development of mature bolls. The flower-buds and bolls which dropped down from these plants were collected, and were examined for determining the proportion of these forms which showed damage caused by the Spotted Boll-worms. (Appendix I, Tables IV, V and VI.)

The flower-bud formation commenced during the month of September in 1925 and 1927, whereas in 1926, it began early in October. The delay in the commencement of the formation of the flower-buds during 1926 was due to the growth of the plant being retarded by the excessive moisture in the soil caused by heavy rains

(Appendix I, Table VII). It was noticed that in all the three years, shedding of flower-buds due to the Spotted Boll-worms continued throughout the season. The earliest shedding of the flower-buds, however, consisted of a large proportion of tiny buds which generally fail to develop. Within a period of about 6 to 8 weeks after the commencement of flower-bud formation, the proportion of the buds attacked by boll-worms increased, and over 50 per cent of the shed buds indicated damage caused by the *Earias* larvae.

During 1925-26, the flower-buds suffered heavily from this damage, almost from the beginning of October till the end of November. In 1927-28, similar damage was observed from the beginning of November till the middle of December. The year 1926-27 turned out to be the worst boll-worm year, and a considerable amount of damage to the flower-buds occurred during that year from the middle of November 1926 to the middle of January 1927.

TABLE XVI


Proportion of the flower-buds damaged by the Spotted Boll-worms

| Year | No. of plants observed | Total No of flower-buds formed per plant | Total No of flowers opened per plant | Percentage of flowers opened to buds formed | Total buds shed per plant | Shed buds examined per plant | Shed buds damaged by boll-worms per plant | Percentage of shed buds due to boll-worms to the total shed |
|---------------|------------------------|--|--------------------------------------|---|---------------------------|------------------------------|---|---|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1925-26 . . . | 49 | 252 | 74 | 29 | 179 | 163 | 79 | 48 |
| 1926-27 . . . | 20 | 297 | 59 | 20 | 238 | 222 | 114 | 51 |
| 1927-28 . . . | 16 | 411 | 96 | 23 | 314 | 244 | 82 | 34 |


Out of the total flower-buds formed, from 20 to 29 per cent opened into flowers and the remaining 71 to 80 per cent shed as buds during the three seasons under consideration. It was further noted that out of this large proportion of flower-buds which dropped away from the plants 34 to 51 per cent (roughly one-third to one half) were injured by the boll-worms (Plate II).

DAMAGE TO FLOWERS AND BOLLS


The shedding of flower-buds due to the boll-worms delayed the period of flower-formation. In fact, intense flower opening was observed during the above three



1925 - 1926 .






1926 - 1927 .



1927 - 1928 .

Showing periodical flower-bud formation, total bud-shedding, and the bud-shedding due to boll-worms from cotton plants at Surat, during three seasons.

PLATE III.



Showing periodical flower opening, boll-shedding and the boll-shedding due to boll-worms from cotton plants at Surat, during three seasons

seasons only after the decline of the shedding of flower-buds due to these insects, (Appendix I, Tables IV, V and VI). This fact was particularly brought out during the year 1926-27 when not only the flower opening commenced very late, but it continued to be very tardy, and a slight intensity of flowering could only be noticed as late as after the end of January.

The Spotted Boll-worms begin to concentrate on the flowers and young bolls as soon as the flower opening begins. When a flower is attacked, the larva, in most cases, enters the ovary and destroys it. Consequently the flower withers and sheds within 3 to 5 days. The developing bolls also are attacked at all stages of their growth. Young bolls, before they are about half grown, drop away from the plants when they are damaged by the Spotted Boll-worms, but the bigger bolls continue to stick to the plant, even when they are injured. These bolls usually yield damaged *kapas*.

It has already been mentioned that the flower-opening is delayed due to the shedding of flower-buds attacked by these larvae. It is further observed that the intensity of attack on the flowers and young bolls at the commencement of the flower-opening delays the boll formation and ultimately the opening of bolls.

It will be interesting to scrutinise the details of the damage to the flowers and young bolls. During 1925-26 the flowering commenced in the week ending on 17th October, and during the first four weeks of the flowering, 76 flowers opened on the 25 plants under observation. Out of these flowers, only 14 bolls developed, and the remaining 62 flowers dropped away, of which 54 (i.e., 87 per cent) were injured by boll-worms. Thereafter, the intensity of the damage of these insects declined and the major crop of bolls was obtained from flowers which opened from the middle of November to the middle of December.

Similarly during 1927-28, fifty-six per cent of the flowers which opened in the first six weeks of the flowering dropped away, and 67 per cent of the sheds were damaged by the boll-worms. The major part of the crop of bolls during that season was gathered from flowers which opened from the middle of December till the 3rd week of January.

The performance of the plants during 1926-27 was, however, particularly interesting. During that year the flower opening commenced at the end of October, but 70 per cent of the flowers which opened till the end of December were destroyed by the boll-worms, and 79 per cent of the final crop was built up from the last flowers which opened after the beginning of January. It was also noticed that due to this continuous destruction of the flowers and young bolls, the flowering was very protracted and continued from the end of October till the end of February (Plate III).

These observations have, therefore, shown very clearly that the success of bolls from the earliest flowers is considerably reduced, due to the shedding of bolls caused by the Spotted Boll-worms, and that the flowering period is prolonged if the severity of attack on the flowers and young bolls continues for a long period.


TABLE XVII

| Year | Plants observed | Total flowers opened per plant | Bolls shed per plant | Bolls shed due to boll-worms per plant | Percentage of the bolls shed due to boll-worms out of the total shed | Percentage of bolls shed due to boll-worms to the flowers opened |
|-------------|-----------------|--------------------------------|----------------------|--|--|--|
| 1925-26 . . | 49 | 74 | 46 | 10 | 22 | 13.5 |
| 1926-27 . . | 20 | 59 | 35 | 24 | 69 | 41 |
| 1927-28 . . | 16 | 96 | 59 | 12 | 20 | 12.5 |

Out of the 3 seasons under consideration it was during 1926-27 that the largest proportion of the shed bolls showed injury caused by the Spotted Boll-worms, and that 41 per cent of the total flowers were destroyed by the boll-worms in that season. During the other two years this damage was restricted to only 13.5 per cent and 12.5 per cent of the total flowers.

It has already been mentioned that some of the developed bolls are also attacked by these worms and that such bolls yield damaged *kapas*. All the dirty *kapas* that we find at the time of pickings, however, is not due to the injury of the Spotted Boll-worms alone, because the Pink Boll-worm (*Platyedra gossypiella* Saund.) is responsible for a greater part of this damage to the developed bolls. The Pink Boll-worm appears in the fields just when the bolls are developing and rapidly increases in numbers. (Appendix VI, Tables VII to XII.)

For ascertaining the total damage to the bolls caused by both the pests, more than 100 plants were kept under observation during 1928-29, 1929-30 and 1930-31 and all the bolls from these plants were carefully picked and the damaged and the sound locks of these bolls were separated out into different classes as mentioned below :—



The results of this classification of the locks during the years mentioned above are summarised in the next table.

TABLE XVIII
(Average per plant)

| Year | No. of plants observed | Total locks picked | LOCKS YIELDING SALEABLE <i>kapas</i> | | | USELESS LOCKS WITH UNSALEABLE <i>kapas</i> | |
|-------------|------------------------|--------------------|--------------------------------------|-------------------|---------------------|--|-----------------------------------|
| | | | Sound locks | Damaged locks | | Damaged by boll-worms | Not developed due to other causes |
| | | | | Due to boll-worms | Due to other causes | | |
| 1928-29 . . | 104 | 202.4 | 144.2 | 38.3 | 8.8 | 6.9 | 4.2 |
| 1929-30 . . | 112 | 170.6 | 156.2 | 9.0 | 2.3 | 2.5 | 0.6 |
| 1930-31 . . | 116 | 211.8 | 190.6 | 17.7 | 1.3 | 2.1 | 0.1 |

The sound and the damaged locks of the saleable *kapas* were collected and weighed separately, and an idea of the extent of the damaged *kapas* can be had from the next statement.

TABLE XIX
(Average per plant) Plants spaced 4' by 4'

| Year | No. of plants observed | Total saleable <i>kapas</i> in Grms. | <i>Kapas</i> from sound locks in Grms. | <i>Kapas</i> from damaged locks in Grms. | Percentage damaged <i>kapas</i> |
|-----------------|------------------------|--------------------------------------|--|--|---------------------------------|
| 1928-29 | 104 | 130.6 | 105.5 | 25.1 | 19 |
| 1929-30 | 112 | 113.2 | 107.9 | 5.3 | 4.7 |
| 1930-31 | 116 | 150.1 | 140.7 | 9.4 | 6.3 |

It is seen that the proportion of the damaged *kapas* was very high during 1928-29. It appears that this was mainly due to a severe attack of the Pink Boll worm during that season. (Table XI of Appendix VI.)






It is very difficult for a casual observer to appreciate the damage caused by the Spotted Boll-worms to the cotton crop as the plant apparently looks quite healthy, and caterpillars feeding on the crop hardly attract the attention of an untrained eye. The observations described above will, however, show the continuous damage to which the cotton crop is subjected by this pest from the beginning to the end of the season.

ABUNDANCE OF THE SPOTTED BOLL-WORMS IN THE COTTON FIELDS AT SURAT

An idea of the comparative abundance of a pest can be obtained if we can estimate its population periodically in a given area of the crop, which is damaged by that pest. In order, therefore, to study the increase or decrease of the Spotted Boll-worms in the cotton fields at Surat, the larval population, on a fixed number of plants, was ascertained weekly. Five small plots of cotton were reserved for obtaining plants for examination, and from these plots 5 plants were collected daily by cutting one plant from each plot. A total number of 25 plants was examined every week, except during the commencement of the growing season when a large number of seedlings had to be examined to trace out the population which naturally used to be very meagre at that time. For these observations all the parts of these plants were separated out in the laboratory and all the shoots, flower-buds and bolls were carefully examined to record the larvae which might be feeding in them. The number of empty forms, from which the larvae had escaped after feeding was also noted. Incidentally notes were also maintained about the presence of shoot rollers (*Phycita infusella* Meyr.) and the Pink Boll-worm (*Platyedra gossypiella* Saund.). The information about these pests is separately presented (Appendices V and VI).

These observations for the larval population of the Spotted Boll-worms were carried out during 6 successive seasons from 1925 to 1931 and the results are tabulated in 6 separate statements in Appendix I, Tables VIII to XIII.

These statements show that the Spotted Boll-worms make their earliest appearance on the shoots of the cotton plants. They divert their attention to the flower-buds as these begin to appear, and for a few weeks the attack continues, both on shoots and flower-buds. When the boll development begins, a part of the population shifts from flower-buds to bolls (Plate IV). It is interesting to note that when both flower-buds and bolls are present there is proportionately a larger number of Spotted Boll-worms feeding on bolls than on flower-buds. This indicates that they prefer bolls to flower-buds. It must, however, be remembered that they feed on the flower-buds for the longest period in the cotton-growing season.



— Earias larvae on 25 plants. - - - - Earias larvae in flower buds,
 - - - - Earias larvae in shoots. - - - - Earias larvae in bolls

Showing the distribution of the population of *Earias* larvae on different parts of 25 plants during 6 consecutive seasons at Siraz.

TABLE XX

Spotted Boll-worm larvae found in every 100 flower-buds and every 100 green bolls present on 25 plants at the time of weekly examinations

1929-30

| Date | No. of buds | No. of larvae in flower-buds | No. of larvae for every 100 flower-buds | No. of green bolls | Total larvae in bolls | No. of larvae for every 100 bolls |
|------------------------|-------------|------------------------------|---|--------------------|-----------------------|-----------------------------------|
| 23rd October 1929 . . | 1,137 | 58 | 5.1 | 6 | 3 | 50.0 |
| 30th October 1929 . . | 1,667 | 86 | 5.2 | 8 | 2 | 25.0 |
| 8th November 1929 . . | 1,993 | 143 | 7.2 | 5 | 2 | 40.0 |
| 14th November 1929 . . | 2,627 | 167 | 6.4 | 17 | 6 | 35.3 |
| 21st November 1929 . . | 2,529 | 143 | 5.7 | 20 | 7 | 35.0 |
| 28th November 1929 . . | 3,381 | 143 | 4.2 | 32 | 7 | 21.9 |
| 5th December 1929 . . | 3,637 | 153 | 4.2 | 72 | 11 | 15.3 |
| 12th December 1929 . . | 3,829 | 92 | 2.4 | 149 | 15 | 10.0 |
| 19th December 1929 . . | 2,673 | 45 | 1.6 | 338 | 24 | 7.1 |
| 26th December 1929 . . | 2,831 | 30 | 1.1 | 493 | 21 | 4.2 |
| 2nd January 1930 . . | 2,537 | 24 | 0.9 | 866 | 29 | 3.3 |
| 9th January 1930 . . | 1,721 | 15 | 0.8 | 1,350 | 23 | 1.7 |
| 16th January 1930 . . | 940 | 3 | 0.3 | 1,814 | 11 | 0.6 |
| 23rd January 1930 . . | 288 | 2 | 0.7 | 1,468 | 7 | 0.5 |
| 30th January 1930 . . | 67 | .. | .. | 1,102 | 3 | 0.3 |

TABLE XXI

Spotted Boll-worm larvae found in every 100 flower-buds and every 100 green bolls present on 25 cotton plants at the time of weekly examinations at Surat

1930-31

| Date | No. of flower-buds | No. of larvae in flower-buds | No. of larvae for every 100 flower-buds | No. of green bolls | Total larvae in bolls | No. of larvae for every 100 green bolls |
|------------------------|--------------------|------------------------------|---|--------------------|-----------------------|---|
| 24th October 1930 . . | 2,260 | 158 | 7 | 50 | 9 | 18.0 |
| 31st October 1930 . . | 1,977 | 155 | 7.8 | 43 | 14 | 32.6 |
| 7th November 1930 . . | 1,777 | 122 | 6.9 | 51 | 12 | 23.5 |
| 14th November 1930 . . | 2,107 | 123 | 5.8 | 33 | 6 | 18.2 |
| 21st November 1930 . . | 2,688 | 78 | 2.9 | 32 | 9 | 28.1 |
| 28th November 1930 . . | 3,098 | 90 | 2.2 | 75 | 15 | 20.0 |
| 5th December 1930 . . | 5,192 | 132 | 2.5 | 208 | 24 | 11.5 |
| 12th December 1930 . . | 4,224 | 91 | 2.1 | 288 | 28 | 9.7 |
| 19th December 1930 . . | 4,207 | 52 | 1.2 | 868 | 33 | 3.8 |
| 26th December 1930 . . | 3,917 | 33 | 0.8 | 1,707 | 19 | 1.1 |
| 2nd January 1931 . . | 2,198 | 13 | 0.6 | 1,895 | 17 | 0.9 |
| 9th January 1931 . . | 927 | 6 | 0.6 | 2,167 | 17 | 0.8 |
| 16th January 1931 . . | 225 | 1 | 0.4 | 1,692 | 13 | 0.7 |
| 23rd January 1931 . . | 18 | .. | .. | 1,248 | 20 | 1.6 |

A compact statement of the weekly larval population on 25 cotton plants during the above mentioned 6 years is given in Table XXII.

TABLE XXII

Weekly population of the Spotted Boll-worms on 25 cotton plants at Surat
 (Plants spaced 3' × 3' during 1925-26, 1926-27 and 1927-28 and 4' × 3' during 1928-29, 1929-30 and 1930-31.)

| Week ending | 1925-26 | 1926-27 | 1927-28 | 1928-29 | 1929-30 | 1930-31 |
|----------------------|---------|---------|---------|---------|---------|---------|
| 5th September . . . | .. | .. | .. | .. | .. | 1 |
| 12th September . . . | .. | .. | .. | 2 | .. | 1 |
| 19th September . . . | 32 | .. | 8 | .. | 1 | 2 |
| 26th September . . . | 48 | .. | 25 | 1 | 1 | 18 |
| 3rd October . . . | 48 | .. | 24 | 4 | 7 | 13 |
| 10th October . . . | 43 | .. | 22 | 20 | 35 | 28 |
| 17th October . . . | 68 | 3 | 74 | 45 | 70 | 78 |
| 24th October . . . | 87 | 5 | 57 | 56 | 81 | 196 |
| 31st October . . . | 73 | 16 | 61 | 32 | 114 | 220 |
| 7th November . . . | 45 | 19 | 51 | 89 | 176 | 167 |
| 14th November . . . | 45 | 38 | 78 | 86 | 204 | 145 |
| 21st November . . . | 32 | 62 | 46 | 95 | 167 | 90 |
| 28th November . . . | 54 | 91 | 69 | 130 | 159 | 108 |
| 5th December . . . | 47 | 76 | 63 | 115 | 165 | 157 |
| 12th December . . . | 44 | 138 | 42 | 123 | 107 | 122 |
| 19th December . . . | 55 | 123 | 34 | 65 | 69 | 85 |
| 26th December . . . | 51 | 144 | 34 | 37 | 51 | 53 |
| 2nd January . . . | 38 | * | 41 | 40 | 54 | 30 |
| 9th January . . . | 30 | 67 | 38 | 19 | 38 | 23 |
| 16th January . . . | 22 | 86 | 35 | 18 | 14 | 14 |
| 23rd January . . . | 25 | 47 | 18 | 20 | 9 | 20 |
| 30th January . . . | 14 | 47 | 12 | 25 | 3 | 28 |
| 6th February . . . | .. | 44 | 11 | 19 | 18 | 37 |
| 13th February . . . | .. | 15 | 4 | 46 | 25 | 23 |
| 20th February . . . | .. | 21 | 2 | 54 | 28 | 17 |
| 27th February . . . | .. | 9 | 6 | 56 | .. | 4 |

*Not examined.

The time of the earliest rise of the population of these larvae in the cotton fields has varied considerably during these 6 years. In 1925-26 this rise was found to be the earliest, while during the next season it was abnormally delayed till the middle of November. In 1927-28 again the boll-worms began to increase in numbers before the end of September, whereas the commencement of similar rise was noticed by about the second week of October during the remaining 3 seasons.


The intensity and distribution of rainfall were important factors in determining the time of the rapid increase of the Spotted Boll-worms, because rain had a depressing effect on their multiplication, and therefore the rise of the population could not begin so long as the rains continued (Plate V). It was also noticed that the population rapidly declined if heavy showers of rain were received even during the later part of the cotton growing season.

Thus on 12th and 13th of November 1927, the rainfall was 2.88 inches. This was followed by a sudden drop in the larval population on the 25 plants from 75 to 44 during the next week. Similarly the population rapidly declined after a storm which occurred on the 29th of October 1930, and which was accompanied by 0.90 inches of rain.


TABLE XXIII
Rainfall at Surat from 1925 to 1930
(In inches)

| Month | Fortnight | 1925 | 1926 | 1927 | 1928 | 1929 | 1930 |
|-----------|-----------|-------|-------|-------|-------|-------|-------|
| June | First | 10.78 | .. | 0.45 | 2.91 | 0.78 | 0.11 |
| | Second | 6.55 | 0.97 | 9.51 | .. | 8.26 | 16.32 |
| July | First | 1.41 | 15.68 | 5.93 | 8.59 | 33.78 | 10.41 |
| | Second | 1.45 | 13.28 | 6.41 | 3.27 | 0.54 | 5.54 |
| August | First | 3.82 | 11.68 | 1.50 | 2.11 | 1.80 | 2.19 |
| | Second | 0.33 | 8.09 | 1.99 | 1.85 | 1.45 | 0.26 |
| September | First | 1.04 | 6.65 | 0.99 | 0.38 | .. | 5.34 |
| | Second | .. | 6.60 | 3.25 | 5.90 | 0.45 | 0.06 |
| October | First | .. | .. | 2.15 | 0.49 | 0.32 | .. |
| | Second | .. | .. | .. | 0.38 | .. | 0.95 |
| November | First | 0.06 | .. | 2.88 | .. | .. | .. |
| | Second | .. | .. | .. | 0.29 | .. | .. |
| TOTAL | | 25.44 | 62.95 | 35.08 | 26.17 | 47.38 | 41.18 |

PLATE V.



Showing incidence of larval-population of *Earias* larvae on 25 cotton plants and weekly rainfall during six consecutive seasons at Surat.



Showing the mean night temperatures and the periodical population of *Earias* larvae during four consecutive seasons at Surat

The year 1925 had a very peculiar type of monsoon. During that year Surat had, from June to November, a total of 25.44 inches of rain, of which 17.33 inches were received in the month of June. The remaining 8 inches were spread over the next 3 months. The Spotted Boll-worms, therefore, got the earliest opportunity of increasing in numbers during that season.

The next season had the heaviest rainfall, and all the 3 months of July, August and September continued to be very wet. The result was that the rapid multiplication of this pest during that year was delayed till the middle of November.

The year 1927 was not quite as dry as 1925 but there were no heavy rains during August and September, and therefore a good beginning of boll-worm population was noticed in the last week of September.

The boll-worms would have increased in numbers before the end of September in 1928, because the period from the beginning of August till the middle of September was not very wet. The heavy showers received during the 2nd fortnight of September, however, delayed their rise till the 2nd week of October.

During the next two seasons the heavy rains in June and July followed by small showers during the next few weeks kept off the serious rise of the pest till the middle of October.

The various indications mentioned above, have, therefore, shown that a heavy rainfall is always effective in checking the rapid increase of the Spotted Boll-worms.

These observations have further shown that the population of the Spotted Boll-worms declines rapidly every year at the end of December or some time in January. Scarcity of food does not seem to be a reason for this, because a large number of flower-buds and bolls can still be found on the cotton plants, when the population of these insects begins to fall. It appears that this decline is partly due to the activities of the parasites of these worms and partly to the effect of climatic conditions.

The following 3 parasites are generally active during the months of November and December, and are responsible to a great extent in controlling the Spotted Boll-worms.

- (1) *Microbracon lefroyi* D. and G.
- (2) *Actia aegyptia* Vill.
- (3) *Rhogas testaceus* Grav.

The cold season generally begins just about the time when the population of the Spotted Boll-worms declines. The cold season prolongs the total life of these worms, and consequently their rate of multiplication is decreased. It is further observed that the period of the decline of these insects and the fall in the night temperature generally coincide (Plate VI). It has been remarked earlier that the capacity of oviposition of the moths declines appreciably with the commencement of the cold weather (Page 9).

In short the Spotted Boll-worms begin to increase rapidly in the cotton fields soon after the close of the monsoon. They feed on the shoots, flower-buds and bolls as they begin to appear in succession. The cotton plant at Surat shows a very great

capacity of getting over all this damage. When the growing shoots are destroyed, fresh growth commences at other points; when the developing flower-buds are damaged, the younger ones begin to develop and open into flowers; and when young bolls drop down due to the injury caused by the caterpillars, the later flowers succeed into bolls (Appendix VIII). The plant cannot, however, be expected to stand this mutilation indefinitely and therefore it is fortunate that ultimately the population of the Spotted Boll-worms declines at the end of December or early in January and allows the cotton plants to complete their performance.

BEHAVIOUR OF THE COTTON PLANT WHEN IT IS GROWN FREE FROM BOLL-WORMS AND ESTIMATION OF THE IMPROVEMENT IN THE YIELD OF *KAPAS*.

Knowledge of the number of shoots destroyed or the amount of shedding of flower-buds and bolls, or even the exact counts of the number of larvae feeding on the plants every week, help but little to estimate the increased yield of *kapas* which should be available, if the boll-worms were absent from the cotton fields. As the economics of control depend upon this, it was necessary to resort to a direct method of studying the plants by growing them free from boll-worm attack.

PLANT CAGES OF WOODEN FRAMES COVERED BY WIRE GAUZE.

To exclude the worms from the plants, wooden cages were prepared, whose sides were 3 ft. in width and $4\frac{1}{2}$ ft. in height. The sides and tops of these cages were covered with zinc coated wire gauze, of 400 meshes to a square inch. The cages were permanently placed on single plants throughout the cotton-growing season (Plate VII, Fig. 1), and the plants were kept under observation.

In addition to these cages, a big permanent cage was constructed which was 15 ft. broad, 20 ft. long and 9 ft. in height. The sides and the top of this cage also were covered with similar wire gauze. This was big enough to accommodate 30 plants spaced 3 ft. by 3 ft. The plants under this cage also were kept under observation.

Within a few weeks, it was observed that all these cages were quite effective in keeping off the boll-worms and for a time it appeared that they would suit our purpose very well. It was, however, not very long before we realised that the plants enclosed in these cages were growing in an abnormal way. They appeared to be tall and lanky, and had thin broad leaves. Most of them reached the tops of the small cages very soon, and some of the plants in the big cage also, to our great surprise, reached its top, which was 9 ft. from the ground (Plate VII, Fig. 2).

In the face of such abnormal growth it was not possible to utilise these plants for getting a correct idea about the maximum capacity of yield of the plants when they were grown free from boll-worms. The need for a better method which would efficiently protect the plant from boll-worms, and at the same time allow them to grow quite normally, was very keenly felt.




FIG. 2. Cotton plant grown free from boll-worms in a plant house with sides and top of wire gauze.




FIG. 1. Plant cage with wire gauze sides and top.

PLATE VIII.




FIG. 1. A cotton plant grown free from boll-worm by night-caging. (Note the tin trench round the stem of the plant.)




FIG. 2. Plant curtains used for night-caging.

NIGHT-CAGING METHOD

In the meanwhile, detailed work of the life-history and habits was in progress in the laboratory which afforded ample evidence that the Spotted Boll-worms were incapable of laying eggs, during the day time, and it was therefore needless to enclose the plants during the day for preventing oviposition by *Earias* moths. The permanent cages were therefore discarded, and it was decided to cage the plants during the night only. Night caging prevented the moths from laying eggs on these plants, but there still remained the possibility of the boll-worm larvae crawling from the non-caged to the caged plants. It was, therefore, necessary to keep small tin trenches, with water and oil in them, round the stem of each plant to catch the worms which would try to encroach. This new method has been hereafter referred to as "Night Caging" method, and its important details are enumerated below :—

(1) Light wooden cages are prepared and are covered with mosquito netting cloth on the sides and the tops. They are placed on the plants every evening before sunset and are removed every morning after sunrise.

(2) Small tin trenches 7 inches in diameter, 3 inches in height and having a circular opening in the centre (3 inches in diameter) for accommodating the stem of the plant, are prepared. These are slipped round the plants before they are 6 to 8 inches in height. A small quantity of water with a thin film of non-volatile oil (such as sesamum oil) is kept in the trenches for catching any of the larvae which may attempt to cross over to the plants selected for caging (Plate VIII, Fig. 1).

(3) The third essential for the success of the method is to shake the plants lightly every evening to drive away the moths, if there be any, before placing the cages, for the night.

To start with, only single plant cages were utilised for this method. It was, however, subsequently modified to cover 3 or 5 plants simultaneously under one cage. The wooden frame work was dispensed with, and iron pipes were fixed in the soil, round groups of plants and mosquito nets of suitable sizes were slipped on these supports (Plate VIII, Fig. 2).

The trenches and the nets were found to be quite effective in keeping off the boll-worms, and the plants grew quite normally as they received full ventilation during the day time. It must be mentioned that this device also prevented the attack of the shoot-roller (*Phycita infusella* Meyr.) and to a great extent of the Pink Boll-worm (*Platyedra gossypiella* Saund.) as well.

A few stray cases are, however, noticed of accidental infection by the boll-worms. The Pink Boll-worm moths especially are smaller in size and therefore they sometime create a good deal of trouble to the developing bolls. A careful watch has to be kept to detect such infection for removing it as soon as it appears.

OBSERVATIONS ON THE CAGED PLANTS

Only ten plants were grown during each of the first two seasons (1925-26 and 1926-27) under night caging. During the third season (1927-28), 20 plants were

similarly grown. During all the three seasons complete records of the growth of the main stem, periodical flower-bud formation, and flower opening were maintained; and the proportion of the periodical flowers to the bolls which finally yielded the *kapas* was also worked out. All these observations were also maintained on an equal number of non-caged plants for comparison.

More extensive trials on a larger number of plants were subsequently undertaken during the next three seasons, especially for determining the yield of *kapas* in the absence of boll-worms. The observations during the first three seasons have helped to give a more or less accurate picture of the cotton plant as it would grow if the boll-worms are eliminated.

GROWTH OF MAIN STEM AND BRANCHES

It has already been mentioned that the tender vegetative shoots of the main stems, as well as, of the branches are destroyed in large numbers by the Spotted Boll-worms and shoot rollers, under normal field conditions. The caged plants, however, escape this injury, and as a result the height of their main stems is greater, and they have a larger number of primary sympodia than the control plants, so that ultimately the caged plants present a more regular conical appearance (Plate IX, Figs. 1 and 2).

TABLE XXIV

Number of primary sympodia per plant and the average height of the caged and the uncaged plants

| — | Year | Plants under observation | No. of primary sympodia per plant | Average height of main stems cms. |
|---------------------|---------|--------------------------|-----------------------------------|-----------------------------------|
| Caged | 1925-26 | 8 | 26.8 | 89.0 |
| Non-caged | 1925-26 | 25 | 5.7 | 38.0 |
| Caged | 1926-27 | 10 | 23.0 | 82.7 |
| Non-caged | 1926-27 | 20 | 13.4 | 59.2 |
| Caged | 1927-28 | 18 | 29.1 | 109.0 |
| Non-caged | 1927-28 | 16 | 8.6 | 54.6 |

FORMATION AND SHEDDING OF FLOWER-BUDS AND BOLLS

The flower-buds begin to appear about the same time both on the caged and the uncaged plants. During some of the seasons, most of the earliest formed flower-buds drop away, due to unfavourable environmental conditions, and then follows a period of rapid development of flower-buds. It is, however, about this time that



FIG. 1. Right:—Cotton plant protected from boll-worm by night-caging.
Left:—Unprotected plant.






FIG. 2. Diagrammatic sketch of a cotton plant grown free from boll-worm by night-caging.



PLATE X.

— weekly flower-buds formed.
 - - - weekly flowers opened.
 - - - bolls matured out of the above flowers.



Showing weekly performance of caged and uncaged plants during three seasons at Surat.

1926-27.



— weekly flower bud formation. —•—•—•— weekly flower opening
---- weekly shedding of flower buds. -x-x-x- weekly boll shedding (relative).
--- weekly shedding of flower buds, . weekly boll shedding (relative)
 showing bollworm injury. showing boll worm injury.

Showing weekly flower-bud formation, shedding of flower-buds, opening of flowers and shedding of bolls from caged and uncaged plants at Surat during 1926-27.

the Spotted Boll-worms begin their attack on the developing buds, and we have seen that the attacked buds shed in large numbers every week from the unprotected plants. In the absence of this damage the number of developing flower-buds on the caged plants soon increases, and an intense flowering can be seen on these plants very much earlier than in the case of the control plants (Appendix I, Tables XIV, XV and XVI).

Later on when the flowering commences on the uncaged plants, the earliest flowers and young bolls fall a victim to the boll-worms in large numbers and hence the boll-development is delayed. It is important to note that a very large percentage of the earliest flowers develops into mature bolls on the caged plants, whereas the success of early flowers into bolls is comparatively very low on the uncaged plants. It is also necessary to note that as soon as a certain number of bolls have begun to develop on the caged plants from the earlier flowers, the later flowers drop away in large numbers (Appendix I, Tables XVII to XIX). (Plate X.)

It can be seen from the tables mentioned above that the difference between the periods of flower-bud formation, flower-opening, and boll-development was very marked during 1926-27. This difference is shown in graph form in Plate XI.

The early intensity of flowering and a larger success from the first flowers naturally leads to an earlier crop of *kapas* from the caged plants, and in some years the *kapas* is ready for picking on the caged plants 4 to 6 weeks earlier than in the general cotton fields.

The numbers of flower-buds which open into flowers and of flowers which develop into mature bolls on the caged and the uncaged plants are analysed below:

TABLE XXV
(Plants spaced 3 feet by 3 feet)

| | No. of plants observed | No. of flower-buds formed per plant | No. of flowers opened per plant | Percentage of flowers to flower-buds | No. of bolls matured per plant | Percentage of mature bolls to flowers | Percentage of mature bolls to flower-buds |
|------------------|------------------------|-------------------------------------|---------------------------------|--------------------------------------|--------------------------------|---------------------------------------|---|
| 1925-26. | | | | | | | |
| Caged plants . | 8 | 229 | 89 | 38.9 | 31 | 35.4 | 13.7 |
| Control plants . | 25 | 257 | 73 | 28.9 | 28 | 38.3 | 10.97 |
| 1926-27. | | | | | | | |
| Caged plants . | 10 | 243 | 89 | 36.6 | 33.0 | 37.7 | 13.7 |
| Control plants . | 20 | 297 | 59 | 19.9 | 24.0 | 40.2 | 7.97 |
| 1927-28. | | | | | | | |
| Caged plants . | 20 | 445 | 166 | 37.4 | 56 | 33.8 | 12.7 |
| Control plants . | 16 | 411 | 96 | 23.4 | 37 | 38.4 | 9.0 |

From column No. 4 of the above statement it is seen that during the 3 seasons under consideration, the percentage of flower-buds that opened into flowers on caged plants varied between 36.6 and 38.9, whereas in the case of the non-caged plants this percentage varied between 19.9 and 28.9. It will be evident from these figures that a very much larger proportion of flower-buds open into flowers on the cotton plants when they are protected from the boll-worms.

Column No. 6 of the same statement, however, indicates that the percentage success from flowers to mature bolls on the caged plants is slightly lower than on the non-caged plants. This is probably due to the fact that a disproportionately higher number of flowers open on the caged plants and hence the percentage of the flowers ultimately retained on them as mature bolls declines to a small extent. The ultimate success of the flower-buds into mature bolls on the caged plants, as can be seen from column No. 7 of the last statement, was higher than on the uncaged plants during all the three seasons. The percentage success of flower-buds to bolls on the caged plants during the three years varied between 12.7 and 13.7, whereas it varied between 7.9 and 10.97 in the case of the control plants.

The last column will further show that the caged plants produced 11 to 51 per cent more mature bolls than their controls during the three seasons under consideration. During 1927-28, *kapas* from these plants was weighed and it was found that the caged plants produced 100 grams of *kapas* per plant, and the uncaged plants 66 grams. The caged plants thus produced 52 per cent more *kapas* during that season.

20 plants were grown by night-caging on the Broach Farm during 1927-28 and parallel observations are given below :

TABLE XXVI

Caged and uncaged plants on Broach Farm

1927-28

| — | Plants observed | Flower- buds per plant | Flowers per plant | Bolls per plant | Percent- age flowers to flower buds | Percent- age bolls to flowers | Percent- age bolls to flower buds | Weight of <i>kapas</i> per plant (Tolas) |
|--------------------|--------------------|------------------------------|----------------------|--------------------|--|--|---|---|
| Caged plants . . | 19 | 677 | 255 | 91 | 37.7 | 35.7 | 13.4 | 13.77 |
| Non-caged plants . | 12 | 530 | 177 | 66 | 33.4 | 37.3 | 12.5 | 8.31 |

These caged plants yielded about 25 bolls more per plant than their controls and produced about 66 per cent more *kapas*.

The observations, mentioned so far, though made on a small number of plants, clearly indicated that if the cotton plants could be grown free from boll-worms, they

would grow into better plants in various respects and would undoubtedly produce a larger number of bolls. The crop would besides be ready about 4 to 6 weeks earlier.

NIGHT CAGING ON A LARGE SCALE

More extensive trials were undertaken during the next three seasons of 1928-29, 1929-30 and 1930-31 at Surat, in order to get a more significant idea about the increase in the yield of *kapas* which would result if the boll-worms were eliminated.

During each of the three seasons, a plot of about one acre was reserved for growing cotton, and the plants were spaced 4 feet by 4 feet. The plot was divided into three blocks, and the central block was used for caging the plants. More than 100 plants were marked out in groups of three, because each mosquito net could only cover 3 plants. A similar number of plants was also marked out in each of the two side blocks in groups of 3 plants, and these groups were located in the same design as in the central block.

These plants were not tampered with, for making any detailed observations, until the bolls matured. The *kapas* from sound and damaged locks from each one of these plants was collected and weighed separately.

MONSOON DURING THE SEASONS, 1928-29, 1929-30, 1930-31

A note upon the peculiarities of the monsoon during the three seasons of these trials will be useful for a better understanding of the performance of the plants under observation.

During the season of 1928-29, the total rainfall was a good deal lower than in the other two seasons (Appendix I, Table VII). It was, however, well distributed, and, the rainfall during the months of July and August being moderate, conditions for vigorous growth of the plants prevailed during the earlier part of the season.

During the next year, 1929-30, out of the total of about 48 inches, 33 inches of rainfall was received during the first fortnight of July. This necessitated the resowing of cotton which was done on the 17th of July 1929.

In 1930-31 also the total rainfall exceeded 40 inches but it was very much better distributed. On the 29th of October, a severe storm passed over Surat accompanied by rainfall (95 cents) and in several plots even the cotton plants were lodged to the ground.

GROWTH OF CAGED PLANTS

Although detailed growth records of these plants were not maintained, the general observations showed that the caged plants during all the three seasons were much earlier in flower-bud and boll-formation than the uncaged plants, and finally it was possible to collect *kapas* from these plants about 4 to 6 weeks earlier than from the general field plants (Appendix I, Table XX).

TABLE XXVII

Period of picking of kapas from the caged and uncaged plants

| Year | From caged plants | From uncaged plants |
|-------------------|---|--|
| 1928-29 | 7th January 1929 to 7th March 1929. | 8th February 1929 to 22nd March 1929. |
| 1929-30 | 1st January 1930 to 11th February 1930. | 20th February 1930 to 24th March 1930. |
| 1930-31 | 5th January 1931 to 10th March 1931. | 20th February 1931 to 19th March 1931. |

During the first two seasons the effective crop of *kapas* from the caged plants was collected before the end of February. It was slightly delayed during 1930-31 due to the storm of 29th October 1930 which destroyed a very large number of developed flower-buds and bolls from the caged plants. It was, therefore, seen that normally January and February would be the months of harvesting of *kapas* instead of February and March if the crop is saved from the boll-worms.

When all the pickings were over, all the plants from the cages and the Northern control were cut at the cotyledonary nodes and were hung up in the fields for being dried for about 3 weeks, and their air-dry-weights were recorded. All the remaining leaves from these plants were nipped out before cutting the plants.

The weights of these plants were recorded for comparing the growth of the caged and uncaged plants and for comparing the *kapas* produced by these plants in relation to their final dry weights.

TABLE XXVIII

Dry weight of scaffolding and crop

| — | No. of plants observed | Average dry weight of plants in grms. | Average weight of <i>kapas</i> per plant in grms. | Amount of <i>kapas</i> produced to unit dry weight |
|--------------------------|------------------------|---------------------------------------|---|--|
| 1928-29. | | | | |
| Caged plants | 89 | 204.0 | 236.6 | 1.16 |
| Uncaged plants | 104 | 137.3 | 130.6 | 0.95 |
| 1929-30. | | | | |
| Caged plants | 113 | 92.1 | 174.0 | 1.89 |
| Uncaged plants | 112 | 99.2 | 113.2 | 1.14 |
| 1930-31. | | | | |
| Caged plants | 118 | 98.7 | 159.7 | 1.62 |
| Uncaged plants | 116 | 126.2 | 150.0 | 1.19 |

It was found that during the first season the dry weight of the caged plants was distinctly higher than the controls. Their weights did not differ to a great extent during the second season, but during the third season the dry weight of the caged plants was distinctly lower. The capacity of the caged plants for producing *kapas* per unit dry weight, or, in other words, the yield efficiency of the caged plants for a given weight of their expanse, was higher during all the seasons and was particularly marked during the last two seasons.

INCREASED YIELD DUE TO CAGING

The caged plants produced more *kapas* than the control field plants during all the three seasons. This increase, however, varied considerably. During 1928-29, the caged plants yielded 81 and 102 per cent more *kapas* than the two controls respectively. During the next season this increase over the control plants amounted to 53 and 54 per cent and during the third year of these trials, the caged plants produced only 6.5 and 15 per cent more *kapas* than each of the two sets of controls respectively.

TABLE XXIX

Yields of caged and uncaged plants at Surat

| | 1928-29 | | | 1929-30 | | | 1930-31 | | |
|---------------------------------|-----------------|---------------------------------|--|-----------------|---------------------------------|--|-----------------|---------------------------------|--|
| | Plants observed | <i>Kapas</i> per plant in grms. | Percentage increase of yield over control plants | Plants observed | <i>Kapas</i> per plant in grms. | Percentage increase of yield over control plants | Plants observed | <i>Kapas</i> per plant in grms. | Percentage increase of yield over the control plants |
| Caged plants | 99 | 236.6 | | 113 | 174.0 | | 118 | 159.7 | |
| Uncaged plants (North-control) | 104 | 130.6 | 81 per cent over the North Control | 112 | 113.2 | 54 per cent over the North Control | 116 | 150.0 | 6.5 per cent over the North Control |
| Uncaged plants (South control). | 100 | 117.4 | 102 per cent over the South Control. | 112 | 113.8 | 53 per cent over the South Control. | 118 | 138.5 | 15 per cent over the South Control |

INCREASE IN THE YIELDS OF CAGED PLANTS IN RELATION TO THE INCIDENCE OF BOLL-WORMS AND OTHER ENVIRONMENTAL CONDITIONS

It might be expected that the differences in yield, between the plants, protected and unprotected from the boll-worms, should vary according to the intensity of the

infestation of the Spotted Boll-worms; and that it may therefore be inferred that the incidence of the boll-worms was the highest during 1928-29 and the least during 1930-31. A careful scrutiny of the weekly records of the population of the Spotted Boll-worms in cotton fields (Table XXII) will not, however, warrant such an inference. In fact the observations show that the intensity of infestation by the Spotted Boll-worms was almost similar during the last two seasons, and that it was a little lower during 1928-29, when the highest differences between the caged and the non-caged plants were recorded.

Information upon the incidence only of the Spotted Boll-worms is insufficient to account for the seasonal differences between the plants, protected and unprotected from this pest, and it is necessary to study the variations in the other environmental conditions as well. It was not within the scope of this enquiry to study in detail the effect of all the various environmental conditions on the plant growth; but the outstanding factors which apparently affected the differences between the caged and uncaged plants must be briefly discussed.

It has been shown that flower-bud formation, flower opening and boll-development take place very much earlier on the caged than on the uncaged plants, and that their *kapas* can be collected about 4 to 6 weeks earlier. This means that the caged plants pass through all their stages of growth a few weeks earlier than the control plants, and consequently the environmental conditions of soil and climate at the various stages of life of both sets of plants are not identical.

Generally, the soil and climatic conditions are more favourable for plant growth during the earlier part of the growing season, and they, therefore, help to increase the differences between the caged and the uncaged plants.

During the month of July at Surat heavy showers of rain are usual, which cause water logging of the soil. These heavy showers were not received during 1928. The total rainfall of the season was not very heavy and it was well distributed. Conditions for plant growth were therefore favourable during the early part of this season, and consequently the caged plants grew vigorously, and ultimately their average dry weight was found to be 204 grams per plant, *i.e.*, about twice the average weight for the other two seasons. The uncaged plants also flourished during this period but their growth was hampered, and their period of flower-bud formation and boll development was delayed, due to the damage caused by the Spotted Boll-worms during the months of September, October and November. They could not, therefore, take full advantage of the favourable environmental conditions which were available during the early part of the season. There was, therefore, a wide difference between the yields of the caged and uncaged plants during this season.

During the next season, there was a very heavy rainfall during July, and the conditions were not quite as favourable for very early growth as during the previous season. The conditions, however, improved considerably after the end of July, and ultimately the caged plants yielded about 54 per cent more *kapas* than their

controls although the final air-dry-weight of the caged plants was slightly less than that of the uncaged plants.

In 1930-31, the caged plants were progressing quite satisfactorily, and flower-opening had commenced on these plants during the month of October, when flower-buds had just commenced to develop on the uncaged plant. A storm passed over Surat on the 29th of October and all the plants were rudely shaken and a few of them had their branches broken. The result of this storm was more disastrous to the caged plants because they had very much advanced in growth and therefore lost many of their developed forms. During the period of about 5 or 6 days after the storm, 8,105 shed flower-buds and 2,520 young shed bolls were collected from all the caged plants. Thus the caged plants after having lost most of their developed forms, had to begin afresh, practically even with the non-caged plants, and hence the difference in the yields of the two sets of plants was meagre.

It is, therefore, seen that the other environmental conditions play an important part in increasing or decreasing the seasonal differences in yields of *kapas* from the plants protected and un-protected from the boll-worms. The balance in favour of the boll-worm free crop may be masked, in abnormal seasons, by the depressing effect of other factors in the environment.

CAGED PLANTS AND THE RED COTTON BUG

Red cotton bug (*Dysdercus cyngulatus*) is not a serious pest at Surat. However, it was found that in November 1929, a large number of these bugs began to appear only on the caged plants, and their numbers increased considerably during the month of December. Nearly 1,450 were collected from the caged plants during that season to save the developed bolls from their injury. In 1930-31, again a very large number appeared on the caged plants during boll development period, whereas they were rarely noticed on the general crop.

The apparent reason for this concentration is that the bugs prefer well developed bolls for their attack, and a large number of these are found on the caged plants at a time when such bolls are rarely met with on the general crop.

CAGING AND THE QUALITY OF *kapas*

It has already been stated that the *kapas* from the sound and the damaged locks from all the bolls of the caged and the uncaged plants was collected and weighed separately. The damaged *kapas* consisted of locks which did not develop normally, either due to the damage of the boll-worms (*Earias fabia*, *Earias insulana* and *Platyedra gossypiella*), or due to other causes.

TABLE XXX

Proportion of kapas from sound and damaged locks

| | SALEABLE <i>kapas</i> PER PLANT IN GRAMS | | Total <i>kapas</i> per plant in grms. |
|--------------------------|---|--------------------------|---|
| | From sound locks | From damaged locks | |
| 1928-29. | | | |
| Caged plants | 221 1 | 15.4 | 236.5 |
| Control plants | 105 5 | 25 1 | 130 6 |
| 1929-30. | | | |
| Caged plants | 171 8 | 2 2 | 174.0 |
| Control plants | 107.9 | 5 3 | 113 2 |
| 1930-31. | | | |
| Caged plants | 152.2 | 7.9 | 160.1 |
| Control plants | 140 7 | 9 4 | 150.1 |

A further classification of the number of locks which yielded saleable and unsaleable *kapas*, and which were partially or completely destroyed by the boll-worms, will be found in Appendix I, Table XXI.

After the pickings of *kapas* from the caged and the uncaged plants were over during 1929-30, we found that the *kapas* from the caged plants appeared distinctly coarser. All the *kapas* from the good locks from both the groups of plants was, therefore, ginned and it was found that the ginning percentage of the *kapas* from the caged plants was 37.3, whereas it was 34.4 in the case of the *kapas* from the uncaged plants.

This difference was naturally attributed to the fact that the environmental conditions for the development of bolls from the caged and uncaged plants were not identical.

In order, therefore, to get a better idea of the effect of caging on the quality of *kapas*, all the lint from the good locks of the caged and the uncaged plants, grown during 1929-30 and 1930-31, was sent to the Matunga Technological Laboratory for spinning tests. The detailed reports are given in Appendix II.

It will be seen from these reports that the lint from the caged plants of 1929-30 was valued at Rs. 150 over the contract rate, whereas the control sample from the

uncaged plants was valued at Rs. 120 over the contract rate. The lint from the caged plants of 1930-31, however, was valued at Rs. 70 over the contract rate, whereas value of the control sample of that season was Rs. 75 over the contract rate.

It is also interesting to note that the staple length of the lint from the protected plants was full $1\frac{1}{2}$ inch, whereas it was $1\frac{1}{8}$ in the case of the lint from the control plants of 1929-30. There was, however, no difference in the staple length of the two samples during the next season.

In spite of all this, it is essential to remember the concluding remarks of the Director of the Technological Laboratory, Matunga. In his first report, he says "The lint from the caged plants, though possessing a somewhat longer staple, is decidedly coarser and rather inferior in spinning performance to the lint from the control plants."

In short, the various observations mentioned in this chapter have shown that if the cotton crop were saved from the Spotted Boll-worms, not only would damage to the various parts of the cotton plant prevented but the crop would be able to make full use of the environmental conditions for growth which are generally more favourable during the earlier part of the growing season ; and that the crop would be ready earlier by about 4 to 6 weeks with the ultimate result that we could normally expect a substantial increase in the production of *kapas*.

STUDY OF CONTROL MEASURES

In the introductory chapter reference was made to some of the control measures, which were advocated by the previous workers. Destruction of the affected cotton shoots at the commencement of the cotton-growing season, use of *bhendi* (*Hibiscus esculentus*) as a trap crop and introduction of parasites were some of these measures. To study their efficacy under the conditions of this tract was, therefore, a part of the programme at the commencement of these investigations.

I. REMOVAL OF ATTACKED SHOOTS

It has already been stated that it is with the cessation of the monsoon that the cotton plant begins to grow vigorously in this tract, and it is then that the Spotted Boll-worms begin to multiply rapidly. A majority of the population of these worms, during the first few weeks of this growing period is found feeding on the tender vegetative shoots (Appendix I, Tables VIII to XIII). Many of the larvae bore into the succulent top internodes and destroy the shoots, which wither and dry up. It was, therefore, necessary to discover whether, if the attacked shoots were collected and destroyed, the further progress of the pest would be checked to a considerable extent.

Two experiments were, therefore, laid out during 1925-26. In the first, an area of one acre was kept under observation during the whole of the period in which the attack was concentrated on shoots. All the attacked shoots, which could easily

be observed by an ordinary labourer from the appearance of the withering tops, were collected once a week and were examined to note the number of *Earias* larvae in them. In the second experiment, a fresh group of 1,000 plants was examined, a little more critically than in the previous case, and all the withering shoots as well as those which showed holes bored by the boll-worms prominently were collected for examination. The number of larvae found every week was compared with the number which was expected to be present. The probable population in these plots was calculated from the number of larvae which were recorded in an exhaustive weekly examination of 25 cotton plants. (Appendix I, Table VIII.)

TABLE XXXI

Number of Earias larvae collected from shoots from the two plots and the expected population of larvae from each of them

| Weeks | Total population of <i>Earias</i> larvae from 25 plants | ONE ACRE PLOT (ABOUT 4,800 PLANTS) | | | 1,000 FRESH PLANTS EVERY WEEK | | |
|-----------------------|---|------------------------------------|-----------------------------|---|-------------------------------|-----------------------------|---|
| | | Withered shoots | No. of <i>Earias</i> larvae | Expected population of <i>Earias</i> larvae | Attacked shoots | No. of <i>Earias</i> larvae | Expected population of <i>Earias</i> larvae |
| 19th September 1925 . | 32 | 86 | 25 | 6,144 | 282 | 139 | 1,280 |
| 26th September 1925 . | 48 | 53 | 19 | 9,216 | 313 | 154 | 1,920 |
| 3rd October 1925 . | 48 | 33 | 6 | 9,216 | 240 | 94 | 1,920 |
| 10th October 1925 . | 43 | 27 | 6 | 8,256 | 248 | 132 | 1,720 |
| 17th October 1925 . | 68 | 25 | 5 | 13,056 | 178 | 97 | 2,720 |
| 24th October 1925 . | 87 | 41 | 15 | 16,704 | 232 | 104 | 3,480 |
| 31st October 1925 . | 73 | 21 | 3 | 14,016 | 194 | 88 | 2,020 |
| 7th November 1925 . | 45 | 16 | <i>Nil</i> | 8,640 | 131 | 30 | 1,800 |

The comparison showed that the number of larvae removed from the plot of an acre was almost insignificant, compared with the population expected to be present in that area and the number of larvae collected from a fresh group of 1,000 plants every week showed that hardly ten per cent of the total population of the boll-worms, calculated from the detailed examination of twenty-five plants, would be destroyed even on careful search.

It is thus seen that only such shoots, whose damage can be easily noticed, can be collected by this method, and that there is a very large percentage of population

which remains over in the fields as the damage caused by them is not easily perceptible. Much significance cannot, therefore, be attached to this method of controlling the Spotted Boll-worms by collecting and destroying such of the attacked shoots as can be easily observed by the cultivators.

2. TRAP CROPS

Bhendi (*Hibiscus esculentus*) is known to be a favourite food plant of the Spotted Boll-worms, and its cultivation in the cotton fields has been advocated for trapping these worms. [Javeri, 1921]. It was therefore necessary to ascertain the effect of using this trap crop as a control measure.

In most of the villages round about Surat, *bhendi* is grown at the commencement of the monsoon and large quantities of *bhendi* pods are available in the Surat market during the months of August and September. The plants from these plots are severely attacked by the Spotted Boll-worms, and by the month of October even the pods which are sent to the market from these plots show a considerable amount of infestation.

TABLE XXXII

Bhendi pods purchased from the Surat market

| Date of examination | Weight of the pods examined | No. of pods examined | No. of <i>Earias</i> larvae |
|-------------------------------|-----------------------------|----------------------|-----------------------------|
| 10th August 1929 | 3 | 144 | 1 |
| 17th August 1929 | 5 | 271 | 4 |
| 24th August 1929 | 5 | 181 | 5 |
| 31st August 1929 | 5 | 215 | 6 |
| 6th September 1929 | 5 | 213 | 7 |
| 13th September 1929 | 10 | 388 | 18 |
| 20th September 1929 | 10 | 443 | 27 |
| 28th September 1929 | 5 | 202 | 30 |
| 4th October 1929 | 5 | 209 | 22 |
| 12th October 1929 | 5 | 259 | 55 |
| 15th October 1929 | 5 | 229 | 60 |

The plants from these plots finish their life by the end of October just when the cotton plants begin their vigorous growth and the Spotted Boll-worms begin to

increase in the cotton fields. All the *Earias* population liberated from the *bhendi* plots naturally transfers its attention to the neighbouring cotton crop.

What actually happens is that the good pods are picked out and the affected pods mostly remain on the plants, and even if the plants are destroyed at the end of the season along with the larvae feeding on them, the pupae in the soil and the adult moths at any rate escape destruction.

Observations at Jalgaon (East Khandesh) during 1929 showed that a few rows of *bhendi* were generally sown in the cotton plots, and that these plants grew luxuriantly and had a large population of Spotted Boll-worms on them (Appendix IV, Table II). Further it was noticed that in Khandesh the life of the *bhendi* plants was over long before the cotton crop was ready and the result was that the *Earias* population on the cotton crop suddenly increased as soon as the *bhendi* crop was over (Appendix IV, Table I and Plate XXII). Similar observation during 1930 showed that the *Earias* population on monsoon *bhendi* was meagre during the months of August and September due to heavy rainfall, and consequently the increase of the Spotted Boll-worms on cotton was more gradual than during the previous season.

It was, therefore, considered that the method of growing *bhendi* in the midst of cotton areas was harmful to the cotton crop, and it was probable that *bhendi* as a trap crop would only be useful if it continued its life and trap these worms till the cotton crop was over; or at least till the end of December when the *Earias* population normally declined.

Different varieties of *bhendi* were, therefore, tried both at Surat and Jalgaon for selecting a type which would be most suitable as a trap crop. An attempt was also made to sow these varieties late in the season in order to delay their growing period. The names of the varieties are given below :—(1) Surat 5, (2) Surat 9, (3) Katai *bhendi* (from Khandesh), (4) American long green, (5) American dwarf green and (6) American white velvet. In Jalgaon in addition to these varieties “Nasik” *bhendi* was also tried.

LATE SOWINGS

At Surat the first sowing of *bhendi* was done on the 26th of June, and these plants came on well. The plants sown in the middle of July made fairly good growth. The germination, however, was very defective in the case of sowings made after the middle of August, and the plants which germinated were very poor in growth.

In Khandesh also the June sowing was successful but the later sowings, which were repeated every fortnight, gave very poor germination and the growth of the few plants which germinated was very poor.

COMPARISON OF THE VARIETIES GROWN AT SURAT

All the varieties, except the Katai *bhendi* suffered heavily from the attack of *Jassids*, which commenced from the middle of August and continued till the end

of September. Aphis infestation also appeared on these plants during this period but was soon checked by lady-bird beetles.

It was further noticed that manuring was necessary at Surat to secure satisfactory growth of these plants. The plants which were manured at the time of sowing with castor cake grew satisfactorily, whereas the plants from the un-manured plots remained stunted.

Twenty-five plants of each of these varieties were kept under observation and the mature pods from these plants were collected, as they became ready, and were examined for *Earias* larvae (Appendix I, Tables XXII, XXIII and XXIV).

Surat 9 began to yield pods in the last week of August, while pods could be collected from Surat 5 from the 5th of September. Surat 5 and Katai *bhendi* from manured plots continued to yield pods practically till the end of January and many of these pods were infested by boll-worms. Katai *bhendi* is a hardy plant, with rough leaves and pods. It is a shy bearer and attracts a smaller number of Spotted Boll-worms than the other types.

All the American varieties turned out to be very late and began to bear pods at Surat from the beginning of December, and continued till the end of January.

On the whole it appeared that Surat 5, if manured at the time of sowing, would continue to trap the worms almost throughout the cotton-growing period and would therefore be a suitable type for trying as a trap crop at Surat.

VARIETIES TRIED IN KHANDESH

At Jalgaon, Nasik type of *bhendi* was the earliest to bear pods and to attract the *Earias* boll-worms. Nasik and Surat 9 finished their life at the end of October, whereas Surat 5 continued to bear till January. The American types began to bear at the end of September and continued till January (Appendix I, Tables XXV, XXVI and XXVII).

It appeared that it would be necessary to sow a mixture of Nasik and Surat 5, or Nasik and one of the American varieties, at the commencement of the monsoon to get a continuous stand of *bhendi* plants for trapping the boll-worms.

INFESTATION ON COTTON SURROUNDED BY *bhendi* PLANTS

These trials indicated a possibility of securing a continuous stand of *bhendi* plants during the cotton-growing season but it was necessary to judge the extent of *Earias* population on the cotton plants growing in close proximity of *bhendi*. A plot of cotton (of about 10 *gunthas*) was laid out at Surat surrounded by two rows of Surat 5, and two rows of Surat 9. The population of *Earias* larvae from 25 plants of cotton from this plot, was determined thrice during the season. Similar population records, from 20 plants each time, were available for comparison from a distant plot of cotton in the same field.

TABLE XXXIII

| 25 PLANTS FROM PLOT SURROUNDED BY <i>bhendi</i> | | | | | 20 PLANTS FROM A DISTANT PLOT OF COTTON | | | | |
|--|--------------------|--|--------------|--------------------------------------|--|--------------------|-------------------------------------|--------------|-------------------------------|
| Date of examination | No. of flower-buds | No. of <i>Earias</i> larvae in flower-buds | No. of bolls | No. of <i>Earias</i> larvae in bolls | Date of examination | No. of flower-buds | <i>Earias</i> larvae in flower-buds | No. of bolls | <i>Earias</i> larvae in bolls |
| 4th November 1930 | 298 | 23 | .. | .. | 31st October 1930 | 290 | 24 | .. | .. |
| 3rd December 1930 | 1,988 | 41 | .. | .. | 29th November 1930. | 1,474 | 22 | .. | .. |
| 10th January 1931 | 1,572 | 8 | 1,050 | 13 | 3rd January 1931 | 2,187 | 6 | 728 | 9 |
| Percentage of flower-buds with <i>Earias</i> larvae=1.8. Percentage of bolls with <i>Earias</i> larvae=1.2. | | | | | Percentage of flower-buds with <i>Earias</i> larvae=1.3. Percentage of bolls with <i>Earias</i> larvae=1.2. | | | | |

These results showed that the attraction of *bhendi* was not strong enough to reduce the infestation on the neighbouring cotton plants. On the contrary, it appeared to be slightly higher; probably because the moths attracted to the *bhendi* plants infested the neighbouring cotton plants as well.

THE UTILITY OF *bhendi* AS A TRAP CROP

Bhendi is no doubt a favourite food of the Spotted Boll-worms, and a large number of these insects can be collected from these plants. If a short season variety is grown at the commencement of the monsoon, it helps to breed a very large number of *Earias* population, and the destruction of these plants with *Earias* larvae and eggs still leaves the pupae and the moths to continue the damage to the cotton crop. It has been found possible to secure a continuous stand of *bhendi* plants during the cotton-growing season, but the attraction of *bhendi* is not so strong as to prevent the infestation on the neighbouring cotton plants.

It would, however, be very difficult to ensure concerted action from the cultivators in picking out and destroying the affected *bhendi* pods at regular intervals before the larvae feeding on them go to the soil for pupation.

It has been shown that *bhendi*, being a more suitable food for these worms, accelerates the multiplication of this pest (page Nos. 9 & 10). Besides this, experience at Surat showed that *Earias* larvae feeding on *bhendi* plants were practically immune from the attack of the two important parasites, viz., *Microbracon lefroyi* and *Actia aegyptia*. Time after time, large quantities of *bhendi* pods were examined during the course of the work of nearly seven years, but the presence of a caterpillar of *Earias*, attacked by *Microbracon lefroyi* or *Actia aegyptia* in these pods was extremely rare, even at a time when the parasites were abundant in the cotton fields. This naturally afforded more favourable conditions for the increase of this pest.

While no definite conclusion, that *bhendi* is effective as a trap crop in the control of *Earias* boll-worms, can be stated upon trials restricted to a single season, the indications are against the possibilities of its successful use and that the disadvantages of growing these plants in the midst of cotton areas would be greater than the advantages.

3. PARASITES OF THE SPOTTED BOLL-WORMS

The following seven parasites of the Spotted Boll-worms were found during our investigations at Surat.

- (1) *Trichogramma* sp.
- (2) *Microbracon lefroyi* D. and G.
- (3) *Rhogas testaceus* Grav.
- (4) *Actia aegyptia* Vill.
- (5) *Apanteles* sp.
- (6) *Chelonus* sp.
- (7) *Melcha nersei* Cam.

We are indebted to M. A. Husain (Punjab) for the identification of most of these parasites.

All the important parasites of the Spotted Boll-worms, noted in other provinces, were thus present in this tract also, and there were no chances, therefore, of importing and establishing a new parasite from any other locality to effect control in Surat.

Observations during the first two or three seasons at Surat showed that the population of the Spotted Boll-worms in the cotton fields considerably declined every year at the end of December. At first though it appeared that the decline was purely due to the cold weather; but it was also observed that during November and December the following three parasites were active at Surat; and therefore, it was considered that the reduction in *Earias* population must be partly, if not wholly, due to the activities of these insects—

- (1) *Rhogas testaceus* Grav.
- (2) *Actia aegyptia* Vill.
- (3) *Microbracon lefroyi* D. and G.

It was, therefore, decided to study the life and habits of these parasites.

Rhogas testaceus Grav.

This parasite is present throughout the year in the cotton fields at Surat, except during July and August. During September and October 1925, out of 1,014 *Earias* larvae collected, 22 caterpillars (i.e., about 2 per cent) were attacked by this parasite.

A few attempts were made to observe the behaviour of this parasite in captivity. Pairs of the adult parasites were placed in round glass jars about 4 inches in diameter and 5 inches in height. Host larvae of different sizes were offered to the parasites, but it was found that they preferred only the small ones, which were only 3 or 4 days old, for laying eggs. The parasite repeatedly stung the larva until it was fully paralysed and then thrust its ovipositor in its body and laid a single egg. Each female laid only 2 or 3 eggs in a day. The larva recovered within a few minutes and began to feed. In September the parasite larva became fully fed within 6 to 8 days after the egg was laid, and was ready to pupate. At this time the host larva succumbed and the parasite pupated in its body. The body of the host caterpillar

swelled to accomodate the full grown parasite and hence its intersegmental chitin was exposed. Before the host caterpillar succumbed, it generally came out of the bud or boll where it was feeding, and fastened itself to a bract of a flower-bud or flower, or to any other part of the plant. From the time of the death of the host larva, it took about 7 to 9 days for the adult parasite to emerge. Thus the whole life-cycle took about 13 to 17 days during September and October. Such parasitised caterpillars could often be seen in the cotton fields and they could be easily recognised by their characteristic appearance. This parasite cannot be considered to be a very important one as its capacity for multiplication is found to be very low.

Actia aegyptia Vill.

This fly was observed at Surat for the first time in November 1923, and it continued to breed till the end of July 1924. It reappeared again in October 1924 and was found to increase in numbers till the end of December 1924. It generally appeared every year in October or November when the *Earias* larvae began to feed on the flower-buds. During September and early in October a large number of *Earias* larvae were feeding on the shoots of the cotton plants but only one larva parasitised by this fly was found in the shoots. So also the *Earias* larvae in *bhendi* pods were never found attacked by this fly, although a very large population of these worms was present on the *bhendi* plants when this parasite was active in the cotton fields.

During September and October 1925, of 1014 *Earias* caterpillars, 69 were found to be parasitised by this fly. Both *Earias fabae* and *Earias insulana* were attacked by this parasite.

All attempts to breed this parasite in the laboratory failed and further information about its life and habits could not be secured.

Microbracon lefroyi D. and G.

This has been a well-known parasite in India since Lefroy recommended its importation in the Punjab after the failure of the cotton crop in 1905 [Lefroy 1906, 2].

In Bombay Presidency the presence of this parasite has been noted in Gujarat, in Khandesh, in Karnatak and in Jamrao canal tract of Sind.

Life of the adult.—The average life of the adult was found to vary from about 11 to 31 days during the different months at Surat. During November and December some of the adults lived for 58 and 59 days, i.e., for nearly 2 months.

Mating generally takes place on the day of emergence, or on the second day, and is repeated several times during the period of their life. No eggs are laid by the female wasp on the day of its emergence, and frequently a week or more elapses before oviposition commences under laboratory conditions, but it is not known if such an interval exists under natural conditions also. Normally oviposition begins on the second or third day after emergence of the female. Eggs are laid parthenogenetically also, but in such cases the progeny consists of males only.

TABLE XXXIV

Period of adult life of Microbracon lefroyi D. and G.

| Month of emergence of the parasite | No. of cases observed | Average life in days | Maximum life of an adult in days |
|------------------------------------|-----------------------|----------------------|----------------------------------|
| August 1926 | 2 | 14 | 21 |
| September 1926 | 17 | 12 | 21 |
| October 1926 | 29 | 15.6 | 38 |
| November 1926 | 49 | 22 | 58 |
| December 1926 | 17 | 31 | 59 |
| January 1927 | 9 | 25 | 48 |
| February 1927 | 22 | 26.6 | 41 |
| March 1927 | 10 | 15 | 27 |
| April 1927 | 3 | 11 | 22 |

Occasionally a few female parasites refused to lay eggs in captivity, although they continued to live for several days.

Oviposition.—The eggs are laid at any hour of the day time but the parasites are most active in the morning between 8 to 10 A.M.

Only fully grown *Earias* caterpillars are selected for oviposition, and both *Earias fabia* and *Earias insulana* are attacked. The host larvae are completely paralysed before eggs are laid. *Earias* caterpillars, freely moving about in the cage, were never attacked by these parasites. Oviposition was observed only when the host larvae were enclosed in their food, such as a piece of *blendi*, a flower-bud or a boll.

The female parasite locates the larva, which is not visible from outside, by tapping the food with its antennae. She then inserts her ovipositor through the hole through which the host larva has entered the food, or if the food material is sufficiently tender, the ovipositor is thrust through the material, and the caterpillar is paralysed. The eggs are then laid in groups of 2 or more on each larva. Instances of caterpillars with single eggs are also noticed but more than one is the general rule. The largest number of eggs laid on a single *Earias* caterpillar was found to be 22, while the largest number of eggs laid by a single parasite in one day was 40.

A few living *Microbracon lefroyi* were secured from Khandesh and the Punjab, and were bred separately at Surat. The capacity of multiplication of all these parasites from different localities are shown in the following tables. The number of eggs laid by these parasites was recorded only in a few cases because the material had to be disturbed for counting the eggs, and often the eggs got dislodged from the body of the host larva. From Table XXXVI it will be seen that the female parasite from pairs No. 14 and 16 laid 310 and 504 eggs respectively. There is a great variation in the number of adults obtained from each pair, which is probably due to the susceptibility of the parasites to the unnatural conditions of breeding captivity.

TABLE XXXV

Reproductive capacity of Khandesh Microbracon lefroyi in Surat in 1925

| PERIOD OF OBSERVATION | | Pair No | No. of eggs laid | Pupae | Males | Females | Total adults |
|-----------------------|---------------------|---------|------------------|-------|-------|---------|--------------|
| From | To | | | | | | |
| 25th August 1925 . | 12th September 1925 | 1 | 55 | 55 | 23 | 25 | 48 |
| 28th August 1925 . | 19th September 1925 | 2 | 87 | 67 | 20 | 38 | 58 |
| 28th August 1925 . | 7th September 1925 | 3 | 65 | 63 | 13 | 43 | 56 |
| 28th August 1925 . | 6th September 1925 | 4 | 16 | 9 | 2 | 7 | 9 |
| 3rd September 1925 . | 27th September 1925 | 5 | 64 | 60 | 22 | 33 | 55 |
| 3rd September 1925 . | 14th September 1925 | 6 | 64 | 58 | 12 | 38 | 50 |
| 5th September 1925 . | 23rd September 1925 | 7 | ..* | 66 | 10 | 43 | 53 |
| 9th September 1925 . | 26th September 1925 | 8 | .. | 50 | 6 | 16 | 22 |
| 9th September 1925 . | 28th September 1925 | 9 | .. | 68 | 19 | 37 | 56 |
| 10th September 1925 | 21st September 1925 | 10 | .. | 37 | 9 | 21 | 30 |
| 13th September 1925 | 5th October 1925 . | 11 | .. | 34 | 8 | 20 | 28 |
| 13th September 1925 | 25th September 1925 | 12 | .. | 27 | .. | 15 | 15 |
| 14th September 1925 | 6th October 1925 . | 13 | .. | 63 | 24 | 16 | 40 |
| 15th September 1925 | 3rd October 1925 . | 14 | .. | 37 | 18 | 9 | 27 |
| 16th September 1925 | 21st September 1925 | 15 | .. | 32 | 8 | 10 | 18 |
| 16th September 1925 | 30th September 1925 | 16 | .. | 22 | 17 | 4 | 21 |
| 17th September 1925 | 3rd October 1925 . | 17 | .. | 28 | 19 | 5 | 24 |
| 24th September 1925 | 5th October 1925 . | 18 | .. | 36 | 8 | 23 | 31 |
| 27th September 1925 | 12th October 1925 . | 19 | .. | 16 | 2 | 8 | 10 |
| 26th September 1925 | 1st October 1925 . | 20 | .. | 13 | 8 | 2 | 10 |
| 28th September 1925 | 16th October 1925 | 21 | .. | 5 | 4 | 1 | 5 |
| 2nd October 1925 . | 16th October 1925 | 22 | .. | 25 | 3 | 10 | 13 |
| 2nd October 1925 . | 7th October 1925 . | 23 | .. | 6 | 1 | 3 | 4 |
| 2nd October 1925 . | 5th October 1925 . | 24 | .. | 6 | 5 | .. | 5 |

Parasites refused to lay eggs after the middle of October.

* Egg counts taken in first six cases only.

TABLE XXXVI

*Reproductive capacity of Microbracon lefroyi at Surat**(Parasites obtained locally.)*

1924-27

| PERIOD OF OBSERVATION | | Pair No. | No. of pupae | No. of males | No. of females | Total adults | REMARKS |
|-----------------------|---------------------|----------|--------------|--------------|----------------|--------------|---------------|
| From | To | | | | | | |
| 27th October 1924 . | 17th November 1924 | 1 | 35 | 6 | 21 | 27 | |
| 30th October 1924 . | 4th November 1924 | 2 | 4 | .. | 4 | 4 | |
| 30th October 1924 . | 13th November 1924 | 3 | 82 | 36 | 26 | 62 | |
| 5th November 1924 . | 24th November 1924 | 4 | 48 | 12 | 23 | 35 | |
| 5th November 1924 . | 22nd November 1924 | 5 | 35 | 26 | .. | 26 | |
| 13th November 1924 | 6th December 1924 | 6 | 141 | 36 | 58 | 94 | |
| 15th November 1924 | 23rd November 1924 | 7 | 12 | 4 | 4 | 8 | |
| 15th November 1924 | 3rd December 1924 | 8 | 131 | 59 | 44 | 103 | |
| 16th November 1924 | 24th November 1924 | 9 | 30 | 23 | .. | 23 | |
| 24th November 1924 | 28th November 1924 | 10 | 8 | 5 | .. | 5 | |
| 26th November 1924 | 13th December 1924 | 11 | 18 | 8 | .. | 8 | |
| 26th November 1924 | 18th December 1924 | 12 | 52 | 42 | .. | 42 | |
| 26th November 1924 | 18th December 1924 | 13 | 41 | 31 | .. | 31 | |
| 29th January 1925 . | 12th March 1925 . | 14 | 125 | 35 | 64 | 99 | Eggs laid 310 |
| 24th February 1925 | 19th March 1925 . | 15 | 38 | 13 | 15 | 28 | Do. 95 |
| 30th April 1925 . | 21st May 1925 . | 16 | | not noted | | | Do. 504 |
| 25th November 1926 | 10th January 1927 . | 17 | .. | 70 | 66 | 136 | |
| 25th November 1926 | 1st January 1927 . | 18 | .. | 40 | 50 | 90 | |
| 28th November 1926 | 8th January 1927 . | 19 | .. | 37 | 59 | 96 | |
| 2nd January 1927 . | 22nd February 1927 | 20 | .. | 68 | 54 | 122 | |

The last two were mated with the Punjab males to see if they interbreed.

TABLE XXXVII

Reproductive capacity of the Punjab Microbracon lefroyi in Surat in 1926-27

| PERIOD OF OBSERVATION | | Pair No. | No. of males | No. of females | No. of adults |
|-----------------------|----------------------|----------|--------------|----------------|---------------|
| From | To | | | | |
| 13th December 1926 . | 30th December 1926 . | 1 | 27 | 26 | 53 |
| 15th December 1926 . | 10th January 1927 . | 2 | 29 | 40 | 69 |
| 23rd December 1926 . | 29th January 1927 . | 3 | 45 | 56 | 101 |
| 23rd December 1926 . | 31st January 1927 . | 4 | 54 | 74 | 128 |
| 1st January 1927 . . | 5th January 1927 . | 5 | .. | 5 | 5 |

The eggs are laid externally on the host and within 20 to 30 hours they hatch out and begin to feed on the paralysed caterpillar.

LARVAL AND PUPAL PERIOD

The larval period consists of 2 to 6 days. It takes 5 to 8 days for the adults to emerge from the pupae. The whole life-cycle thus covers from about 8 to 13 days.

TABLE XXXVIII

Period of total life-cycle of Microbracon lefroyi during different months

| Months | No. of cases observed | Period of life-cycle in days (Average) | Maximum period of life-cycle in that month |
|-------------------------|-----------------------|--|--|
| November 1926 | 18 | 10 | 11 |
| December 1926 | 45 | 12 | 14 |
| January 1927 | 76 | 13 | 16 |
| February 1927 | 45 | 11 | 13 |
| March 1927 | 53 | 9.5 | 10 |
| April 1927 | 16 | 7.5 | 8 |

METHOD OF REARING THE PARASITES IN THE LABORATORY

A flat piece of glass, about 6 to 8 inches square was taken and a piece of blotting paper of similar size was placed on it. A round glass jar about 4 inches in diameter and about 5 inches in height was inverted and placed on this blotting paper. The adult parasites were liberated in the jars, and fully grown *Earias* larvae feeding in flower-buds, bolls or pieces of *bhendi* were inserted. A piece of blotting paper was placed on a small piece of glass (1 inch square) and the blotting paper was wetted with solution of sugar or honey. These pieces were placed inside the jars for providing food for the adult parasites. Fresh solution was offered twice or thrice a day. The host larvae along with their food were removed every day, and were placed in small sample tubes for recording the emergence of the parasites.

The food of the host larvae usually gave out a lot of moisture which used to accumulate on the sides of the glass cages. Calcium chloride in small sample tubes, closed with cotton wadding, was used for absorbing the moisture.

INCIDENCE OF *Microbracon lefroyi* AT SURAT

The Spotted Boll-worms begin to appear on the new cotton crop generally in August, and during September and October a large number of them are found feeding on the cotton shoots. *Microbracon lefroyi*, however, does not appear till flower-buds begin to develop, late in October or early in November. Thousands of shoots attacked by *Earias* larvae were examined but only one larva parasitised by *Microbracon lefroyi* was found in shoots. It appeared that the parasite was incapable of attacking the *Earias* larvae tunnelling into the cotton shoots. Larvae feeding in cotton shoots were offered to this parasite in the laboratory. The result was that it did parasitise the larvae feeding in tender shoots but not the ones which were feeding in comparatively harder internode. The probable explanation of the immunity of the *Earias* larvae feeding on the shoots in the cotton fields seems to be that it is only the younger larvae which feed on the tender parts of the shoots while the older ones, which are selected by the parasite for oviposition, generally penetrate the shoot by making a hole at a lower node where the shoot is very much harder.

These parasites increase in numbers after the month of October, and in December and January most of the larvae, in the cotton fields, which have reached the parasitisable stage, are found attacked. During 1928-29 a large number of *Earias* larvae were collected from the shed forms gathered from the cotton fields and the number of the caterpillars of the parasitisable age was recorded; as also the number of the larvae which were attacked by this parasite. It was seen that during the first fortnight of December, 66 per cent of the big larvae were parasitised by *Microbracon lefroyi*, whereas during the second fortnight of December this percentage increased to 90. In January the proportion of the fully grown larvae decreased very considerably, and the few big larvae which could be collected were all parasitised by *Microbracon lefroyi*.

TABLE XXXIX

Larvae parasitised by Microbracon lefroyi out of the total larvae of the parasitisable size collected from shed forms in the cotton fields

| Period of observation | Total No. of larvae collected | Larvae big enough to be parasitised by <i>M. lefroyi</i> | Larvae actually parasitised by <i>M. lefroyi</i> | Percentage of the parasitised larvae out of the big ones |
|--|-------------------------------|--|--|--|
| 1st fortnight of December 1928 | 128 | 21 | 14 | 67 |
| 2nd fortnight of December 1928 | 157 | 10 | 9 | 90 |
| January 1929 | 178 | 5 | 5 | 100 |
| February 1929 | 22 | 2 | 2 | 100 |

It is interesting to note that *Earias* larvae feeding in pieces of *bhendi* pods were freely parasitised by this wasp in the laboratory but the *Earias* larvae feeding on *bhendi* plants were never attacked by this parasite, even during November, December and January when the parasite was very common in the cotton fields. A large number of big larvae could thus be obtained from *bhendi* pods, at a time when they were very rare on cotton plants.

No explanation can be offered why this parasite, as well as *Actia aegyptia*, does not attack the *Earias* larvae feeding on *bhendi* plants. The fact, however, supports the conclusion that the cultivation of *bhendi* must be considered to be harmful to cotton, as it allows the rapid multiplication of this pest undisturbed by these two useful parasites.

In the cotton fields at Surat this parasite is present throughout the cotton-growing season except during the first two or three months, *viz.*, July, August and September.

Microbracon lefroyi D. AND G. IMPORTED FROM KHANDESH, PUNJAB AND KARNATAK.

It has been shown that *Microbracon lefroyi* appeared in the cotton fields at Surat in October or November. It was considered that if this parasite were present in the cotton fields a little earlier, or its numbers in October and November could be artificially increased by liberating a large number of parasites bred in the laboratory, the rapid increase of the *Earias* population on cotton in October and November could probably be effectively checked. It was known that this parasite appeared very much earlier in Khandesh and in the Punjab. It was decided to get specimens from these places and try to breed them at Surat.

Microbracon lefroyi FROM KHANDESH

A few of these parasites were brought from Khandesh in August 1925, for multiplication at Surat. They bred normally until October but refused to lay eggs afterwards. (Table XXXV.)

Live material was again secured from Khandesh in 1925, 1926 and in 1927. During 1925 and also in 1926, this was received at Surat in October and practically no eggs were laid after arrival. In 1927, the parasites were received in September and a few eggs were laid by them in that month. The adults from these eggs, however, turned out to be all males. Thus during all the four years the breeding of this parasite from Khandesh did not continue beyond October at Surat. These parasites did not interbreed with the local parasites or those from the Punjab.

Microbracon lefroyi FROM THE PUNJAB

Consignments of *Microbracon lefroyi* were received from the Punjab during August both in 1925 and 1926 through the courtesy of the Government Entomologist, Punjab. These parasites bred quite successfully in captivity at Surat. During both the seasons it was found that by April their oviposition declined very considerably, and the few eggs which were then laid all developed into males.

TABLE XL

Rearing of Microbracon lefroyi from the Punjab at Surat during different months in 1925 and 1926

| Month of the emergence of the female | No. of pairs | PROGENY. | | | No. of females which did not lay eggs | REMARKS |
|--------------------------------------|--------------|----------|---------|-------|---------------------------------------|--|
| | | Males | Females | Total | | |
| August 1926 . | 1 | 20 | 7 | 27 | 1 | The progeny is from only one pair received from the Punjab. |
| September 1926 . | 9 | 75 | 31 | 106 | 6 | |
| October 1926 . | 26 | 463 | 155 | 618 | 2 | |
| November 1926 . | 42 | 654 | 120 | 774 | 27 | From November onwards a large number of females and also males were taken out for various other experiments. |
| December 1926 . | 15 | 120 | 45 | 165 | 6 | |
| January 1927 . | 10 | 140 | 32 | 172 | 5 | |
| February 1927 . | 17 | 127 | 56 | 183 | 6 | |
| March 1927 . | 5 | 18 | 5 | 23 | 6 | |
| April 1927 . | 1 | 3 | .. | 3 | 2 | |

Microbracon lefroyi FROM KARNATAK

A few of these parasites were secured from Karnatak. They could also breed at Surat quite satisfactorily, but it was not possible to get them very early in the season as the cotton crop in Karnatak is sown even later than the crop at Surat.

It was observed that these parasites from the Punjab, Karnatak and Gujarat could interbreed without any difficulty.

Specimens of these parasites from all the localities, viz., the Punjab, Khandesh, Gujarat and Karnatak were sent to Waterston for identification. He reported them all to be *Microbracon lefroyi* D. and G. Ramkrishna Iyer from Coimbatore, however, was of opinion that the Punjab ones might be slightly different, but there was no difference between the Surat and Khandesh ones. He attributed the apathy of the Khandesh ones to breed at Surat to different climatic conditions.

Microbracon REARED FROM OTHER SOURCES

A few specimens of *Microbracon* were reared from *Earias* larvae feeding on *Abutilon indicum* and from shoot rollers (*Phycita infusella* Meyr.) feeding on cotton shoots. A few more specimens were also collected from brinjal (*Solanum melangona*) on bud-worms (*Phthorimæa blaspisigona*). These were also sent to Waterston who identified them as under:—

Parasites reared from *Earias* larvae from *Abutilon indicum* were *Microbracon lefroyi* D. and G.

Parasites from *Phycita infusella* were *Microbracon brevicornis* and

Parasites from brinjal bud-worm were *Microbracon gelechediphagus*.

ATTEMPTS AT CONTINUOUS REARING OF LOCAL PARASITES

The trials of importing *Microbracon lefroyi* from distant localities showed that the parasites from Khandesh were not useful as they could not breed at Surat beyond October. In Karnatak they appeared very late so it was not possible to get the earliest supply of these parasites from that locality. It was possible to secure small consignments of *Microbracon lefroyi* from the Punjab, as early as August, and it was also found possible to breed them successfully at Surat. It was clear that large consignments of these parasites must be available to be able to liberate a very large number of them in the cotton fields in September or October. This was not, however, considered to be very practicable. An attempt was, therefore, made to find out if a large stock of the local *Microbracon lefroyi* could be carried from one season to another. Continuous rearing of the local *Microbracon lefroyi* was, therefore, arranged in the laboratory.

CONTINUOUS REARING OF LOCAL *Microbracon lefroyi* AT SURAT

A few pairs of these parasites were kept in separate jars for rearing. One of the difficulties which was experienced in this rearing was that after an interval

of a few days the proportion of males used to increase in the total progeny. A few trials showed that this was the result of breeding together of closely related parents, which naturally occurred under laboratory conditions of rearing.

TABLE XLI

Proportion of males and females in the progeny of parents closely related and of those which were not related.

| | Proportion of males to every 100 females in 1st genera- tion. | Proportion of males to every 100 females in 2nd genera- tion. | Proportion of males to every 100 females in 3rd genera- tion. |
|--|---|---|---|
| Result of the breeding of closely related parents . | 50 | 79 | 566 |
| Result of the breeding of parents not related to one another. | 50 | 58 | 32 |

MASS REARING

An attempt was made to devise suitable cages for mass rearing of these parasites. They were, however, not very successful.

Another method tried was the liberation of a large number of the caterpillars of the proper size on the cotton plants, enclosed in big muslin cages, along with the adult parasites. In one case 107 big *Earias* larvae were placed on four cotton plants bearing flower-buds and bolls in December 1926, and 20 females and a few males of *Microbracon lefroyi* were liberated inside. A wooden trench with water and oil was kept all round to prevent the escape of the *Earias* larvae. A week later, it was found that seventy-three of these larvae were drowned in the trench, 17 were seen feeding on the plants, 8 pupated, one was found paralysed, and there was only one which was found attacked by the parasites. This method was repeated on two more occasions without any better results. The adults were found always sitting on the thin cloth, more anxious probably to gain their liberty than to oviposit under those conditions.

REARING AFTER MAY

The rearing of these parasites in separate rearing cages could be continued till the month of May. The oviposition by this time declined considerably and most of the eggs which were laid in this month developed into males. Besides this, at about this time the cotton crop was over, and difficulty of supplying host larvae to these parasites was badly felt. Pink Boll-worm (*Platyedra gossypiella* Saund.) larvae which were found in the resting stage, and also a few caterpillars of *Jowar*

stem borer (*Chilo simplex*) which were found hibernating in *Jowar* stalks, were therefore offered. The parasites did lay their eggs on both kinds of caterpillars, but the larvae emerging from the eggs failed to develop on these hosts.

CARRY-OVER OF THE PUPAE AND THE ADULTS OF *Microbracon lefroyi* IN COLD CHAMBERS

To see if the life of these parasites could be prolonged so as to get them to reach the next cotton season safely, a few pupae, which had passed 5, 7 and 9 days after pupation, were kept in the first four compartments at the cold end of the multiple temperature incubator (App. III), during February 1927.

TABLE XLII

Maximum period of pupation in the different compartments after the pupae were placed in them

(10 pupae in each compartment)

| | 1st compartment temperature 48°F. to 54°F. | 2nd compartment temperature 60° to 64°F. | 3rd compartment temperature 64° to 68°F. | 4th compartment temperature 70° to 76°F. |
|--|--|--|--|--|
| Maximum number of days for emergence from the five days old pupae. | 33 | 11 | 9 | 3 |
| No. of adults emerged from the above. | 6 | 7 | 9 | 10 |
| Maximum number of days for emergence from 7 days old pupae. | 11 | 9 | 5 | 4 |
| No. of adults emerged from the above. | 6 | 9 | 7 | 6 |
| Maximum number of days for emergence from 9 days old pupae. | 13 | 3 | emerged on the day they were kept. | |
| No. of adults emerged from the above. | 8 | 9 | 10 | 10 |

The parasites emerging from these pupae could breed normally afterwards in the laboratory.

Similarly 18 adults (10 males and 8 females) were kept in each of the above four compartments for noting the total period of their life. They were fed with sugar solution every day.

TABLE XLIII

Length of life of adult Microbracon lefroyi under continuous low temperatures of the first four compartments

(See Table XLII)

| — | First compartment | Second compartment | Third compartment | Fourth compartment |
|--|----------------------|-----------------------|----------------------|-----------------------|
| Maximum No. of days any of the adults lived. | More than 84 | 71 | 64 | 45 |
| Minimum No. of days any of the adults lived. | 30 | 25 | 3 | 2 |
| Average life of the adults . . . | 44 | 38 | 31 | 14 |

Only six of these parasites lived for more than 84 days in the first compartment. These were then removed for observing the capacity of oviposition. It was, however, found that all of them died without laying any eggs. Besides this, 3 pairs of parasites were kept for egg laying in each of these four compartments. All of them laid eggs except the ones kept in the first chamber where the temperature varied between 48° and 55°F.

These experiments showed that it was possible to prolong the life of the pupae as well as the adults considerably under low temperatures, but the procedure was impracticable under ordinary conditions; and besides there were indications that the adults which survived long enough would probably not breed normally.

An attempt was also made to keep the parasites alive in the pupal stage by keeping them at a very low temperature. A small cold chamber was, therefore, improvised by keeping a small round tin immersed in ice cold water continuously. The temperature inside this tin varied between 34 and 42°F. Forty *Microbracon* pupae were placed in this tin in a glass tube. Twenty of them were removed after 12 days and 4 adults emerged from this lot. The remaining 20 pupae were removed after 34 days and it was found that all of them were dead by that time. Thus it appeared that the pupae would not stand very low temperatures for a considerable length of time.

Finally, therefore, it must be admitted that after making a large number of efforts no easy and cheap method could be arrived at of securing a large stock of these parasites for making an attempt at augmenting the natural population of *Microbracon lefroyi* in the early part of the cotton-growing season.

Melcha nersei CAM. (PARASITE ON THE PUPAE OF *Earias*)

Out of the remaining four parasites, only *Melcha nersei* Cam. calls for comment.

Husain and Mathur [1923] have mentioned that these big parasites are seen flying about in the fields of cotton, lucerne, *senji* and in the gardens on hollyhocks in the Punjab. But they are not so commonly seen at Surat and it was only on a couple of occasions that their presence was noticed in the cotton fields. The marked absence of this parasite at Surat appears to be due to the fact that the *Earias* larvae invariably go to the soil for pupation in this tract and the pupae are not, therefore, exposed to the attack of this parasite. In the Punjab, on the other hand, a large number of *Earias* pupae are probably to be found on the parts of the cotton plants above the soil.

Only one female of this parasite was bred from a few pupae, collected from *bhendi* plants (*Hibiscus esculentus*) from Adajan, a small village near Surat, on 15th October 1924.

This female was kept in a glass rearing jar and *Earias* pupae were offered to it for oviposition. From the pupae, which were attacked, the first male emerged on the 31st of October and it was introduced in the cage of this female wasp. The female wasp escaped on 4th November 1924. During this interval we had offered a total of 176 *Earias* pupae, out of which 55 *Earias* moths emerged. Eight female and 37 male parasites emerged from the remaining 121 pupae.

A few of the off-spring of this parasite were kept under observation for recording the life-history during November and December at Surat.

Pair No. 1.—The female and male lived for 20 and 22 days respectively. During this period 168 *Earias* pupae were offered (ten fresh pupae were offered on most of the days). From these pupae 40 *Earias* moths and 12 female and 56 male parasites emerged. The period between the day on which the *Earias* pupae were attacked and the day on which the parasites emerged varied between 15 to 22 days.

Pair No. 2.—The female lived for 26 days and the male for 11 days. In all 222 *Earias* pupae were offered to the female. From these, 50 *Earias* moths and 82 male parasites emerged. The progeny consisted of males only, although a male parasite was present in the female cage for nearly 10 days and they were seen to mate. The period from egg to adult in this case varied between 15 to 24 days.

4. DETERRENTS

Attempts were made to find out if flowers of sulphur and cynogas dust (calcium cyanide) would be of any use as deterrents for the *Earias* moths.

A large number of cotton plants were dusted with flowers of sulphur mixed with road dust in the last week of December of 1923, and it was decided to keep

SPOTTED BOLL-WORMS OF COTTON IN SOUTH GUJARAT

these plants under observation. After about 4 days, however, a very large number of flower-buds, young bolls and leaves were found to be dropping down from the dusted plants. The shedding continued for some time, and after a few days only a few stray leaves and developed bolls remained on these plants.

This experiment was repeated in the last week of February 1924, when a few plants of cotton were dusted with flowers of sulphur alone. This was also followed by shedding of forms, as in the above case.

Not only did the dusted plants suffer but heavy shedding of leaves, flower-buds and young bolls occurred on a large number of neighbouring plants over which the sulphur fumes were carried by wind.

These trials and subsequent experience of other workers [Thakar and Desai, 1929] showed that the plants of herbaceous cottons at Surat were extremely susceptible to the slightest traces of sulphur, and it was dangerous to expose the plants to any substance which was likely to give out even traces of free sulphur.

Cynogas dust (calcium cyanide A, supplied by Messrs. Shaw Wallace and Co., Bombay) was another substance which was tested for keeping the *Earias* moths from the cotton plants. During October 1926 the shoots of 200 plants were lightly dusted with this powder. The tender leaves of these plants were considerably scorched by it. An attempt was, therefore, made to spread a small quantity of this powder on the soil near stems of cotton plants, instead of applying it to them directly. Even this change did not prevent the scorching of the leaves, and hence the attempt to use this material as a deterrent was given up.

5. ATTRACTANTS

It had been observed that a few *Earias* moths came to the lights in the residential quarters during January and February. It was found that ordinary Dietz lanterns as well as acetylene lamps exposed in the fields failed to attract any of these moths during the period of the heaviest infestation.

Aromatic substances such as, amyl acetate, methyl butyrate, acetic acid, formaldehyde, clove oil, molasses with yeast, vinegar, turpentine, rum, country liquor were exposed in the cotton fields. They were placed in small enamel dishes, which were kept in the centre of two feet square pans, containing water. None of these substances showed any attraction for these moths.

In the meanwhile we found that Power and Chestnut [1925] had determined the odorous constituents of cotton plants and had shown that ammonia and trimethylamine were present in appreciable quantities in the distillate obtained from these plants, and that they were also found in the emanations of the living cotton plants.

Ammonia, ammonium carbonate and rotting fish were, therefore, exposed, in bait pans. Rotting fish were used instead of trimethylamine as the smell of both the substances is said to resemble very closely. These substances also did not attract the *Earias* moths.

Attention was, therefore, diverted to find out if the cotton seeds, at any stage of their decomposition, would be able to attract the moths. A small quantity of seed was, therefore, moistened and was exposed in the same way as the substances mentioned above. The seed was again moistened at an interval of every 3 or 4 days. It was found that on some of the days a few moths of the Spotted Boll-worms could be found in the water pans. The attraction, however, was not constant and it appeared that very moist or very dry seeds were not effective in attracting the moths (App. I, Table XXVIII).

Bhendi (*Hibiscus esculentus*) seed was crushed and moistened and was similarly exposed in the cotton fields and the observations showed that it proved to be slightly superior to cotton seed (App. I, Table XXVIII).

It was not possible to say if it was the decomposition of the seeds of the host plants, which caused the attraction or if any similar substance would also be able to attract the moths. By analogy of the composition of these substances, therefore, moistened sesamum cake was exposed in the bait pans. The moths of the Spotted Boll-worms did respond to it, and it was found that the number of moths attracted to the cake pans, was more than that at the cotton or *bhendi* seeds. On the 31st of January 1928, as many as 38 moths were found round a single pan of sesamum cake, while in the pans of cotton and *bhendi* seeds there were only one and three moths respectively on that day (App. I, Table XXVIII).

The cakes of groundnut and cotton, in addition to that of sesamum, were tested and it was found that all three could attract a small number of *Earias* moths. Sufficient water was added to the powder of these cakes to form a paste before exposing them in the enamel dishes. Later on, however, it was found that these substances showed a better attraction when about 2 or 3 ounces of their powder was added directly to the bait pans containing about 2 or 3 gallons of water, instead of only moistening the cakes.

A few pans with these substances were exposed during most of the cotton growing season in 1929-30 and 1930-31, both at Surat and at Jalgaon in East Khandesh district (App. I, Tables XXIX to XXXII). The number of moths attracted to these pans at Jalgaon was considerably greater than at Surat.

It was rather peculiar that the largest number of moths were collected from these pans when the cotton crop was about to be over both at Surat and Jalgaon. Probably it meant that the attraction of these baits could only be strong enough to attract the moths in the absence of the vigorously growing crop of cotton, having tender shoots, flower-buds and young bolls.

It is interesting to note that these baits not only attracted the moths of the Spotted Boll-worms but also of the Pink Boll-worm (*Platyedra gossypiella* Saund.) and the shoot roller (*Phycita infusella* Meyr.). A very large number of Pink Boll-worm moths were attracted to these pans at Jalgaon (App. I, Tables XXXI and XXXII).

Paste of sandal wood and nutmeg, prepared by rubbing these substances on a rough stone with little water, also attracted a few moths of the Spotted and Pink Boll-worms.

The number of moths attracted to any of these substances was not sufficiently large to warrant the use of any of them as a control measure. The trials may, however, indicate possibilities of evolving a substance which may prove to be a powerful bait, not only for the Spotted Boll-worms, but also for the allied pests of cotton.

6. SOIL MULCH AS A CONTROL MEASURE

A careful search in the cotton fields at Surat showed that it was extremely rare to find a pupa of the Spotted Boll-worms on the plants. This fact indicated that the larvae which are about to pupate, moved away from the cotton plants for spinning their cocoons. An examination of the soil, round the cotton plants showed that they were to be found in the soil, from about 2 to 8 inches below the surface of the soil.

It has already been mentioned that the fully fed larvae enter cracks and crevices for finding a suitable place for pupation. Soon after the close of the monsoon the soil of the locality opens into small cracks, which slowly widen and have several sub-cracks, going in all directions from them. These cracks allow very easy access for the larvae to the soil and the pupae are often found in such small cracks, by the side of small clods (Plate I, Fig. 1). It has also been stated that the full-fed larvae descend the plants in the forenoon and immediately begin to search for a place for pupation. By about 2 p.m., most of them are found to have begun the work of spinning their cocoons. This happens to be the hottest part of the day, and it is natural that the larvae should try to take shelter in the cooler layers of the soil.

The facts that the larvae almost invariably go to the soil for pupation in this tract, that they have to enter the soil through the existing cracks and crevices, that the moths, when they emerge have to make their way out of the soil through similar cracks and crevices, and that the larvae choose the hottest part of the day for selecting a place for pupation, all pointed out that this was likely to be a vulnerable point in the life of the Spotted Boll-worms.

What was required for successfully attacking the worms at this stage was to prevent them by some means from going deep in the soil, thus exposing them to the heat of the sun and to the attack of their predators; or to prevent the emergence of the moths from the pupae in the soil. It was thought that this would be possible by maintaining a fine soil mulch in the cotton fields, and therefore trials were undertaken for testing the effect of soil mulch on the pupation and the emergence of moths of the Spotted Boll-worms.

The preliminary laboratory trials consisted of observing the emergence of moths from the pupae, which were covered over with a shallow layer of loose soil. Three wire gauge cages (one foot square) were taken and were filled in with field soil, which contained a few clods also. Twenty full-grown larvae were liberated in each one of them and they were allowed to pupate in the soil. The first cage was reserved as a control. In the other cages a layer of one inch and two inches of loose soil was added. The emergence of moths from these cages was noted and later the soil was examined to find out if any of the moths were caught in the loose soil.

TABLE XLIV

Emergence of the moths from the above three cages

(20 larvae liberated in each case)

| Cage No. | No. of larvae pupated in the soil | Moths emerging above the soil | Moths dead in the soil | Percentage mortality |
|---------------------------------------|-----------------------------------|-------------------------------|------------------------|----------------------|
| No. I, Control | 19 | 19 | Nil | Nil. |
| No. II, 1 inch layer of loose soil . | 18 | 6 | 12 | 72 |
| No. III, 2 inches layer of loose soil | 13 | Nil | 13 | 100 |

All the moths in the first case came out normally; in the second case, however, 72 per cent of them could not extricate themselves from the loose soil and were stifled below the surface of the soil. In the third case all the moths died in the soil and were found with their wings crumpled up, when the soil was examined.

These trials only showed that if there is a small layer of loose soil over the pupae, most of the moths would fail to emerge. It was yet to be ascertained as to where the larvae would pupate, if a good soil mulch was maintained all over the cotton fields. Information on the following points was considered to be essential for determining the efficiency of the soil mulch as a control measure :—

- (1) Will the larvae pupate on the surface of the soil if the area is mulched with fine soil ?

- (2) Can they successfully pass all their pupation period in the above case and be able to emerge as moths normally ?
- (3) Is the temperature of the surface layer of the soil high enough to kill the pupae ?
- (4) Can the implements pulverise all the soil right up to the collar of the plants ? If not, what proportion of the larvae would pupate in the untreated space ?

A trial in the laboratory showed that when 20 full-grown larvae were liberated in a wire-gauge cage, containing pulverised soil (passed through a sieve with 16 meshes to a square inch) 17 of them pupated in the surface layer of the soil by binding together a few particles of the soil. Moths emerged normally from all these pupae.

This trial was later repeated in the fields, where 106 larvae were liberated in a small area (6 feet square) surrounded by a tin trench, containing water and oil. The temperature of the surface layer of the soil in the above space was found to be 108° F. at 2 P.M. on that day. Most of the larvae tried to escape from the enclosed space as they could not get a suitable place for pupation, and 58 of them were drowned in the trench in trying to go beyond. Later, when the soil was examined, no pupa could be traced and therefore it appeared that the remaining larvae must have been killed by the heat or were destroyed by their predators.

In another case 50 fresh pupae were scattered in the surface layer of an inch of the soil during the month of September. They were allowed to be there for a period of one week. An examination of the soil then gave a recovery of 31 cocoons out of the total of fifty. Fourteen of these were almost completely eaten up by predators, and 12 more were partially damaged. Only five pupae were found alive. Thus out of the fifty, only 31 could be secured and the remaining 19 were entirely missing. They were either devoured completely or were bodily removed by their enemies.

In the meanwhile, to study the effect of heat on the pupae, a few of them were subjected to different temperatures in the multiple temperature incubator (App. III). It was found that they could not stand a continuous temperature of more than 100°F., as all the pupae which were placed in the chambers with temperatures ranging above 100°F. failed to emerge as moths and were all found dried up.

Such continuous temperatures of about 100°F. or more are not naturally available in the fields. The soil temperature in the fields is the hottest between 12 noon and 4 P.M. Surface temperatures of the soil recorded at 12 noon, 2 P.M. and 4 P.M. once in a week, during October, November and December 1926, are given below for showing the range of temperature during the hottest part of the day.

TABLE XLV

Soil temperatures of the surface 1 inch layer in cotton fields

(Fahrenheit-1926)

| Date of recording | Mid-day | 2 P.M. | 4 P.M. |
|-------------------------|---------|--------|--------|
| 9th October | 104 | 108 | 106 |
| 16th October | 101 | 106 | 105 |
| 23rd October | 105 | 112 | 108 |
| 30th October | 105 | 111 | 109 |
| 6th November | 100 | 106 | 103 |
| 13th November | 109 | 110 | 109 |
| 20th November | 103 | 112 | 106 |
| 27th November | 100 | 108 | 103 |
| 4th December | 101 | 114 | 112 |
| 11th December | 100 | 111 | 108 |
| 18th December | 100 | 109 | 102 |
| 25th December | 100 | 109 | 105 |

These observations indicated that temperatures between 100° and 110°F. are available for a few hours during the day in October, November and December, in the surface layer of the field soil. The effect was then examined of exposing the pupae to about 110°F. for varying periods in the multiple temperature incubator. The emergence of moths from these pupae was noted.

TABLE XLVI

Emergence of moths from pupae exposed to 110°F.

| No. of hours the pupae were exposed | Number of pupae under observation | No. of moths emerged |
|-------------------------------------|-----------------------------------|----------------------|
| 4 | 10 | 9 |
| 8 | 10 | 7 |
| 24 | 10 | 5 |
| 48 | 10 | Nil |
| 72 | 10 | Nil |

} All dried up.

A few of the pupae could endure the temperature of 110°F. for about 24 hours, but all those which were kept at that temperature for 48 hours or more, were shrivelled up and dead due to the heat.

Attempts were next made to find out if it would be practicable to secure the desired soil-mulch with different implements. The blade harrow was found to be the best implement for pulverising the soil when it was used in the optimum condition of the soil. All the implements, however, left a small space of a few inches undisturbed near the collar of the plants. It was clear that some of the larvae would take advantage of this unmulched area for pupation. Further trials of artificially liberating the larvae in the fields showed that when a space of about 6 inches round each plant was left untreated, 20 per cent of the larvae (16 out of 80) pupated in this space. When this space was reduced to 2 inches, 10 per cent (10 out of 102) of the larvae pupated in that small space.

The last few trials indicated that the presence of well pulverised soil in the fields would prevent the larvae from going below the surface layers if the soil could be mulched right up to the collar of the plants and that some of these, which would pupate in the surface layer, would be destroyed by the heat of the sun or by the predators.

In order, therefore, to get a definite idea about the efficiency of the soil-mulch as a control measure, it was decided to lay out small plots with and without mulch, and to observe the natural pupation under those conditions.

Four plots of cotton, about 4 or 5 *gunthas* each, were selected at the commencement of the cotton-growing season during 1928-29. One of these plots received all the usual tillage operations as in the case of the whole cotton field, and was reserved as a control. All the soil in the second plot was carefully pulverised, and fine soil-mulch was maintained in this plot throughout the season by stirring the soil repeatedly, and by breaking the clods which came up. Similar soil-mulch was maintained in the third plot, but a space of about 1 inches round each plant was left untreated in that plot. No cultural treatment was given to the fourth plot and the soil was allowed to crack to the maximum extent. The soil around 9 plants (plants spaced 4'×3') was carefully examined in each one of these plots, once a month, for pupae of the Spotted Boll-worms (Tables XLVII to XLIX).

During the next season, 1929-30, only 3 plots were laid out :

1st plot.—Control. Normal tillage.

2nd plot.—With good soil mulch, except 4" round each plant.

3rd plot.—With clods. The clods were purposely turned up and were not pulverised in order to have a very good contrast with the plot having well pulverised soil.

Soil round 20 plants (spaced 4'×3') was examined once in a month from each one of these plots.

The plot with the soil-mulch going right up to the collar of the plants was not attempted as it was found impracticable to pulverise the soil with any implement without leaving at least some space untreated round the plants.

The soil was removed for observation by separating out small clods and these were carefully examined. The positions of the pupae found in the soil were recorded. The loose soil was again passed through sieves to prevent the loss of any of the pupae through oversight.

TABLE XLVII

Depth at which the cocoons were found in the soil

| Depth in inches | 1928-29 | | | | 1929-30 | | |
|------------------------------|--------------|------------------|--|--------------------|--------------|----------------------------|---|
| | Control plot | Plot with cracks | Plot fully mulched leaving 4" round plants | Plot fully mulched | Control plot | Plot with clods and cracks | Plot fully mulched except 4" round plants |
| 1 | 14 | 14 | 18 | 35 | 13 | 3 | 10 |
| 2 | 5 | 8 | 6 | 5 | 9 | 10 | 8 |
| 3 | 2 | 2 | .. | 2 | 7 | 23 | 9 |
| 4 | 1 | 2 | .. | 1 | .. | 12 | 2 |
| 5 | 1 | 2 | .. | .. | .. | 5 | .. |
| 6 | 1 | 1 | 1 | .. | 4 | 3 | 5 |
| 7 | 1 | 3 | 2 | .. | 5 | .. | 8 |
| 8 | .. | 1 | 1 | .. | .. | 1 | .. |
| 9 | .. | 1 | 1 | .. | 2 | 1 | 1 |
| 10 | 1 | 1 | .. | .. | .. | 1 | .. |
| 11 | .. | .. | .. | .. | 1 | 1 | .. |
| 12 | .. | 1 | .. | .. | .. | .. | .. |
| 13 | .. | 2 | .. | .. | .. | .. | .. |
| Cocoons caught in the sieve. | 1 | .. | 2 | 2 | 4 | 3 | 5 |
| TOTAL | 27 | 40 | 31 | 45 | 45 | 63 | 43 |

TABLE XLVIII

Proportion of the cocoons in the surface one-inch layer of the soil.

| Plots | Total cocoons | Cocoons in 1st 1-inch layer | Percentage of the cocoons in surface 1-inch layer |
|--|---------------|-----------------------------|---|
| 1928-29. | | | |
| Control | 26 | 14 | 54 |
| With cracks | 40 | 14 | 35 |
| Mulched leaving 4" round plants | 29 | 18 | 62 |
| Fully mulched | 43 | 35 | 81 |
| 1929-30. | | | |
| Control | 41 | 13 | 32 |
| With cracks and clods | 60 | 3 | 5 |
| With mulch leaving 4" round plants | 38 | 10 | 26 |

The cocoons which were found in these plots were examined and were placed in three classes :—

- (1) Live pupae.
- (2) Damaged cocoons (destroyed by predators).
- (3) Empty cocoons, which showed that moths had emerged out normally from them.

TABLE XLVIX

| Plots | Total cocoons | Live pupae. | Dead or damaged cocoons | Percent-age damaged cocoons | Empty cocoons showing normal emergence | Percent-age empty cocoons |
|---------------------------------------|---------------|-------------|-------------------------|-----------------------------|--|---------------------------|
| 1928-29. | | | | | | |
| Control plot | 27 | 5 | 4 | 14.8 | 18 | 66.6 |
| Plot with cracks | 40 | 8 | 10 | 25.0 | 22 | 55.0 |
| Plot mulched leaving 4" round plants. | 31 | 4 | 11 | 35.5 | 16 | 51.6 |
| Fully mulched plot | 45 | 11 | 15 | 33.3 | 19 | 42.2 |
| 1929-30. | | | | | | |
| Control plot | 45 | 2 | 17 | 37.8 | 26 | 57.8 |
| Plot with cracks and clods | 63 | 5 | 22 | 34.9 | 36 | 57.1 |
| Plot mulched leaving 4" round plants. | 43 | 5 | 20 | 46.5 | 18 | 41.9 |

It was found that the surface of one-inch layer of the mulched plots had comparatively a larger number of cocoons than those which were found in similar layers of the other plots, having cracks and clods. But it was also true that in these plots, some of the cocoons were found in the deeper layers also, in spite of the surface soil-mulch.

Secondly, it was observed that the mulched plots had a larger proportion of damaged cocoons and that in these plots a smaller number of them showed signs of normal emergence, as compared with the other plots. The difference, however, was not substantial and it appeared that moths did emerge successfully from the mulched plots also.

Thus it was found that the pupation in the mulched plots was not restricted only to the thin surface layer of the pulverised soil, and that it appeared that a good number of moths could emerge normally from cocoons in these plots. This was mostly due to the following reasons.

A large number of cracks began to appear in the mulched plots as the soil began to dry up. It was, therefore, necessary to stir the soil very often in the mulched plots for preventing the cracks. In stirring the soil, the fine particles of the soil settled down deeper and small clods came up to the surface. After the surface soil had dried up, it was impossible to pulverise these clods and therefore they provided very good shelter for the pupae in these plots.

It was also found that some of the pupae in the soil were destroyed by predators, such as lizards, which lived in the cracks. They could not, however, be active in the mulched plots in the absence of cracks.

All these observations, therefore, showed that soil mulch, as far as it can be maintained in the field conditions at Surat, could not be considered to be an effective control measure against the Spotted Boll-worms, though the preliminary trials indicated possibilities of its success.

7. INSECTICIDES

Stomach poisons are generally recommended for insects which have a biting habit of feeding. The possibility of this kind of insecticide being effective against the Spotted Boll-worms was doubtful, because they bore into their food and then continued feeding from inside the food. The only chance, therefore, of their getting the poison is while making an entrance hole.

A few preliminary trials in the laboratory, however, indicated that a very high mortality ensued when these worms were fed on flower-buds and bolls which were previously sprayed or dusted with lead arsenate or Paris Green,

TABLE L

Mortality of larvae fed on flower-buds sprayed with lead arsenate and Paris Green

| Poison | No. of worms fed | Mortality on each day after feeding | | | | | | | TOTAL |
|--|------------------|-------------------------------------|----|----|----|----|----|----|------------------------|
| | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | |
| Lead arsenate 2 ozs. to 4 gallons of water. | 100 | 28 | 22 | 28 | 10 | .. | 11 | .. | 99 dead and 1 pupated. |
| Paris Green $\frac{1}{2}$ oz. to 4 gallons of water. | 100 | 54 | 24 | 16 | 6 | .. | .. | .. | 100 |

TABLE LI

Mortality of larvae fed on flower-buds dusted with lead arsenate and Paris Green

| Poison | No. of worms fed | Maximum number of days for which any of these larvae lived | No. of larvae pupated | Percentage mortality |
|--------------------------|------------------|--|-----------------------|----------------------|
| Lead arsenate dust . . . | 50 | 16 | 6 | 88 |
| Paris Green dust . . . | 50 | 14 | 12 | 76 |
| Untreated | 50 | .. | 32 | 36 |

Experiments in the fields were, therefore, undertaken in plots of about half a *guntha* each, for testing the practical efficiency of these insecticides. Both sprayings and dustings were attempted at the interval of one, two and three weeks. The treatments were restricted to the months of November and December and a part of January when the boll-worms were particularly active on the cotton plants. Lead arsenate was sprayed at the rate of 2 ounces for 4 gallons of water and Paris Green at half an ounce to 4 gallons of water. Lead arsenate and Paris Green were

mixed with powdered lime in the proportion of 1 to 8 before dusting. The results of these treatments are tabulated below :--

TABLE LII

Cotton plants sprayed and dusted with lead arsenate and Paris Green.


1924-25

| Treatment | Plants under observation | Flowers per plant | Mature bolls per plant | Percentage flowers to bolls | Kapas per plant in Tolas |
|---|--------------------------|-------------------|------------------------|-----------------------------|--------------------------|
| Check plot | 54 | 86 | 40 | 46 | 7.1 |
| Lead arsenate sprayed once in a week . | 54 | 71 | 31 | 43 | 5.0 |
| Lead arsenate sprayed once in two weeks | 52 | 67 | 33 | 49 | 5.5 |
| Lead arsenate sprayed once in three weeks | 54 | 71 | 33 | 46 | 5.5 |
| Lead arsenate dusted once in a week . | 51 | 82 | 37 | 44 | 6.1 |
| Lead arsenate dusted once in two weeks . | 53 | 75 | 36 | 48 | 5.9 |
| Lead arsenate dusted once in three weeks | 54 | 82 | 37 | 45 | 6.2 |
| Paris Green sprayed once in two weeks . | 54 | 64 | 28 | 43 | 4.1 |
| Paris Green dusted once in two weeks . | 53 | 72 | 31 | 42 | 4.7 |

These trials showed that no advantage could be seen as a result of these treatments at any stage of the plant growth or in the final yields of *kapas*. Paris Green was, on the contrary, found to be harmful as it affected the leaves, to a certain extent, and ultimately reduced the number of bolls and the yield of *kapas*.


In the meanwhile, it was observed that the tiny *Earias* larvae as soon as they come out of the eggs have the habit of wandering about on the plant for some time before they begin to feed. It was considered that these larvae would get the poison very easily if the plants were dusted with a very finely powdered insecticide. It was, therefore, decided to try calcium arsenate which was being extensively used against the boll-weevil (*Anthonomus grandis*) in America. During 1926-27, therefore, three plots of 100 plants each, were laid out for these trials. One of these was dusted with calcium arsenate without any dilution; and in the other, a mixture of calcium arsenate and slaked lime was used in the proportion of 1 to 3. The third plot was reserved as a control. These dustings were started in the last week of November and were repeated at the interval of 5 days till the end of December.

The beneficial results of these applications were soon apparent. The dusted plots showed a number of developed flower-buds within a few days, and the vigorous flower formation reached its height 5 weeks earlier than in the control plot. (Plate XII.) This indicated that the earlier flower-buds in dusted plots had escaped the



Showing periodical flower opening and relative success of bolls from dusted and undusted plants at Surat during 1926-27

PLATE XIII.



Early opening of bolls in plot (left) dusted with calcium arsenate.

damage of the boll-worms and had succeeded into flowers in large proportions (Appendix I, Table XXXIII). It was also found that the earlier flowers were retained on these plants in larger proportions than on the control plants. Consequently a large number of bolls opened on the dusted plants very much earlier than the general field plants. (Plate XIII.)

In order to obtain comparative data upon the number of worms in the dusted and the control plots, 25 plants were examined from each of the 3 plots, 4 weeks after the dustings were started. The results of these examinations are summarised in the next two tables.

TABLE LIII

Examination of 25 cotton plants in each case

| Treatment | Total flower-buds | Total bolls | Live larvae | Dead larvae (very tiny) | Live eggs |
|-------------------------------------|-------------------|-------------|-------------|-------------------------|-----------|
| Calcium arsenate | 2,992 | 139 | 60 | 463 | 596 |
| Calcium arsenate and lime | 2,935 | 128 | 117 | 97 | 205 |
| Control | 2,116 | nil | 123 | nil | 28 |

TABLE LIV

The number of bolls and the proportion of the flower-buds of different sizes from the above plants

| Treatment | Smallest buds | Small buds | Medium buds | Big buds | Bolls |
|-------------------------------------|---------------|------------|-------------|----------|-------|
| Calcium arsenate | 1,364 | 651 | 597 | 205 | 139 |
| Calcium arsenate and lime | 1,845 | 646 | 338 | 106 | 128 |
| Control | 1,711 | 244 | 112 | 25 | nil |

The following points are noticeable in the above tables :—

- (1) The plants from the dusted plots had a few bolls on them at a time when the non-dusted plants had none. Besides they carried a large number of big sized flower-buds.
- (2) Dusted plants had a smaller number of live boll-worm larvae than the control plants.

- (3) A very large number of tiny dead larvae could be seen all over the dusted plants.
- (4) The dusted plants had an unexpectedly high number of live eggs of the Spotted Boll-worms.

It must be noted that the above facts were more markedly seen in the plot dusted with calcium arsenate only than in the other which was dusted with calcium arsenate diluted with slaked lime.

The conclusions which these facts could warrant were that the calcium arsenate was successful in killing a large number of boll-worm larvae, and most of them were killed, just after they had emerged out of their eggs and before they got into their food. This further indicated that the larvae got a dose of the poison while crawling in search of their food on the parts of the plants, which were covered with the dust. It was the large mortality of these larvae which allowed the appearance of better developed flower-buds and bolls, earlier in the dusted plots. The presence of these advanced forms on the dusted plants, in larger numbers than what were to be found anywhere in the surrounding cotton plots, appeared to have attracted a very large number of moths to oviposit on them, with the result that the number of live eggs on the dusted plants swelled to an unexpected extent. This was a very significant fact, because it emphatically pointed out that in spite of the continuous abnormal oviposition, the dust could afford protection to the flower-buds and bolls by keeping the larval population very low. It would probably have been still lower if the oviposition had continued to be normal.

The treatment appeared to be very hopeful, and to promise a large increase in the yield of *kapas*. However, with the commencement of vigorous flowering, a heavy infestation of aphid (*Aphis gossypii*) developed on the dusted plants, and they were soon covered by the sugary secretions of this pest. As a result of this trouble the dusted plants finally did not even yield the normal number of bolls, found on the control plants.

TABLE LV




Dusted and control plants

1926-27

(Average of 20 plants in each case)

| Treatment | Total flowers per plant | Total mature bolls per plant |
|-------------------------------------|-------------------------|------------------------------|
| Calcium arsenate | 87 | 24 |
| Calcium arsenate and lime | 76 | 25 |
| Control | 82 | 31 |

PLATE XIV.



— weekly flower opening
 --- bolls which matured from above flowers

Showing weekly flower-opening and relative success of bolls in cotton plots dusted and undusted with insecticides.

A few more trials were again laid out during the next season (1927-28) to find out, if the aphid infection would recur during every season, and if it would be possible to escape it by some means. The following were some of the treatments which were attempted. They were carried out during the period from the middle of October to the middle of December, which generally happens to be the period of heavy boll-worm damage—

- (1) Dusting with calcium arsenate at the interval of 5 days, from the end of October till the middle of December.
- (2) Dusting as above and 4 sprayings with fish-oil rosin soap, after the appearance of aphid.
- (3) Dusting with calcium arsenate at intervals of 10 days, from the middle of October till the middle of December.
- (4) Two dustings of calcium arsenate at the commencement of the flower-bud development for protecting the flower-buds, and two more at the flowering period for securing success from the earliest flowers.
- (5) Only two dustings of calcium arsenate at the period of flowering.
- (6) Dustings with calcium arsenate diluted with gram flour (in the proportion of 1 of calcium arsenate to 2 of flour) at intervals of 5 days from the end of October till the middle of December.

About 100 plants were treated in each case and 20 plants were selected in the middle of each of the plots for noting the daily flower opening, boll shedding and the final number of mature bolls. These results have been included in Appendix I, Table XXXIV (Plate XIV). Some of the important points seen from these figures are enumerated below :—

- (1) Intense flowering was much earlier in the dusted plots than in the control ones.
- (2) The plots dusted with calcium arsenate at 5 and 10 days intervals respectively had about the same period of flowering but the intensity of flowering in the plot dusted at intervals of 10 days was lower.
- (3) The plot, which received calcium arsenate diluted with gram flour, did not show as much intensity of flowering as the one which received arsenate alone.
- (4) The plot which was dusted with calcium arsenate at intervals of 5 days and was also sprayed with fish-oil rosin soap (against aphid) showed an advantage over the similarly dusted plot, which was not sprayed. It retained proportionately a larger number of flowers, during the weeks which followed the sprayings.
- (5) The plot which was dusted twice only with calcium arsenate showed a remarkable rise in flowering immediately after the commencement of

dustings; and the early flowers gave a higher percentage of success of flowers to bolls than the untreated plots.

These trials, therefore, again pointed out that as a result of the dustings of calcium arsenate, an intense early flowering took place and the early flowers were retained on the plants in larger proportions.

Soon after the commencement of the flowering it was, however, found that during this season also infestation by aphis began to appear, and it was found to be very serious in the dusted plots during December. This affected very considerably the production of *kapas* in the dusted plots as can be seen from the next table. Yields of *kapas* from 8 untreated plots (with about 100 plants in each) selected at random, in the same field, are also shown for judging the normal variation of yield in that field.

TABLE LVI

| Treatments | No. of plants observed | Total yield of <i>kapas</i> in tolas |
|---|------------------------|--------------------------------------|
| 1. Dusting calcium arsenate every 5 days | 99 | 263 |
| 2. Dusting calcium arsenate every 5 days and spraying with fish-oil rosin soap. | 98 | 310 |
| 3. Dusting calcium arsenate with gram flour | 101 | 270 |
| 4. Dusting calcium arsenate twice at flower-bud development and twice at flower-opening period. | 99 | 209 |
| 5. Dusting calcium arsenate at every 10 days | 95 | 236 |
| 6. Dusting calcium arsenate twice only | 93 | 258 |
| 7. Untreated plot | 99 | 540 |
| 8. Do. | 99 | 531 |
| 9. Do. | 98 | 410 |
| 10. Do. | 100 | 497 |
| 11. Do. | 100 | 488 |
| 12. Do. | 100 | 517 |
| 13. Do. | 99 | 425 |
| 14. Do. | 99 | 343 |

The yield of *kapas* in all the dusted plots was markedly reduced in spite of the early intense flowering and the better retention of the earliest bolls in these plots. This effect was mainly due to the serious aphid infestation. No foliage injury was noticed on the plants dusted with calcium arsenate. It is significant to note that of all the dusted plots, the one in which the aphid infestation was partially controlled by using fish-oil rosin soap, yielded the largest amount of *kapas*.

It appeared that it would be necessary to give up the pursuit of this insecticide because it was found that its use on the cotton plants was always followed by severe aphid infestation which obliterated its beneficial effect. In the meanwhile, however, W. J. Folsom [1927] pointed out that the heavy infestation of aphid in small experimental plots dusted repeatedly with calcium arsenate was the result of the positive photo-tropic response of the winged aphid to the white dust, and of the immunity which they got from some of their enemies in the presence of this insecticide. He summarised his experience in the following words :—

“ Excessive applications of calcium arsenate are often followed by heavy infestations of the cotton aphid. The initial infestations are due to the positive photo-tropic reaction of winged females to white substances, such as, calcium arsenate, calcium carbonate, starch or flour. A heavy infestation is built up, not by the destruction of predators by calcium arsenate, but by the killing of hymenopterous parasites when they emerge in the presence of the arsenical. They are killed also, though more slowly, by calcium hydroxide, calcium carbonate, corn and starch.”

It was, therefore, decided that the white powder of the insecticide should be blackened with finely-powdered charcoal in order to find out if the aphid infestation could be prevented.

It was also decided to lay out a few trials for testing another insecticide, viz., sodium silico-fluoride, along with calcium arsenate during 1928-29; because Osborne [1926] had found that sodium fluosilicate was more toxic to the cotton boll-weevil than calcium arsenate.

The plants could not be dusted uniformly with the Vermorel Knapsack dusting machine used so far, and therefore, a new machine called “ The Peerless Dusting gun ” was purchased.

The calcium arsenate previously used contained more than .75 per cent soluble arsenic acid, which was undesirable according to Coad and Cassidy [1920], and a fresh stock of calcium arsenate was secured from Grasselli Chemical Co. of New York, U. S. A. (for analysis see Appendix I, Table XXXV). A stock of sodium silico-fluoride was also secured from the same firm.

It was further realised that it would not be possible to protect the cotton plants from the Spotted Boll-worms during all the period of three or four months of the activities of the pest. It was, therefore, necessary to ascertain the stage of plant

growth, at which the use of the insecticides would be most beneficial upon the yield of *kapas*. The period of plant growth was divided into three parts as under :—

- (1) Early part of the season, when most of the *Earias* population is concentrated on the shoots of cotton.
- (2) Period when the flower-bud development is most vigorous.
- (3) Period when the flower-opening is at its height.

In order, therefore, to determine the most suitable period for using the insecticides and in order to find out if the aphid infestation could be prevented by blackening the insecticides, the following trials were laid out during 1928-29 :—

Plot No. 1.—Dusting with blackened calcium arsenate when the Spotted Boll-worms are feeding on cotton shoots. •

Plot No. 2.—Dusting with blackened calcium arsenate when vigorous flower-bud development is in progress.

Plot No. 3.—Dusting with blackened calcium arsenate when the boll-development has commenced.

Plot No. 4.—Dusting with calcium arsenate when vigorous flower-bud development is in progress.

Plot No. 5.—Dusting with sodium silico-fluoride when flower-bud development is in progress.

Plot No. 6.—Dusting with blackened sodium silico-fluoride when flower-bud development is in progress.

Blackened calcium arsenate was prepared by mixing calcium arsenate and finely-powdered charcoal in equal proportions. Sodium silico-fluoride, is a heavier salt and therefore, one part of finely-powdered charcoal was mixed with two parts of this insecticide.

Each of the plots consisted of 100 plants, and in each case an adjoining plot with about the same number of plants was kept as a control.

The actual dates of dusting are given in Table XXXVI of Appendix I.

Foliage injury.—Showers of rain after dusting with sodium silico-fluoride were harmful to the foliage. The leaves developed brownish red spots but did not shed. They were seen on these plants until the bolls began to open. Some of the flower-buds, however, dropped away from these plants. There was no such injury noticed in the plots where calcium arsenate was used.

Insecticides and Spotted Boll-worms.—Examination of plants for *Earias* larvae from the dusted and the control plots showed that sodium silico-fluoride was distinctly more effective in reducing the *Earias* population in the plots where that insecticide was used.

TABLE LVII

Population of Earias larvae on dusted and control plants

1928-29

| Date of examination | Plot No. | No. of plants examined | No. of flower-buds | No. of bolls | Live <i>Earias</i> larvae | Dead <i>Earias</i> larvae |
|------------------------|------------------------|------------------------|--------------------|--------------|---------------------------|---------------------------|
| 13th November 1928 . . | 4 | 10 | 1,157 | 12 | 1 | 7 |
| 14th November 1928 . . | 2 | 10 | 1,193 | 17 | 17 | 14 |
| 15th November 1928 . . | 5 | 10 | 1,247 | 30 | 1 | 7 |
| 16th November 1928 . . | 6 | 10 | 1,250 | 30 | 6 | 6 |
| 16th November 1928 . . | Control (Non-treated). | 10 | 1,098 | 29 | 33 | .. |
| 1st December 1928 . . | 2 | 10 | 1,442 | 191 | 45 | 30 |
| 3rd December 1928 . . | 4 | 10 | 1,905 | 340 | 37 | 49 |
| 5th December 1928 . . | 5 | 10 | 596 | 227 | .. | 15 |
| 6th December 1928 . . | 6 | 10 | 747 | 246 | 2 | 9 |
| 6th December 1928 . . | Control | 10 | 1,333 | 108 | 38 | .. |

For the above examinations ten plants were up-rooted from each one of the above plots, once during the middle of November 1928, eight days after the commencement of the dusting in the respective plots and a second round of samples of ten plants from each plot was taken in the first week of December 1928, i.e., nine days after the 3rd application of the insecticide in each plot.

APHIS INFESTATION

In order to estimate the comparative infestation of this pest under different treatments, twenty cotton leaves were collected at random once in a week from plot Nos. 2, 4, 5 and 6 and also from a control plot. The aphid on these leaves were counted as accurately as possible.

TABLE LVIII

Population of aphids on 20 leaves from cotton plots dusted with white and blackened powders of insecticides

1928-29

| Plot No. | Treatment | No. of aphids on 20 leaves | | | | | | | |
|----------|--|----------------------------|-----------|-----------|-----------|----------|-----------|-----------|-----------|
| | | 8th Dec. | 15th Dec. | 22nd Dec. | 29th Dec. | 5th Jan. | 12th Jan. | 19th Jan. | 26th Jan. |
| 2 | Blackened calcium arsenate at bud-development. | traces | 7 | 147 | 110 | 274 | 252 | 575 | 772 |
| 4 | White calcium arsenate at bud-development. | " | 155 | 188 | 499 | 948 | 820 | 1,006 | 1,767 |
| 5 | White sodium silico-fluoride at bud-development. | " | 280 | 361 | 561 | 617 | 882 | 1,182 | 1,673 |
| 6 | Blackened sodium silico-fluoride at bud-development. | — | — | 13 | 28 | 106 | 104 | 78 | 270 |
| | Control (Not dusted) | — | 1 | 66 | 17 | 56 | 37 | 7 | 48 |

All the plots were sprayed once with fish-oil rosin soap on 9th January.

A small amount of aphid infestation was present on the general cotton crop from the middle of December till the end of January. This probably served as a nucleus for the higher infestation which was built up in the plots dusted with blackened calcium arsenate and sodium silico-fluoride. Out of these two plots the plants dusted with blackened sodium silico-fluoride showed a much less infestation, probably because this mixture was slightly darker in colour than the powder of blackened calcium arsenate. The plants dusted with white calcium arsenate and sodium silico-fluoride suffered from a very serious aphid infestation, because they must have attracted winged aphids from the general cotton area in addition to the traces of original infestation which existed on them. These trials, therefore, showed that blackening the insecticide would not altogether prevent the aphid infestation, but that it would not be as serious as on the plants dusted with white calcium arsenate or sodium silico-fluoride.

BOLL-DEVELOPMENT AND YIELD OF *kapas*

Twenty plants in each of the treated plots and their controls were reserved for studying the effect of the treatments on the flower opening and the development of bolls. The weekly flower opening and the relative number of bolls which matured is given in Appendix I, Table XXXVII. Besides this, the number of mature bolls

from all the plants from the plots were counted, and the *kapas* from each of the plants was collected and weighed separately. The average yield of bolls and *kapas* from these plants is given below.

TABLE LIX

Number of bolls and yield of kapas from the dusted plots and their controls

1928-29

| Plot No. | Treatment | Number of plants observed | Number of bolls per plant | Weight of kapas per plant in grms. | Percentage increase or decrease of kapas per plant over the control |
|----------|---|---------------------------|---------------------------|------------------------------------|---|
| 1 | Blackened calcium arsenate at shoot attack | 100 | 28.3 | 47.7 | —21.1 |
| 1-A | Control for plot No. 1 | 100 | 24.7 | 60.5 | |
| 2 | Blackened calcium arsenate at flower-bud development. | 100 | 28.2 | 55.2 | 8.6 |
| 2-A | Control for plot No. 2 | 99 | 27.8 | 50.8 | |
| 3 | Blackened calcium arsenate at boll-development. | 100 | 35.8 | 66.3 | 14.9 |
| 3-A | Control for plot No. 3 | 100 | 30.6 | 57.7 | |
| 4 | White calcium arsenate at flower-bud development. | 100 | 40.3 | 64.2 | 1.5 |
| 4-A | Control for plot No. 4 | 100 | 35.5 | 63.2 | |
| 5 | White sodium silico-fluoride at flower-bud development. | 100 | 25.4 | 43.5 | —43.2 |
| 5-A | Control for plot No. 5 | 99 | 41.1 | 76.6 | |
| 6 | Blackened sodium silico-fluoride at flower-bud development. | 100 | 32.5 | 60.4 | 12.3 |
| 6-A | Control for plot No. 6 | 100 | 29.0 | 53.8 | |

All the plots, where the plants were dusted with blackened insecticides yielded more *kapas* than their controls except plot No. 1 and plot No. 5.

Plot No. 1 received three dustings with blackened calcium arsenate on 8th, 18th and 29th October 1928. These plants were not dusted afterwards. As a result of these dustings, these plants began to flower vigorously early in November (Appendix I, Table XXXVIII); but the application of the insecticide to these plants had already stopped, and therefore, the big flower-buds and the young bolls

shed in large numbers due to the damage of the Spotted Boll-worms (Plate XV). It was, therefore, found that the use of the insecticides during the early part of the season would not be practicable because the dustings could not be continued till about the end of December when the Spotted Boll-worms normally declined, and if they were discontinued earlier, all the advantage gained by the early dustings would be lost, because the Spotted Boll-worms would concentrate in these advanced plots and would damage the developed forms. It might, therefore, be stated, in general, that the use of insecticides like calcium arsenate and sodium silico-fluoride during the early part of the plant growth would be distinctly harmful under Surat conditions.

The plot of blackened calcium arsenate at boll-development yielded more *kapas* than the plot which received the same insecticide at the time of flower-bud development.

White sodium silico-fluoride at the time of flower-bud development turned out to be distinctly harmful, but gave 12.3 per cent more *kapas* when it was blackened before use.



The plants, dusted with white calcium arsenate at flower-bud development, yielded nearly five bolls more than the control plants, but they could hardly yield two per cent additional *kapas*. This was apparently due to the fact that the bolls on these plants did not develop normally due to excessive aphid infestation during the period of their development.

SLAKED LIME AND APHIS INFESTATION

Powder of slaked lime is quite as white as the powder of calcium arsenate. A test was made to see if the use of slaked lime alone would lead to serious aphid infestation. During the course of the above trials, two groups of twenty plants each were selected, and one was dusted with calcium arsenate and the other with slaked lime. After an interval of about ten days both the groups showed the commencement of aphid attack, but the plants dusted with lime had comparatively very few aphids on them. Within another ten days, the plants with calcium arsenate were seriously infested with aphids, whereas, their increase on the plants dusted with lime was very slow. It was, therefore, clear that calcium arsenate afforded very much better conditions for the multiplication of aphids.

TRIALS DURING 1929-30

In the light of the experience of the previous season, it was decided that the use of the insecticides should be concentrated during the boll-development period only.



1928 - 29

- weekly flower opening
- x — x — bolls shed
- - - shed bolls showing boll-worm attack.

Showing weekly flower-opening, shedding of bolls and bolls shed due to boll-worms in a cotton plot dusted with calcium arsenate during the period of shoot attack; and also in the control plot.

It appeared that during the previous trials excessive quantities of insecticides were applied, and hence it was decided to regulate the doses.

Nicotine sulphate ("Black leaf 40") was also to be used for attempting the control of aphids.

The following trials were, therefore, laid out :—

Plot No. 1.—A cotton plot of one acre to be dusted with blackened calcium arsenate at the rate of 20 lbs. of mixture per acre. Nicotine sulphate to be used when aphids appear.

Plot No. 2.—One hundred plants to be dusted with blackened calcium arsenate at the rate of 20 lbs. of mixture per acre.

Plot No. 3.—One hundred plants to be dusted with white calcium arsenate at 20 lbs. to an acre and nicotine sulphate to be used when aphids appear.

Plot No. 4.—One hundred plants to be dusted with white calcium arsenate at 10 lbs. per acre.

Plot No. 5.—One hundred plants to be dusted with blackened sodium silico-fluoride at 40 lbs. of mixture per acre.

Plot No. 6.—One hundred plants to be dusted with blackened sodium silico-fluoride at 40 lbs. of mixture per acre and nicotine sulphate to be used for controlling the aphids.

Plot No. 7.—One hundred plants to be dusted with white sodium silico-fluoride at 20 lbs. per acre. Details of the dates of dustings, the quantities of insecticide actually used and their cost calculated per acre are given in Appendix I, Table XXXIX.

APHIS INFESTATION DURING 1929-30

Traces of aphids were noticed during this season also from the month of December onwards on the general crop of cotton. On the dusted plots they began to increase rapidly within four weeks after the commencement of the dustings.

Small plots of about 100 plants, when they were dusted with calcium arsenate or sodium silico-fluoride, developed a serious infestation of aphids. An area of one acre of cotton (Plot No. 1) was taken for dusting with blackened calcium arsenate in order to observe the effect upon a very much larger area when under this treatment. Aphid infestation turned out to be the worst in this plot, and could not be checked even with two applications of nicotine sulphate (Table LX).

In plots No. 3, 4 and 7 the white powder of insecticides was used without any dilution. In all these plots, the infestation had considerably advanced by the end of December. Nicotine sulphate was then used in plot No. 3, but the reduction in the population of aphids was only temporary.

In the remaining plots No. 2, 5 and 6 where the insecticides were blackened, the infestation was comparatively much lower by the middle of January, and it remained fairly low throughout the season in plot No. 6, where besides blackening the dust, nicotine sulphate also was used twice.

TABLE LX

Weekly population of aphids on the dusted plants in 1929-30

(20 leaves examined each time)

| Plot No. | Treatment | No. of aphids on 20 leaves every week | | | | | | | | REMARKS |
|----------|--|---------------------------------------|-----------|-----------|----------|-----------|-----------|----------|-----------|---|
| | | 15th Jan. | 22nd Jan. | 29th Jan. | 5th Feb. | 13th Feb. | 21st Feb. | 2nd Mar. | 12th Mar. | |
| 1 | One acre plot blackened calcium arsenate at 20 lbs per acre | 637 | 776 | 934 | 732 | 1,132 | 1,378 | 952 | 130 | Nicotine sulphate used on 2nd and 21st Jan. |
| 2 | Blackened calcium arsenate at 20 lbs per acre (10th plants) | 53 | 112 | 84 | 100 | 242 | 1,245 | 356 | 133 | |
| 3 | White calcium arsenate at 20 lbs per acre (100 plants). | 544 | 142 | 48 | 211 | 349 | 587 | 174 | 65 | Nicotine sulphate used on 24th Dec., 3rd Jan. and 21st Jan. |
| 4 | White calcium arsenate at 10 lbs. per acre (100 plants). | 392 | 205 | 288 | 471 | 827 | 1,863 | 453 | 192 | |
| 5 | Blackened sodium silico-fluoride at 40 lbs. per acre (100 plants). | 40 | 56 | 107 | 87 | 126 | 430 | 243 | 78 | |
| 6 | Blackened sodium silico-fluoride at 40 lbs. per acre (100 plants). | 24 | 17 | 41 | 85 | 54 | 73 | 15 | traces | Nicotine sulphate used on 4th and 21st Jan. |
| 7 | White sodium silico-fluoride at 20 lbs. per acre (100 plants). | 132 | 139 | 561 | 565 | 653 | 956 | 339 | 108 | |
| | Control (Not dusted) | traces all along. | | | | | | | | |

It is thus seen that aphid infestation cannot be altogether prevented by blackening the insecticides, but it can be prevented from being as serious as when the

insecticides are used without any dilution. Blackened sodium silico-fluoride, being darker than blackened calcium arsenate, was found to be comparatively more effective in preventing serious infestation.

This drawback of these insecticides is likely to be removed to a considerable extent if the powders can be made jet black by adding small quantities of an inert colour. Nicotine sulphate mixed with these insecticides or with lime is not quite effective in removing the infestation entirely, and the use of such a costly insecticide on a large scale is prohibitive.

YIELD FROM DUSTED PLANTS AND THEIR CONTROLS

Acceleration of flowering and better retention of the earlier flowers was noticed during this season also (Appendix I, Table XL). But, as already stated, the aphid infestation appeared this year also and prevented any accurate measure of the effectiveness of the insecticides. Plots No. 5 and 7 alone showed an increase of 7 and 20 per cent respectively in the yield of *kapas*, whereas the increase was insignificant in plot No. 4, and there was a decrease in the yield of the remaining four plots.

TABLE LXI

Yield of kapas from the plot of one acre dusted with blackened calcium arsenate and also from the three adjoining control plots

1929-30

| Treatment | Area under trial | Yield of kapas per acre |
|---|------------------|-------------------------|
| | Acres | Pounds |
| Dusted with blackened calcium arsenate at 20 lbs. per acre and used nicotine sulphate twice (Plot No. 1). | 1 | 370 23 |
| Control No. 1 | 1 | 382-10 |
| Control No. 2 | 1 | 386-0 |
| Control No. 3 | 1 | 362-65 |

TABLE LXII

Yield of mature bolls and kapas from the dusted plots and their controls, in 1929-30

(100 plants in each case)

| Plot No. | Treatments | Total No. of bolls | Total yield of salcable kapas in grms. | Percentage increase or decrease over control |
|----------|---|--------------------|--|--|
| 2 | Blackened calcium arsenate at 20 lbs. per acre | 2,871 | 5,645 | |
| 2A | Control | 3,034 | 5,874 | -3.9 |
| 3 | White calcium arsenate at 20 lbs. per acre and nicotine sulphate against aphis. | 2,650 | 5,154 | |
| 3A | Control | 2,818 | 5,634 | -8.5 |
| 4 | White calcium arsenate at 10 lbs. per acre . . | 2,647 | 5,092 | |
| 4A | Control | 2,494 | 4,976 | 2.3 |
| 5 | Blackened sodium silico-fluoride at 40 lbs. per acre. | 2,781 | 5,400 | |
| 5A | Control | 2,579 | 5,062 | 6.6 |
| 6 | Blackened sodium silico-fluoride at 40 lbs. per acre and nicotine sulphate against aphis. | 2,690 | 5,360 | |
| 6A | Control | 2,710 | 5,551 | -3.4 |
| 7 | White sodium silico-fluoride at 20 lbs. per acre . | 3,231 | 6,232 | |
| 7A | Control | 2,600 | 5,185 | 20 |

The first four plots were devoted to studying the most effective way of using calcium arsenate. Out of these plots, No. 4 only showed a slight increase in the yield of *kapas*.

Sodium silico-fluoride was tried in the remaining three plots, and two of them (Plots No. 5 and 7) yielded 6.6 and 20 per cent more *kapas* respectively than their controls. The increase in yield in both the cases was not large enough to cover the cost of the insecticides, as can be seen from the following considerations :—

- (1) Plot No. 5 yielded 338 grms. more *kapas* than its control. At this rate, the increased yield of *kapas* per acre would be 28 pounds and would fetch about Rs. 4-3-0, while the cost of the insecticides required would be about Rs. 27-12-0.

- (2) Plot No. 7 yielded 1048 grms. of *kapas* more than its control. At this rate the extra yield of *kapas* would be about 86 pounds per acre, and would fetch about Rs. 12-14-0. The cost of the insecticides required for the treatment of one acre would be about Rs. 23-8-0.

Plots No. 5 and 6 both received similar treatments, and plot No. 6 received the further advantage of nicotine sulphate in the last two dustings. It is, however, strange to find that plot No. 5 has yielded more *kapas* than its control, whereas the yield of *kapas* from plot No. 6 has been less than that of its own control.

The experiments indicate that it is uneconomic to use these insecticides for controlling the Spotted Boll-worms. As the trials have been described, more or less in chronological order, the more important conclusions are summarised below.

Conclusions

(1) Lead arsenate and Paris Green, applied to the cotton plants in the form of dust or spray, are not useful in preventing damage by Spotted Boll-worms.

(2) Calcium arsenate dusted on the cotton plants, without any dilution, is effective in killing a large number of the *Earias* larvae. The tiny larvae, as soon as they emerge from the eggs, have the habit of wandering about on the plants, for some time, before they start feeding. These larvae come in contact of the dust and are killed in large numbers.

(3) The protection afforded by this insecticide to the cotton plants from the *Earias* larvae, results in the earlier flowering on the dusted plants and a better retention of bolls from these flowers.

(4) Within a period of about four weeks from the commencement of dusting of calcium arsenate, a serious infestation of aphids develops on the cotton plants. As a result of this infestation, the yield of *kapas* from these plants is generally reduced.

(5) It is found that the white colour of this insecticide attracts the winged aphids, which multiply rapidly on the plants. Blackening the dust of calcium arsenate, by adding an equal quantity of finely powdered charcoal before using, reduces the infestation but does not prevent it altogether; because the small local infestation, which exists on most of the cotton plants, gets extremely favourable conditions for multiplication, as they are protected from some of their important enemies by this insecticide.

(6) Sodium silico-fluoride also, in the form of white dust, is more effective in killing *Earias* larvae on the cotton plants than calcium arsenate.

(7) Earlier flower-opening and better retention of bolls from the early flowers is noticed in plots dusted with sodium silico-fluoride.

(8) One part of finely powdered charcoal and two parts of sodium silico-fluoride form a mixture which is darker than blackened calcium arsenate, and probably due to this reason, the aphid infestation remains at a slightly lower level on the plants dusted with this mixture than on those which receive blackened calcium arsenate.

(9) The leaves of cotton are scorched, and develop reddish brown patches, if showers of rain are received after the plants are dusted with sodium silico-fluoride. These leaves, however, do not shed but some of the flower-buds drop away due to this injury.

(10) The use of calcium arsenate or sodium silico-fluoride, only during the early part of the cotton-growing season, is harmful because the flower-buds and bolls saved from the Spotted Boll-worms during the above period are seriously damaged soon after the dustings are discontinued. It is, therefore, advisable to start the use of these insecticides late in the season and to continue it till the *Earias* population normally declines by the end of December or early in January.

(11) Nicotine sulphate used after mixing with calcium arsenate, sodium silico-fluoride or slaked lime (5 parts nicotine sulphate to 100 parts of one of the three powders) helped to dislodge most of the grown-up aphids from the plants, and thus to reduce the infestation to a great extent. This reduction was, however, only temporary.

(12) In several cases the yields of the plants, which were dusted either with white or blackened calcium arsenate or sodium silico-fluoride, were greatly reduced, in spite of the advantages mentioned above. This was apparently due to the serious aphid infestations which developed on the dusted plants.

(13) Higher yields of *kapas* were obtained from the plots, where the following treatments were attempted, than their respective controls. The extra yields, however, did not cover the cost of the insecticides, and hence the treatments were found to be uneconomic.

- (1) 8.6 per cent higher yield of *kapas* during 1928-29 by using blackened calcium arsenate at flower-bud development (4 dustings).
- (2) 14.9 per cent higher yield during 1928-29 by using blackened calcium arsenate during boll-development (4 dustings).
- (3) 12.3 per cent higher yield during 1928-29 by using blackened sodium silico-fluoride at flower-bud development (4 dustings).
- (4) 6.6 per cent higher yield during 1929-30 by using blackened sodium silico-fluoride at 40 lbs. of mixture per acre for each dusting (4 dustings).
- (5) 20 per cent higher yield during 1929-30 by using white sodium silico-fluoride at 20 lbs. per acre for each dusting (4 dustings).

(14) There is a great possibility of these insecticides (calcium arsenate and sodium silico-fluoride) being very useful in controlling the Spotted Boll-worms if by some means the aphid infestation, which develops after their use, can be prevented altogether; or if a very cheap and effective method can be found for destroying the aphids.

(15) Of the two insecticides, sodium silico-fluoride is the more efficient.

8. CLEAN-UP MEASURES

The experience of Gough in Egypt [Gough, 1919] showed that there was a great possibility of controlling the Spotted Boll-worms on cotton, if suitable conditions are available for keeping all the cotton tract free from the food of these insects during the period between the two cotton-growing seasons.

At the Third Entomological Conference at Pusa [1919] Dr. Gough stated his experience as under :—

“.....We had legislation against *Earias* and the result is that what was a bad pest in 1912 is only a quite minor pest now. In 1910-12 at the electric light on our verandah we used to attract twenty or thirty *Earias insulana* moths every evening, but now-a-days this species is comparatively a rarity, and the percentage of cotton bolls attacked by *Earias* is well below ten per cent. The legislation laid down demands that all cotton sticks be pulled out and burnt; and this applies also to *Hibiscus esculentus* and *Hibiscus cannabinus*. Thus for five months there is no food available for *Earias* as we do not leave any cotton or *Hibiscus* or any other food plant. Legislation on such lines forms the best means of control.”

The attempts at direct control having failed to provide a practical remedy, it became necessary to consider if a preventive measure like the clean-up campaign could be profitably used for controlling this pest on cotton in Gujarat.

The last picking of cotton is generally over in this tract before the end of April, and sowing of the cotton crop of the next season commences some time from the middle of June to the middle of July according to the condition of the monsoon. Usually it is not till August that the plants grow to a size when the Spotted Boll-worms can thrive on them. It is, therefore, necessary to understand what happens to these insects during this period of nearly three months, and what may be the sources of fresh infection to the next crop. The possibilities are :—

- (1) Aestivation between cotton seasons in one stage or another of their life.
- (2) Active breeding throughout this period, in places where food may be available.

Nothing is so far known which would indicate the possibilities of aestivation of the Spotted Boll-worms. They have been collected in active condition throughout the year and have been reared in the laboratory during all the months. It was, therefore, decided to study the role of different plants in providing food for the pest in the field during the period of about three months between seasons.

- (1) Wild food plants.
- (2) Cultivation of *bhendi*.
- (3) Fresh growth from cotton plants and their stumps.

Wild food-plants

The following wild food-plants of the Spotted Boll-worms are met with in the locality :—

- (1) *Hibiscus rugosus*.
- (2) *Hibiscus manihot*.
- (3) *Hibiscus ficulneus*.
- (4) *Hibiscus panduræformis*.
- (5) *Abutilon indicum*.
- (6) *Abutilon graveolens*.

All these plants spring up after the commencement of the monsoon, and it is not till about the end of August that *Earias* population can be found on them. The pest appears almost simultaneously on these weeds and the fresh crop of cotton (Appendix I, Tables XLI to XLIV). *Earias* larvae can easily be collected from these weeds during October and November, and the *Abutilons*, in particular, are severely attacked during these months (Appendix I, Table XLIV).

By about the month of January, however, most of these plants dry up. Only a few stray plants of *Abutilon* are noticed with green leaves and pods, after January, in hedges or in neglected parts of gardens (Appendix I, Table XLV). A few plants of *Hibiscus panduræformis* are also found in the fields during summer, but the population of boll-worms on these plants is very meagre.

Barring the stray plants of *Abutilon* and *Hibiscus panduræformis*, the weeds do not afford any help for the Spotted Boll-worms to tide over the off-season.

CULTIVATION OF *bhendi* (*Hibiscus esculentus*)

It is usually with the first showers of the monsoon that the *bhendi* crop is sown on a large scale, and quantities of *bhendi* pods are available in the market by the middle of August. Though this is mainly a monsoon crop, a few scattered plots of this vegetable are met with even during the summer in the irrigated areas around wells. These plants are profusely attacked by the Spotted Boll-worms (Appendix I, Table XLVI).

This happens to be a favourite host for *Earias* caterpillars and the cultivation of *bhendi* certainly helps the carry-over of a very large number of these insects from one cotton-growing season to another.

Besides these stray plots of *bhendi* in summer, there is another possibility of the infection of the pest being carried by these pods. During May, June and July, the local supply of this vegetable becomes almost negligible, and pods are, therefore, imported in the important market centres, like Surat, from distant places like Ahmedabad, Nasik and Poona. These pods in the market also show the presence of *Earias* larvae in them. A coastal village named Hajira near Surat also specially grows *bhendi* during the summer and supplies these pods to Surat market in June

TABLE LXIII

Examination of bhendi pods from Surat market

| Date of examination | Quantity of pods examined | No. of pods examined | No. of <i>Earias</i> larvae | Sources of pods |
|-------------------------|---------------------------|----------------------|-----------------------------|-------------------------|
| | Lbs. | | | |
| 6th May 1929 | 3 | 200 | 3 | Ahmedabad. |
| 18th May 1929 | 5 | 240 | 3 | Do. |
| 25th May 1929 | 5.5 | 362 | 13 | Hajira, District Surat. |
| 1st June 1929 | 5 | 329 | 8 | Do. |
| 8th June 1929 | 4 | 255 | 12 | Do. |
| 15th June 1929 | 2 | 114 | 9 | Do. |
| 22nd June 1929 | 5 | 294 | 15 | Do. |
| 29th June 1929 | 5 | 260 | 8 | Do. |
| 5th July 1929 | 5 | 222 | 4 | Poona or Nasik side. |
| 12th July 1929 | 5 | 191 | 17 | Do. |
| 19th July 1929 | 5 | 189 | 5 | Do. |
| 27th July 1929 | 5 | 192 | 2 | Do. |
| 2nd August 1929 | 5 | 176 | nil. | Do. |
| 3rd May 1930 | 4 | 275 | 8 | Ahmedabad side. |
| 8th May 1930 | 5 | 274 | 13 | Do. |
| 15th May 1930 | 4 | 173 | 11 | Do. |
| 24th May 1930 | 3 | 232 | 10 | Do. |
| 31st May 1930 | 5 | 328 | 18 | Do. |
| 7th June 1930 | 5 | 290 | 11 | Hajira, District Surat. |
| 14th June 1930 | 5 | 228 | 3 | Do. |
| 21st June 1930 | 5 | 199 | 2 | Do. |
| 28th June 1930 | 5 | 170 | 6 | Bombay side. |
| 5th July 1930 | 5 | 243 | 3 | Do. |
| 12th July 1930 | 5 | 228 | 3 | Do. |

Bhendi is thus seen to be an important source of food of the Spotted Boll-worms during the period between the two cotton-growing seasons, and it is, therefore, a source of infection to the cotton crop of the next season, not only in places where it is cultivated but also in distant places where its pods may be exported for table purposes.

FRESH GROWTH FROM THE COTTON PLANTS

When the last picking of cotton is over, the green growth on the cotton plants, consisting of tender shoots, flower-buds and young bolls, is practically absent, except on a few border plants. A few days later, however, most of these plants begin to sprout, and fresh vegetative shoots, flower-buds and young bolls begin to arise. The fresh growth from the standing cotton plants provides food and shelter to the Spotted Boll-worms until the plants are removed by the cultivators (Appendix I, Table XLVII), who do not show any particular anxiety to clean-up their cotton fields very early and the operation of the removal of the cotton plants proceeds slowly.

TABLE LXIV

The progress of the removal of cotton plants in a village called Piplod near Surat

| Time of removal | Area cleared up in acres | REMARKS |
|---------------------------|--------------------------|---|
| 26th April 1928 | 30 | Total area 116 acres. All picking of <i>kapas</i> over on the 10th April 1928. |
| 9th May 1928 | 46 | |
| 23rd May 1928 | 12 | |
| 7th June 1928 | 22 | |
| 23rd March 1929 | 5 | Total area 93.25 acres. All pickings of <i>kapas</i> over by the last week of March 1929. |
| 6th April 1929 | 21 | |
| 20th April 1929 | 59 | |
| 3rd May 1929 | 7 | |
| 17th May 1929 | 1 | |
| 1st June 1929 | $\frac{1}{2}$ | |

Fields are often seen where cotton plants are allowed to stand until the monsoon and a few cases are also found where the plants continue to stand even during the next season, the plots being neglected by the cultivators due to some reason or the other.

In one case a ratoon crop of cotton was observed in a plot at Bharthana near Surat during 1928-29. The plants were cut off, leaving a stump of about 8 to 12 inches from the ground. These plants put up luxuriant growth with the commencement of the monsoon, and flowering was noticed on these plants in the second fortnight of September, when the flower-bud development on the new crop had just begun. Ten plants from this plot were kept under observation, and another group of ten plants from the new crop was reserved for similar observations for comparison (Appendix 1, Table XLVIII). The first flush of vigorous flowering on the ratoon plants appeared very early and continued till the middle of October, while the flowering on the new crop commenced in the middle of November. The result of the early flowering on the ratoon crop was that the Spotted Boll-worms concentrated on these plants, and destroyed most of the young bolls and the developed flower-buds and the vigorous flowering abruptly came to a close at the end of the second week of October and could not commence again for about four weeks. The second flush of flowering which appeared on the ratoon plants along with the normal flowering of the new crop was very poor.

TABLE LXV

Flowering and boll-shedding on the ratoon plants and the new crop of cotton

(10 plants in each case)

| | Flowers opened | Bolls matured | Bolls shed | Bolls shed due to <i>Earias</i> |
|-----------------------|-------------------|------------------|---------------|------------------------------------|
| Ratoon crop | 234 | 42 | 192 | 136 |
| New crop | 548 | 228 | 320 | 106 |

These observations indicate that the ratoon crop of cotton cannot yield a good crop of *kapas* as it is severely damaged by the Spotted Boll-worms, and also that it is harmful to the new crop of cotton as it affords a good breeding ground for this pest in the earliest part of the season.

It is fortunate that the fresh growth from the standing cotton plants in summer from most of the open fields (not fenced) is generally destroyed by the cattle; but the cotton plant shows a remarkable tenacity of life and the cotton sticks grazed by the cattle quickly develop fresh shoots.

Next to *bhendi*, therefore, these standing cotton plants are responsible for supplying food to these larvae for passing over from one cotton-growing season to another.

FRESH GROWTH FROM STUMPS

The cotton plants are removed by the cultivators by hacking them with a *kudali* (light pick-axe) near the ground level (Plate XVI, Fig. I). A small stump of each one of these plants remains in the soil. Some of these stumps put up fresh growth within a few days after cutting, and plots from which the plants are removed the earliest usually contain a large number of sprouts, even before many of the other cotton fields are cleared off. Thus there is an overlapping of the sprouting from the standing cotton sticks and the stumps.

TABLE LXVI

Sprouts of cotton stumps from a plot on Surat Farm
(Plants removed by hacking with *kudali* on 23rd April 1928)

| Date of observation | Total No. of hills in the plot | No. of hills showing sprouts | Percentage of sprouting |
|---|--------------------------------|------------------------------|-------------------------|
| 21st May 1928 | 2,105 | 812 | 39 |
| 8th June 1928 | 2,105 | 1,286 | 62 |
| Plot harrowed on the 14th of June preparing the land for <i>jowar</i> | | | |
| 28th June 1928 | 2,105 | 186 | 9 |
| 11th July 1928 | 2,105 | 184 | 9 |
| <i>Jowar</i> sown in the third week of July | | | |
| 31st July 1928 | 2,105 | 1,065 | 51 |

TABLE LXVII

Sprouts from cotton stumps from a cultivator's plot
(Plants removed by hacking with a *kudali* on 3rd April 1929)

| Date of observation | Total No. of plants removed | No. of stumps showing sprouts | Percentage of sprouting |
|------------------------------------|-----------------------------|-------------------------------|-------------------------|
| 7th May 1929 | 6,091 | 2,239 | 37 |
| 22nd May 1929 | 6,091 | 3,243 | 53 |
| 30th May 1929 | 6,091 | 3,516 | 58 |
| Not harrowed in the middle of June | | | |
| 26th June 1929 | 6,091 | 947 | 16 |



FIG. 1. Removing the cotton plants with a light pick-axe.




FIG. 2. A shoot sprouting from a stump of cotton plant.

PLATE XVII.




FIG. 1. A shoot sprouting from a side root of a stump of cotton plant.




FIG. 2. A cotton plant-puller.

In another case, ninety-three acres of cotton were kept under observation during 1929 in a village called Piplod. The last picking of cotton was over at the end of March 1929.

TABLE LXVIII

Rough estimate of sprouting from stumps in 93 acres of cotton at Piplod 1929

| Date of observation | ACRES SHOWING PERCENTAGE SPROUTING FROM STUMPS | | | |
|---------------------------|--|----------------------------|----------------------------|-----------------------|
| | Less than 25 per cent | Between 25 and 50 per cent | Between 50 and 75 per cent | More than 75 per cent |
| 27th April 1929 | 10 | .. | .. | .. |
| 3rd May 1929 | .. | .. | 8 | .. |
| 10th May 1929 | 8 | .. | 3 | .. |
| 17th May 1929 | 28 | .. | 5 | .. |
| 25th May 1929 | 61 | 5 | 5 | .. |
| 1st June 1929 | 54 | 7 | 2 | 5 |
| 12th June 1929 | 51 | 13 | 9 | 5 |
| 22nd June 1929 | 35 | 18 | .. | 1 |
| 27th June 1929 | 52 | 21 | 13 | .. |
| 20th July 1929 | 67 | 8 | .. | 7 |

It is interesting to note that most of the sprouts arise from the adventitious buds, because the plants are cut off below the cotyledonary nodes (Plate XVI, Fig. 2). More interesting still is the fact that we often come across shoots which have sprung up from the side roots of these stumps (Plate XVII, Fig. 1).

Earias larvae can be collected from these shoots as soon as they are about four to six inches in height (Appendix I, Tables XLIX and L). It is, therefore, seen that the fresh growth from the standing cotton plants and their stumps, the cultivation of *bhendi*, and the presence of stray plants of a couple of malvaceous weeds constitute the supply of food, which enable the Spotted Boll-worms to pass over from one cotton-growing season to another.

The cultivation of *bhendi* during this period is most harmful, because it harbours a very large population of these worms and the presence of even a few plants is enough to re-infect large areas of cotton. The responsibility of standing sticks, for providing food and shelter to these worms during the off-season, comes next.

The weeds and the sprouts from cotton stumps are comparatively less dangerous because the *Earias* population on them, during this critical period, is generally very meagre (Plates XVIII and XIX).

Having traced the means of carry-over of this pest from one season to another in Gujarat, where the period between the two cotton-growing seasons is comparatively short, observations were made in Khandesh where the off-season between the two crops is more than six months, and where the climatic and soil conditions are unsuited for the sprouting of the cotton plants or their stumps (Appendix IV).

These observations showed that the cotton areas in Khandesh are cleaned up before the end of March, and the stumps do not put forth new shoots, as they do in Gujarat, and hence there is no food available for the caterpillars in general cotton areas. There are, however, many irrigation wells in several villages, and the irrigated plots around these wells are found to be chiefly responsible for helping the carry-over of this pest in Khandesh, because old plants of cotton and plants of *Abutilon* are found growing in some of these plots, and *bhendi* is also grown in some during the period between the two cotton-growing seasons.


The knowledge of these facts leads to the conclusion that if the above sources of food, during the period between the two cotton-growing seasons, can be destroyed, the presence of the Spotted Boll-worms on the next crop of cotton can be eliminated. This would only be possible if a perfect clean-up campaign were organised over a very large area, and the cleaning-up is thoroughly carried out. It is essential to attend to the following points while arranging such a campaign :—


(1) DESTRUCTION OF WEEDS

It has already been mentioned that out of the several wild food plants of the Spotted Boll-worms, only a very few plants of *Abutilon* and *Hibiscus panduræformis* are met with in green condition during the summer. These must be destroyed. It must, therefore, be brought home to the cultivators that these plants growing in their plots and hedges are distinctly harmful to the cotton crop, and must, therefore, be removed as soon as they make their appearance.

(2) ELIMINATION OF *bhendi*

Bhendi in summer can only be grown where irrigation facilities are available. Even with these facilities it is very difficult to get a good crop in summer, because the growth of the plants is stunted due to the extreme heat; and also they are seriously infested by the Spotted Boll-worms. It is, therefore, advisable that the plants should not be grown by the cultivators during the months of April, May and June, in the interest of the large areas of their cotton crop.





(3) PREVENTION OF GROWTH FROM THE STANDING COTTON PLANTS AND THEIR STUMPS

Fresh growth from the standing cotton plants will not be available if the plants are removed immediately after the pickings of *kapas* are over. The absence of these plants will also prevent unnecessary drain of the plant food from the soil.

It is found that when the plants are removed by the usual method, fresh growth appears from the stumps which remain in the soil. A trial was made of removing the plants a little deeper, by hacking them after removing one or two clods near the stems with a light pick-axe.

TABLE LXIX

Sprouting from stumps in a plot where the plants were dug deeper on 27th April 1928

Surat Farm

| Date of observation | Total No. of hills in the plot | No. of hills showing sprouts | Percentage of the sprouts |
|--|--------------------------------|------------------------------|---------------------------|
| 23rd May 1928. | 2,771 | 44 | 1.6 |
| 8th June 1928 | 2,771 | 129 | 4.7 |
| Plot harrowed on 14th June for preparing the land for <i>jowar</i> . | | | |
| 28th June 1928 | 2,771 | 43 | 1.6 |
| 11th July 1928 | 2,771 | 48 | 1.7 |
| <i>Jowar</i> sown in the third week of July. | | | |
| 31st July 1928 | 2,771 | 53 | 1.9 |

TABLE LXX

Sprouting from stumps in a plot where the plants were dug deeper on 3rd April 1929


Cultivator's plot

| Date of observation | No. of plants removed | No. of stumps showing sprouts | Percentage sprouting |
|--------------------------------------|-----------------------|-------------------------------|----------------------|
| 7th May 1929 | 5,707 | 187 | 3 |
| 22nd May 1929 | 5,707 | 560 | 10 |
| 30th May 1929. | 5,707 | 837 | 15 |
| Plot harrowed in the middle of June. | | | |
| 26th June 1929 | 5,707 | 190 | 3 |

The sprouting from stumps was considerably reduced by this method, but it was not eliminated. It was, therefore, considered that the sprouting could be prevented completely, only if the plants were pulled out with their roots. In several cases it was impossible to pull out the cotton plant in this locality even if two men tugged at it. It was, therefore, decided to prepare a mechanical device for doing this work. A large number of small tools of iron and wood were devised but they were not found to be quite suitable for this work. In the meanwhile we came across an old model prepared some time in 1911 at Coimbatore, devised on the principle of a nail-puller such as is used for extracting nails from deal-wood boxes. It could pull out the plants, but it had a very heavy handle of iron, and it often went out of order, as it was defective in several places. A large number of trials were made to perfect this tool, and the construction had to be radically changed before a satisfactory implement for uprooting cotton plants was evolved.

With this plant puller (Plate XVII, Fig. 2 and Plate XX), the cotton plants can be pulled out very easily. Most of the tap root and the main side roots are pulled out, while the thinner roots are left in the soil. A cultivator can work with this implement for the whole day without any abnormal exertion, and can clear up at least as much area as he can do with his *kudali* by the old method. The implement at present costs Rs. 1-4-0.

Trials are now in progress for testing these measures by organising a clean-up campaign in an area of about 480 square miles, which has about 80,000 acres under cotton in the Broach District. The results of these trials will be available in due course.



A cotton-plant puller in action.

APPENDIX I

TABLE I

Length of life of Earias moths and the eggs laid by them, 1930-31

| Pair No. | Date of emergence of moths | No. of days the female moth lived | No. of days the male moth lived | Total No. of eggs laid by the female | REMARKS |
|----------|----------------------------|-----------------------------------|---------------------------------|--------------------------------------|---|
| 1 | 4th November 1930 . . . | 23 | 20 | 367 | 341 eggs per female moth, which emerged in November 1930. |
| 2 | 5th November 1930 . . . | 23 | 12 | 655 | |
| 3 | 7th November 1930 . . . | 23 | 23 | 185 | |
| 4 | 7th November 1930 . . . | 20 | 20 | 415 | |
| 5 | 8th November 1930 . . . | 21 | 21 | 617 | |
| 6 | 10th November 1930 . . . | 14 | 14 | 332 | |
| 7 | 10th November 1930 . . . | 18 | 18 | 335 | |
| 8 | 22nd November 1930 . . . | 16 | 13 | 414 | |
| 9 | 24th November 1930 . . . | 20 | 20 | 400 | |
| 10 | 27th November 1930 . . . | 24 | 10 | 424 | |
| 11 | 27th November 1930 . . . | 17 | 17 | 100 | |
| 12 | 28th November 1930 . . . | 10 | 4 | 160 | |
| 13 | 28th November 1930 . . . | 14 | 14 | 134 | |
| 14 | 29th November 1930 . . . | 15 | 5 | 231 | |
| 15 | 3rd December 1930 . . . | 28 | 27 | 357 | 298 eggs per female moth, which emerged in December 1930. |
| 16 | 7th December 1930 . . . | 27 | .. | 608 | |
| 17 | 7th December 1930 . . . | 24 | .. | 238 | |
| 18 | 8th December 1930 . . . | 13 | .. | 229 | |
| 19 | 8th December 1930 . . . | 24 | 22 | 101 | |
| 20 | 8th December 1930 . . . | 32 | 10 | 423 | |
| 21 | 8th December 1930 . . . | 34 | .. | 354 | |
| 22 | 12th December 1930 . . . | 8 | 7 | 110 | |
| 23 | 18th December 1930 . . . | 14 | 14 | 232 | |
| 24 | 20th December 1930 . . . | 27 | .. | 589 | |
| 25 | 21st December 1930 . . . | 29 | 4 | 192 | |
| 26 | 21st December 1930 . . . | 16 | 12 | 373 | |
| 27 | 31st December 1930 . . . | 11 | 5 | 171 | |
| 28 | 31st December 1930 . . . | 14 | 14 | 198 | |
| 29 | 1st January 1931 . . . | 8 | .. | 182 | 185 eggs per female moth, which emerged in January 1931. |
| 30 | 1st January 1931 . . . | 11 | 4 | 114 | |
| 31 | 3rd January 1931 . . . | 11 | .. | 279 | |
| 32 | 7th January 1931 . . . | 7 | .. | 156 | |
| 33 | 9th January 1931 . . . | 12 | 9 | 96 | |
| 34 | 9th January 1931 . . . | 20 | .. | 363 | |
| 35 | 11th January 1931 . . . | 18 | .. | 205 | |
| 36 | 11th January 1931 . . . | 13 | 9 | 81 | |

SPOTTED BOLL-WORMS OF COTTON IN SOUTH GUJARAT

TABLE II

Time of oviposition of the Earias moths. (Observed for 5 days)

| Day of observation | 1st pair | | 2nd pair | | 3rd pair | | 4th pair | | 5th pair | | 6th pair | | 7th pair | | 8th pair | | 9th pair | | 10th pair | | 11th pair | |
|--|---------------|--------------|---------------|--------------|----------------|---------------|----------------|---------------|----------------|---------------|---------------|--------------|---------------|--------------|---------------|--------------|---------------|--------------|---------------|--------------|---------------|--------------|
| | Before 8 P.M. | After 8 P.M. | Before 9 P.M. | After 9 P.M. | Before 10 P.M. | After 10 P.M. | Before 11 P.M. | After 11 P.M. | Before 12 P.M. | After 12 P.M. | Before 1 A.M. | After 1 A.M. | Before 2 A.M. | After 2 A.M. | Before 3 A.M. | After 3 A.M. | Before 4 A.M. | After 4 A.M. | Before 5 A.M. | After 5 A.M. | Before 6 A.M. | After 6 A.M. |
| 1st . | 1 | 15 | .. | 31 | 54 | 12 | 59 | 2 | 42 | 12 | 67 | .. | 70 | 7 | 52 | . | 70 | 4 | 36 | .. | 78 | .. |
| 2nd . | 39 | 82 | 23 | 40 | 42 | 24 | 62 | 3 | 110 | 2 | 108 | 10 | 79 | 8 | 78 | 1 | 55 | 4 | 101 | 2 | 109 | .. |
| 3rd . | 76 | 6 | 46 | 8 | 66 | 21 | 101 | 4 | 96 | 3 | 98 | 15 | 102 | 5 | 59 | 5 | 67 | 7 | 93 | 4 | 90 | .. |
| 4th . | 50 | 35 | 55 | 13 | 60 | 11 | 75 | 11 | 61 | 6 | 59 | 7 | 62 | 4 | 61 | 5 | 39 | 3 | 89 | 1 | 86 | .. |
| 5th . | 64 | 22 | 30 | 7 | 52 | 9 | 62 | 4 | 65 | .. | 52 | .. | 55 | . | 56 | 3 | 10 | .. | 67 | 3 | 71 | .. |
| Total eggs . | 230 | 160 | 154 | 99 | 274 | 77 | 359 | 24 | 374 | 23 | 384 | 32 | 383 | 24 | 396 | 14 | 250 | 18 | 398 | 10 | 454 | .. |
| Percentage oviposition before and after the allotted hour for each pair. | 59 | 41 | 60.9 | 39.1 | 78 | 22 | 93.7 | 6.3 | 94.2 | 5.8 | 88.4 | 11.6 | 93.3 | 6.7 | 92.6 | 4.4 | 93.3 | 6.7 | 97.4 | 2.6 | 100 | .. |

TABLE IV

Periodical bud formation, flower-opening, and shedding of flower-buds and bolls ; and the causes of shedding

(49 plants observed at Surat during 1925-26)

| | 5th September. | 12th September. | 18th September. | 26th September. | 3rd October. | 10th October. | 17th October. | 24th October. | 31st October. | 7th November. | 14th November. | 21st November. | 28th November. | 5th December. | 12th December. | 19th December. | 26th December. | 2nd January. | 9th January. | 16th January. | 23rd January. | 30th January. | 6th February. | TOTAL. |
|---------------------------------------|----------------|-----------------|-----------------|-----------------|--------------|---------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|---------------|----------------|----------------|----------------|--------------|--------------|---------------|---------------|---------------|---------------|--------|
| Flower-bud formation. | 96 | 225 | 339 | 465 | 761 | 837 | 1,192 | 1,494 | 1,254 | 1,415 | 1,523 | 1,294 | 770 | 423 | 186 | 47 | 6 | 5 | . | . | . | . | . | 12,341 |
| Flower-bud shedding . | 8 | 85 | 184 | 266 | 359 | 456 | 829 | 1,050 | 920 | 740 | 758 | 675 | 447 | 493 | 646 | 512 | 927 | 98 | 15 | . | . | . | . | 8,786 |
| Flower-buds shed due to boll-worms. | .. | . | 41 | 71 | 110 | 201 | 436 | 571 | 503 | 502 | 371 | 357 | 203 | 196 | 146 | 77 | 22 | 2 | .. | .. | .. | .. | .. | 3,899 |
| Flower-buds shed due to other causes. | 5 | 62 | 117 | 173 | 217 | 216 | 358 | 447 | 290 | 211 | 233 | 286 | 221 | 252 | 424 | 381 | 170 | 66 | .. | .. | .. | .. | .. | 4,129 |
| Flower opening . | .. | .. | .. | .. | .. | .. | 22 | 22 | 15 | 17 | 85 | 214 | 562 | 757 | 764 | 643 | 378 | 128 | 25 | 3 | .. | .. | .. | 3,835 |
| Total boll shedding . | .. | .. | .. | .. | .. | .. | 7 | 20 | 15 | 11 | 17 | 34 | 56 | 161 | 278 | 425 | 595 | 484 | 118 | 24 | 11 | 1 | .. | 2,257 |
| Bolls shed due to boll-worms. | .. | .. | .. | .. | .. | .. | 7 | 18 | 13 | 8 | 15 | 21 | 40 | 88 | 75 | 66 | 74 | 43 | 14 | 3 | 1 | .. | .. | 486 |
| Bolls shed due to other causes. | .. | .. | .. | .. | .. | .. | .. | 2 | 2 | 3 | 2 | 13 | 18 | 73 | 203 | 359 | 521 | 411 | 102 | 21 | 10 | 1 | .. | 1,771 |

TABLE V

Periodical flower-bud formation, flower-opening, and shedding of flower-buds and bolls; with causes of their shedding
(20 plants observed during 1926-27 at Surat)

| | 2nd October. | 9th October. | 10th October. | 23rd October. | 30th October. | 6th November. | 13th November. | 20th November. | 27th November. | 4th December. | 11th December. | 18th December. | 25th December. | 1st January. | 8th January. | 15th January. | 22nd January. | 29th January. | 5th February. | 12th February. | 19th February. | 26th February. | Total. | |
|--|--------------|--------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|---------------|----------------|----------------|----------------|--------------|--------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|--------|-------|
| Flower-bud formation . . . | 29 | 55 | 110 | 284 | 452 | 511 | 614 | 656 | 506 | 677 | 544 | 407 | 225 | 254 | 130 | 119 | 86 | 45 | 26 | 8 | 12 | 15 | .. | 5,945 |
| Flower-bud shedding (total) . . . | .. | 6 | 34 | 31 | 96 | 148 | 274 | 323 | 441 | 414 | 455 | 528 | 476 | 324 | 281 | 136 | 161 | 183 | 194 | 157 | 68 | 18 | 15 | 4,766 |
| Flower-buds shed due to boll-worms . . . | . | . | 6 | 3 | 14 | 40 | 109 | 157 | 260 | 247 | 267 | 328 | 276 | 189 | 162 | 67 | 62 | 49 | 39 | 11 | 2 | .. | .. | 2,238 |
| Flower-buds shed due to other causes . . . | .. | 3 | 26 | 17 | 52 | 85 | 129 | 140 | 140 | 143 | 164 | 185 | 183 | 122 | 112 | 62 | 93 | 127 | 149 | 143 | 61 | 16 | 13 | 2,165 |
| Flower opening | .. | .. | .. | .. | 3 | 7 | 32 | 40 | 40 | 71 | 67 | 88 | 80 | 68 | 96 | 85 | 92 | 114 | 125 | 115 | 40 | 8 | .. | 1,180 |
| Boll-shedding | . | .. | .. | . | . | .. | 8 | 14 | 23 | 37 | 55 | 62 | 81 | 65 | 60 | 45 | 38 | 38 | 55 | 46 | 64 | 10 | 5 | 706 |
| Bolls shed due to boll-worms | .. | . | . | . | . | . | 6 | 10 | 21 | 32 | 46 | 56 | 74 | 58 | 52 | 36 | 29 | 17 | 27 | 12 | 6 | .. | .. | 482 |
| Bolls shed due to other causes | .. | .. | . | .. | .. | . | 2 | 4 | 2 | 5 | 9 | 6 | 7 | 7 | 8 | 9 | 9 | 21 | 28 | 34 | 58 | 10 | 5 | 224 |

TABLE VI

Periodical flower-bud formation, flower-opening and shedding of flower-buds and bolls ; with causes of their shedding

(16 plants observed at Surat during 1927-28)

| | 17th September. | 24th September. | 1st October. | 8th October. | 15th October. | 22nd October. | 29th October. | 5th November. | 12th November. | 19th November. | 26th November. | 3rd December. | 10th December. | 17th December. | 24th December. | 31st December. | 7th January. | 14th January. | 21st January. | 28th January. | 4th February. | 11th February. | 18th February. | 25th February. | Total. |
|---------------------------------------|-----------------|-----------------|--------------|--------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|---------------|----------------|----------------|----------------|----------------|--------------|---------------|---------------|---------------|---------------|----------------|----------------|----------------|--------|
| Flower-bud formation . | 114 | 187 | 317 | 317 | 279 | 349 | 278 | 574 | 612 | 473 | 493 | 573 | 563 | 474 | 427 | 239 | 156 | 68 | 49 | 24 | 3 | .. | 1 | .. | 6,570 |
| Flower-bud shedding (total) . | 26 | 98 | 225 | 225 | 180 | 352 | 239 | 239 | 299 | 319 | 310 | 302 | 351 | 219 | 187 | 290 | 276 | 301 | 243 | 195 | 70 | 54 | 27 | .. | 5,027 |
| Flower-buds shed due to boll-worms. | 3 | 24 | 12 | 12 | 18 | 75 | 55 | 88 | 131 | 95 | 116 | 153 | 163 | 91 | 65 | 46 | 69 | 37 | 26 | 13 | 9 | 2 | .. | .. | 1,308 |
| Flower-buds shed due to other causes. | 14 | 52 | 121 | 121 | 124 | 234 | 143 | 78 | 84 | 84 | 117 | 69 | 130 | 102 | 105 | 207 | 167 | 210 | 191 | 154 | 51 | 39 | 17 | .. | 2,594 |
| Flower opening . | .. | .. | .. | .. | .. | .. | .. | .. | 1 | 6 | 12 | 31 | 67 | 67 | 116 | 142 | 193 | 279 | 223 | 221 | 109 | 46 | 22 | 2 | 1,542 |
| Bolls shed (Total) . | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | 5 | 4 | 23 | 30 | 24 | 68 | 49 | 91 | 124 | 128 | 156 | 156 | 71 | 273 | 950 |
| Bolls shed due to boll-worms . | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | 5 | 3 | 17 | 19 | 17 | 84 | 18 | 31 | 25 | 11 | 4 | 4 | .. | 3 | 192 |
| Bolls shed due to other causes . | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | 6 | 11 | 7 | 24 | 31 | 60 | 99 | 117 | 152 | 152 | 71 | 243 | 758 |

TABLE VII

Rainfall at Surat for 6 years. (June to December)

| 1925 | | | 1926 | | | 1927 | | | 1928 | | | 1929 | | | 1930 | | |
|-------------|---|-------|-------------|---|-------|-------------|---|-------|-------------|---|-------|-------------|---|-------|-------------|---|-------|
| Weeks | | Rain | Weeks | | Rain | Weeks | | Rain | Weeks | | Rain | Weeks | | Rain | Weeks | | Rain |
| 6th June | • | 8.40 | 6th June | • | • | 4th June | • | 0.45 | 2nd June | • | • | 1st June | • | • | 30th May | • | • |
| 13th June | • | 9.41 | 13th June | • | • | 11th June | • | 4.73 | 9th June | • | • | 8th June | • | • | 6th June | • | • |
| 20th June | • | 3.76 | 20th June | • | • | 18th June | • | 3.86 | 16th June | • | • | 15th June | • | • | 13th June | • | • |
| 27th June | • | • | 27th June | • | • | 25th June | • | • | 23rd June | • | • | 22nd June | • | • | 20th June | • | 1.15 |
| 4th July | • | 1.27 | 4th July | • | 0.17 | 2nd July | • | 0.98 | 30th June | • | • | 29th June | • | • | 27th June | • | 7.55 |
| 11th July | • | 0.59 | 11th July | • | 4.45 | 9th July | • | 3.39 | 7th July | • | • | 6th July | • | • | 4th July | • | 7.55 |
| 18th July | • | 1.12 | 18th July | • | 6.73 | 16th July | • | 2.48 | 14th July | • | • | 13th July | • | • | 11th July | • | 15.36 |
| 25th July | • | 0.53 | 25th July | • | 0.85 | 23rd July | • | 1.18 | 21st July | • | • | 20th July | • | • | 18th July | • | 0.85 |
| 1st August | • | 0.53 | 1st August | • | 11.99 | 30th July | • | 5.14 | 28th July | • | • | 27th July | • | • | 25th July | • | 3.03 |
| 8th August | • | 3.32 | 8th August | • | 0.76 | 6th August | • | 0.91 | 4th August | • | • | 3rd August | • | • | 1st August | • | 1.88 |
| 15th August | • | 0.50 | 15th August | • | 10.66 | 13th August | • | 0.97 | 11th August | • | • | 10th August | • | • | 8th August | • | 0.61 |
| 22nd August | • | 0.25 | 22nd August | • | 1.77 | 20th August | • | 1.13 | 18th August | • | • | 17th August | • | • | 15th August | • | 0.17 |
| 29th August | • | 0.08 | 29th August | • | 5.12 | 27th August | • | 0.70 | 25th August | • | • | 24th August | • | • | 22nd August | • | • |
| 5th Sept. | • | 0.45 | 5th Sept. | • | 3.05 | 3rd Sept. | • | 0.55 | 1st Sept. | • | • | 31st August | • | • | 29th August | • | • |
| 12th Sept. | • | 0.60 | 12th Sept. | • | 9.59 | 10th Sept. | • | 0.42 | 8th Sept. | • | • | 7th Sept. | • | • | 5th Sept. | • | 0.16 |
| 19th Sept. | • | • | 19th Sept. | • | 9.42 | 17th Sept. | • | 0.46 | 15th Sept. | • | • | 14th Sept. | • | • | 12th Sept. | • | 1.07 |
| 26th Sept. | • | • | 26th Sept. | • | 1.25 | 24th Sept. | • | 0.74 | 22nd Sept. | • | • | 21st Sept. | • | • | 19th Sept. | • | 4.20 |
| 3rd Oct. | • | • | 3rd Oct. | • | • | 1st Oct. | • | 2.51 | 29th Sept. | • | • | 28th Sept. | • | • | 26th Sept. | • | 0.06 |
| 10th Oct. | • | • | 10th Oct. | • | • | 8th Oct. | • | 2.15 | 6th Oct. | • | • | 5th Oct. | • | • | 3rd Oct. | • | • |
| 17th Oct. | • | • | 17th Oct. | • | • | 15th Oct. | • | • | 13th Oct. | • | • | 12th Oct. | • | • | 10th Oct. | • | • |
| 24th Oct. | • | • | 24th Oct. | • | • | 22nd Oct. | • | • | 20th Oct. | • | • | 19th Oct. | • | • | 17th Oct. | • | • |
| 31st Oct. | • | • | 31st Oct. | • | • | 29th Oct. | • | • | 27th Oct. | • | • | 26th Oct. | • | • | 24th Oct. | • | • |
| 7th Nov. | • | • | 7th Nov. | • | • | 5th Nov. | • | • | 3rd Nov. | • | • | 2nd Nov. | • | • | 31st Oct. | • | 0.95 |
| 14th Nov. | • | • | 14th Nov. | • | • | 12th Nov. | • | 1.18 | 10th Nov. | • | • | 9th Nov. | • | • | 7th Nov. | • | • |
| 21st Nov. | • | 0.06 | 21st Nov. | • | • | 19th Nov. | • | 1.70 | 17th Nov. | • | • | 16th Nov. | • | • | 14th Nov. | • | • |
| 28th Nov. | • | • | 28th Nov. | • | • | 26th Nov. | • | • | 24th Nov. | • | • | 23rd Nov. | • | • | 21st Nov. | • | • |
| 4th Dec. | • | • | 4th Dec. | • | • | 3rd Dec. | • | • | 1st Dec. | • | • | 30th Nov. | • | • | 28th Nov. | • | • |
| 11th Dec. | • | • | 11th Dec. | • | • | 10th Dec. | • | • | 8th Dec. | • | • | 7th Dec. | • | • | 5th Dec. | • | 0.10 |
| 18th Dec. | • | • | 18th Dec. | • | • | 17th Dec. | • | • | 15th Dec. | • | • | 14th Dec. | • | • | 12th Dec. | • | • |
| TOTAL | • | 25.44 | | • | 62.95 | | • | 35.06 | | • | 26.21 | | • | 48.26 | | • | 41.28 |

TABLE VIII

Population of Earias larvae on cotton plants at Surat

(25 cotton plants examined every week, 1925-26)

| Week ending | No. of plants | <i>Earias</i> larvae in | | | Total <i>Earias</i> | Empty buds | Total buds | Empty bolls | Total bolls |
|--------------------------|---------------|-------------------------|------|-------|---------------------|------------|------------|-------------|-------------|
| | | Shoots | Buds | Bolls | | | | | |
| 19th September 1925 . | 25 | 28 | 4 | .. | 32 | 56 | 248 | .. | .. |
| 25th September 1925 . . | 25 | 44 | 4 | .. | 48 | 80 | 355 | .. | .. |
| 2nd October 1925 . . . | 25 | 42 | 6 | .. | 48 | 99 | 427 | .. | .. |
| 9th October 1925 . . . | 25 | 27 | 16 | .. | 43 | 106 | 662 | .. | .. |
| 16th October 1925 . . . | 25 | 54 | 14 | .. | 68 | 136 | 633 | .. | .. |
| 23rd October 1925 . . . | 25 | 58 | 29 | .. | 87 | 230 | 692 | .. | .. |
| 30th October 1925 . . . | 25 | 81 | 42 | .. | 73 | 200 | 856 | .. | .. |
| 7th November 1925 . . . | 25 | 11 | 34 | .. | 45 | 185 | 1,042 | .. | .. |
| 14th November 1925 . . . | 25 | 5 | 40 | .. | 45 | 156 | 1,040 | 1 | .. |
| 21st November 1925 . . . | 25 | 2 | 27 | 3 | 32 | 202 | 2,221 | 4 | 44 |
| 28th November 1925 . . . | 25 | .. | 27 | 27 | 54 | 90 | 2,341 | 22 | 181 |
| 4th December 1925 . . . | 25 | .. | 26 | 21 | 47 | 40 | 2,460 | 20 | 334 |
| 11th December 1925 . . . | 25 | .. | 18 | 26 | 44 | 31 | 1,684 | 16 | 420 |
| 18th December 1925 . . . | 25 | .. | 18 | 37 | 55 | 18 | 1,577 | 35 | 606 |
| 25th December 1925 . . . | 25 | .. | 13 | 38 | 51 | 12 | 927 | 60 | 1,186 |
| 1st January 1926 . . . | 25 | .. | 3 | 35 | 38 | 4 | 401 | 54 | 1,127 |
| 8th January 1926 . . . | 25 | .. | 3 | 27 | 30 | 1 | 97 | 32 | 966 |
| 15th January 1926 . . . | 25 | .. | .. | 22 | 22 | .. | 49 | 44 | 821 |
| 22nd January 1926 . . . | 25 | .. | .. | 25 | 25 | .. | .. | 54 | 838 |
| 29th January 1926 . . . | 25 | .. | .. | 14 | 14 | .. | 24 | 31 | 795 |

TABLE IX

Population of Earias larvae on cotton plants at Surat

(25 plants examined every week, 1926-27)

| Week ending | No. of plants | Earias larvae in | | | Total Earias | Empty buds | Total buds | Empty bolls | Total bolls |
|--------------------------|---------------|------------------|------|-------|--------------|------------|------------|-------------|-------------|
| | | Shoots | Buds | Bolls | | | | | |
| 9th October 1926 . . . | 25 | .. | .. | .. | .. | 1 | 56 | .. | .. |
| 16th October 1926 . . . | 25 | 2 | 1 | .. | 3 | 2 | 97 | .. | .. |
| 23rd October 1926 . . . | 25 | 5 | .. | .. | 5 | 2 | 162 | .. | .. |
| 30th October 1926 . . . | 25 | 15 | 1 | .. | 16 | 9 | 517 | .. | .. |
| 6th November 1926 . . . | 25 | 9 | 10 | .. | 19 | 75 | 804 | .. | .. |
| 13th November 1926 . . . | 25 | 9 | 28 | 1 | 38 | 84 | 1,065 | .. | 4 |
| 20th November 1926 . . . | 25 | 18 | 44 | .. | 62 | 103 | 1,349 | .. | 2 |
| 27th November 1926 . . . | 25 | 17 | 73 | 1 | 91 | 178 | 1,367 | 1 | 4 |
| 4th December 1926 . . . | 25 | 15 | 59 | 2 | 76 | 157 | 1,445 | .. | 6 |
| 11th December 1926 . . . | 25 | 18 | 115 | 5 | 138 | 167 | 1,505 | .. | 9 |
| 18th December 1926 . . . | 25 | 21 | 99 | 3 | 123 | 305 | 2,116 | 1 | 10 |
| 25th December 1926 . . . | 25 | 12 | 124 | 8 | 144 | 311 | 1,947 | 7 | 24 |
| 1st January 1927 . . . | 25 | .. | .. | .. | .. | .. | .. | .. | .. |
| 8th January 1927 . . . | 25 | 2 | 60 | 5 | 67 | 160 | 2,114 | 4 | 25 |
| 15th January 1927 . . . | 25 | 1 | 73 | 12 | 86 | 215 | 2,220 | 4 | 58 |
| 22nd January 1927 . . . | 25 | .. | 44 | 3 | 47 | 114 | 2,165 | 3 | 99 |
| 29th January 1927 . . . | 25 | 1 | 31 | 15 | 47 | 95 | 2,090 | 8 | 229 |
| 5th February 1927 . . . | 25 | .. | 21 | 23 | 44 | 74 | 1,890 | 19 | 439 |
| 12th February 1927 . . . | 25 | .. | 10 | 5 | 15 | 53 | 1,991 | 11 | 794 |
| 19th February 1927 . . . | 25 | .. | 9 | 12 | 21 | 28 | 1,271 | 32 | 1,081 |
| 26th February 1927 . . . | 25 | .. | 2 | 7 | 9 | 10 | 682 | 15 | 1,071 |
| 5th March 1927 . . . | 25 | .. | 1 | 4 | 5 | 1 | 195 | 11 | 888 |
| 12th March 1927 . . . | 25 | .. | .. | 5 | 5 | .. | 9 | 21 | 801 |
| 19th March 1927 . . . | 25 | .. | .. | 4 | 4 | .. | .. | 40 | 781 |
| 26th March 1927 . . . | 25 | .. | .. | 7 | 7 | .. | .. | 78 | 690 |

TABLE X

Population of Earias larvae on cotton plants at Surat

(25 plants examined every week, 1927-28)

| Week ending | <i>Earias</i> larvae in | | | Total <i>Earias</i> | Empty buds | Total buds | Empty bolls | Total bolls | REMARKS |
|---------------------|-------------------------|------|-------|------------------------|---------------|---------------|----------------|----------------|--------------------|
| | Shoots | Buds | Bolls | | | | | | |
| 17th September 1927 | 7 | 1 | .. | 8 | .. | 166 | .. | .. | |
| 24th September 1927 | 22 | 3 | .. | 25 | 6 | 333 | .. | .. | |
| 1st October 1927 | 23 | 1 | .. | 24 | 15 | 432 | .. | .. | |
| 8th October 1927 | 21 | 1 | .. | 22 | 28 | 588 | .. | .. | |
| 15th October 1927 | 68 | 6 | .. | 74 | 56 | 777 | .. | .. | |
| 22nd October 1927 | 40 | 8 | .. | 57 | 71 | 819 | .. | .. | |
| 29th October 1927 | 46 | 12 | .. | 61 | 169 | 1,158 | .. | .. | 3 moving on plant. |
| 5th November 1927 | 31 | 20 | .. | 51 | 141 | 1,263 | .. | .. | |
| 12th November 1927 | 38 | 40 | .. | 78 | 227 | 1,976 | .. | .. | |
| 19th November 1927 | 6 | 33 | 1 | 46 | 171 | 2,392 | 1 | 2 | 1 moving on plant. |
| 26th November 1927 | 3 | 65 | 1 | 69 | 197 | 2,421 | .. | 13 | |
| 3rd December 1927 | 5 | 55 | 3 | 63 | 122 | 2,302 | 5 | 38 | |
| 10th December 1927 | .. | 35 | 7 | 42 | 123 | 2,496 | 3 | 107 | |
| 17th December 1927 | .. | 30 | 4 | 34 | 130 | 3,501 | 7 | 170 | |
| 24th December 1927 | .. | 23 | 11 | 34 | 111 | 3,541 | 8 | 189 | |
| 31st December 1927 | .. | 29 | 12 | 41 | 79 | 3,530 | 23 | 406 | |
| 7th January 1928 | .. | 22 | 16 | 38 | 66 | 3,204 | 11 | 487 | |
| 14th January 1928 | .. | 12 | 23 | 35 | 51 | 2,763 | 60 | 1,156 | |
| 21st January 1928 | .. | 3 | 15 | 18 | 32 | 1,840 | 44 | 1,164 | |
| 28th January 1928 | .. | 1 | 11 | 12 | 34 | 968 | 86 | 1,357 | |
| 4th February 1928 | .. | 1 | 10 | 11 | 26 | 607 | 80 | 1,629 | |
| 11th February 1928 | .. | .. | 4 | 4 | 1 | 105 | 70 | 1,270 | |
| 18th February 1928 | .. | .. | 2 | 2 | .. | 56 | 47 | 1,001 | |
| 25th February 1928 | .. | 1 | 7 | 8 | .. | 14 | 59 | 857 | |
| 3rd March 1928 | .. | .. | 2 | 2 | .. | 4 | 51 | 527 | |
| 10th March 1928 | .. | .. | 9 | 9 | .. | .. | 39 | 300 | |
| 17th March 1928 | .. | .. | 15 | 15 | .. | .. | 23 | 183 | |
| 24th March 1928 | .. | .. | 7 | 7 | .. | .. | 10 | 72 | |
| 31st March 1928 | 1 | .. | 6 | 7 | 2 | 27 | 3 | 25 | |

TABLE XI

Population of Earias larvae on cotton plants at Surat
(1928-29)

| Week ending | No. of plants | Earias larvae in | | | Total Earias | Empty buds | Total buds | Empty bolls | Total bolls | REMARKS |
|---------------------|---------------|------------------|------|-------|--------------|------------|------------|-------------|-------------|-----------|
| | | Shoots | Buds | Bolls | | | | | | |
| 4th August 1928 | 1,000 | .. | .. | .. | .. | .. | .. | .. | .. | |
| 11th August 1928 | 1,000 | .. | .. | .. | .. | .. | .. | .. | .. | |
| 18th August 1928 | 1,000 | .. | .. | .. | .. | .. | .. | .. | .. | |
| 25th August 1928 | 750 | .. | .. | .. | .. | .. | 3 | .. | .. | |
| 1st September 1928 | 750 | 5 | .. | .. | 5 | .. | 9 | .. | .. | |
| 8th September 1928 | 250 | 4 | .. | .. | 4 | .. | 21 | .. | .. | |
| 15th September 1928 | 125 | 9 | .. | .. | 9 | .. | 90 | .. | .. | |
| 22nd September 1928 | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 29th September 1928 | 100 | 4 | .. | .. | 4 | 2 | 274 | .. | .. | |
| 6th October 1928 | 50 | 6 | 1 | .. | 7 | 2 | 203 | .. | .. | |
| 13th October 1928 | 25 | 26 | 3 | .. | 29 | 14 | 562 | .. | .. | |
| 20th October 1928 | 25 | 30 | 15 | .. | 45 | 40 | 1,113 | .. | .. | |
| 27th October 1928 | 25 | 31 | 24 | .. | 56 | 61 | 1,845 | .. | .. | |
| 3rd November 1928 | 25 | 10 | 22 | .. | 32 | 92 | 1,758 | .. | 1 | |
| 10th November 1928 | 25 | 13 | 75 | 1 | 89 | 135 | 2,758 | .. | .. | |
| 17th November 1928 | 25 | 6 | 67 | 10 | 86 | 108 | 2,767 | 4 | 50 | 3 moving. |
| 24th November 1928 | 25 | 2 | 84 | 8 | 95 | 86 | 3,096 | 7 | 119 | 1 moving. |
| 1st December 1928 | 25 | .. | 100 | 30 | 130 | 90 | 3,235 | 12 | 210 | |
| 8th December 1928 | 25 | .. | 72 | 43 | 115 | 75 | 3,201 | 26 | 314 | |
| 15th December 1928 | 25 | 2 | 80 | 32 | 123 | 74 | 2,797 | 14 | 338 | |
| 22nd December 1928 | 25 | .. | 34 | 30 | 65 | 59 | 2,915 | 20 | 610 | |
| 29th December 1928 | 25 | .. | 22 | 15 | 37 | 12 | 3,162 | 19 | 863 | |
| 5th January 1929 | 25 | .. | 21 | 19 | 40 | 6 | 1,744 | 12 | 863 | |
| 12th January 1929 | 25 | .. | 8 | 11 | 19 | 5 | 1,025 | 26 | 851 | |
| 19th January 1929 | 25 | .. | 8 | 10 | 18 | 5 | 804 | 42 | 1,062 | |
| 26th January 1929 | 25 | .. | 4 | 16 | 20 | 4 | 681 | 26 | 1,005 | |
| 2nd February 1929 | 25 | .. | 1 | 24 | 25 | 2 | 188 | 16 | 1,027 | |
| 9th February 1929 | 25 | .. | .. | 19 | 19 | .. | 30 | 5 | 794 | |
| 16th February 1929 | 25 | .. | 1 | 45 | 46 | .. | 3 | 1 | 761 | |
| 23rd February 1929 | 25 | .. | 1 | 53 | 54 | .. | 1 | .. | 586 | |
| 2nd March 1929 | 25 | .. | .. | 56 | 56 | .. | .. | 12 | 416 | |
| 9th March 1929 | 25 | .. | .. | 28 | 28 | .. | .. | .. | 175 | |
| 16th March 1929 | 25 | .. | .. | 17 | 17 | .. | 5 | .. | 35 | |
| 23rd March 1929 | 25 | .. | .. | 4 | 4 | .. | 17 | .. | 10 | |
| 30th March 1929 | 50 | .. | 2 | .. | 2 | 2 | 75 | .. | 2 | |

Population of Earias larvae on cotton plants at Surat
(1929-30)

[illegible]

TABLE XIII

Population of Earias larvae on cotton plants at Surat

(1930-31)

| Week ending | No. of plants | Earias larvae in | | | Total Earias | Empty buds | Total buds | Empty bolls | Total bolls | REMARKS |
|---------------------|---------------|------------------|------|-------|--------------|------------|------------|-------------|-------------|-------------------------|
| | | Shoots | Buds | Bolls | | | | | | |
| 13th August 1930 | 1,000 | .. | .. | .. | .. | .. | .. | .. | .. | |
| 20th August 1930 | 1,000 | 2 | .. | .. | 2 | .. | .. | .. | .. | |
| 27th August 1930 | 1,000 | 5 | .. | .. | 5 | .. | .. | .. | .. | |
| 3rd September 1930 | 500 | 18 | .. | .. | 18 | .. | .. | .. | .. | |
| 10th September 1930 | 200 | 8 | .. | .. | 8 | .. | 124 | .. | .. | |
| 17th September 1930 | 100 | 6 | .. | .. | 7 | .. | 62 | .. | .. | 1 moving |
| 24th September 1930 | 50 | 32 | 1 | .. | 36 | .. | 470 | .. | .. | 3 moving |
| 3rd October 1930 | 25 | 8 | 4 | .. | 13 | 11 | 565 | .. | .. | 1 <i>Earias</i> moving. |
| 10th October 1930 | 25 | 9 | 17 | .. | 28 | 36 | 1,301 | .. | .. | 2 <i>Earias</i> moving. |
| 17th October 1930 | 25 | 17 | 57 | .. | 78 | 86 | 2,202 | .. | 3 | 4 <i>Earias</i> moving. |
| 24th October 1930 | 25 | 24 | 158 | 9 | 196 | 155 | 2,260 | 7 | 50 | 5 <i>Earias</i> moving. |
| 31st October 1930 | 25 | 43 | 155 | 14 | 220 | 186 | 1,977 | 5 | 43 | 8 <i>Earias</i> moving. |
| 7th November 1930 | 25 | 31 | 122 | 12 | 167 | 174 | 1,777 | 7 | 51 | 2 <i>Earias</i> moving. |
| 14th November 1930 | 25 | 11 | 123 | 6 | 140 | 190 | 2,107 | 14 | 33 | |
| 21st November 1930 | 25 | 1 | 78 | 9 | 90 | 97 | 2,688 | 2 | 32 | 2 <i>Earias</i> moving. |
| 28th November 1930 | 25 | .. | 90 | 15 | 108 | .. | 3,998 | .. | 75 | 3 <i>Earias</i> moving. |
| 5th December 1930 | 25 | .. | 132 | 24 | 167 | 94 | 5,192 | 7 | 208 | 1 <i>Earias</i> moving. |
| 12th December 1930 | 25 | .. | 91 | 28 | 122 | 67 | 4,224 | 16 | 288 | 3 <i>Earias</i> moving. |
| 19th December 1930 | 25 | .. | 52 | 33 | 85 | 71 | 4,207 | 35 | 868 | |
| 26th December 1930 | 25 | .. | 33 | 19 | 53 | 53 | 3,910 | 37 | 1,707 | 1 <i>Earias</i> moving. |
| 2nd January 1931 | 25 | .. | 13 | 17 | 30 | 17 | 2,198 | 24 | 1,895 | |
| 9th January 1931 | 25 | .. | 6 | 17 | 23 | 5 | 927 | 33 | 2,167 | |
| 16th January 1931 | 25 | .. | 1 | 13 | 14 | 1 | 225 | 37 | 1,692 | |
| 23rd January 1931 | 25 | .. | .. | 20 | 20 | .. | 18 | 26 | 1,248 | |
| 30th January 1931 | 25 | .. | .. | 28 | 28 | .. | 1 | 38 | 1,165 | |

TABLE XIV

Weekly flower-bud formation, flower-opening and shedding of flower-buds and bolls from the caged and control plants

(1925-26)

| | 4th September | 11th September | 18th September | 25th September | 2nd October | 9th October | 16th October | 23rd October | 30th October | 6th November | 13th November | 20th November | 27th November | 4th December | 11th December | 18th December | 25th December | 1st January | 8th January | 15th January | TOTAL |
|---------------------------------------|---------------|----------------|----------------|----------------|-------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|-------------|-------------|--------------|--------|
| CAGED PLANTS 8. | | | | | | | | | | | | | | | | | | | | | |
| Flower-bud formation . . . | 27 | 72 | 91 | 109 | 223 | 239 | 292 | 263 | 237 | 172 | 60 | 18 | 20 | 8 | .. | .. | .. | .. | .. | .. | 1,831 |
| Flower-bud shedding . . . | 2 | 19 | 34 | 47 | 36 | 46 | 68 | 101 | 111 | 179 | 193 | 228 | 14 | 28 | 16 | .. | .. | .. | .. | .. | 1,122 |
| Flower-opening . . . | .. | .. | .. | .. | .. | 1 | 16 | 46 | 133 | 178 | 188 | 113 | 29 | 3 | 3 | .. | .. | .. | .. | .. | 709 |
| UNCAGED PLANTS 40. | | | | | | | | | | | | | | | | | | | | | |
| Flower-bud formation . . . | 96 | 255 | 339 | 465 | 761 | 837 | 1,192 | 1,494 | 1,254 | 1,115 | 1,523 | 1,204 | 779 | 423 | 186 | 47 | 6 | 5 | .. | .. | 12,341 |
| Flower-bud shedding (Total) . . | 8 | 85 | 184 | 266 | 359 | 456 | 829 | 1,059 | 920 | 749 | 758 | 675 | 1,147 | 493 | 646 | 512 | 227 | 98 | 15 | .. | 8,786 |
| Flower-buds shed due to boll-worms . | .. | .. | 41 | 71 | 110 | 201 | 436 | 571 | 593 | 592 | 371 | 357 | 203 | 196 | 146 | 77 | 22 | 2 | .. | .. | 3,999 |
| Flower-buds shed due to other causes. | 5 | 62 | 117 | 173 | 217 | 216 | 353 | 447 | 290 | 211 | 233 | 286 | 221 | 252 | 424 | 381 | 170 | 66 | .. | .. | 4,129 |
| Flower-opening . . . | .. | .. | .. | .. | .. | .. | 22 | 22 | 15 | 17 | 85 | 214 | 562 | 757 | 764 | 643 | 378 | 128 | 25 | 3 | 3,645 |

TABLE XV

Weekly flower-bud formation, flower-opening and shedding of flower-buds and bolls from the caged and control plants

(1926-27)

| | 2nd October | 9th October | 17th October | 23rd October | 30th October | 6th November | 13th November | 20th November | 27th November | 4th December | 11th December | 18th December | 25th December | 1st January | 8th January | 15th January | 22nd January | 29th January | 5th February | 12th February | 19th February | 26th February | 5th March | Total |
|--------------------------------------|-------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|-------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|-----------|-------|
| CAGED PLANTS 10. | | | | | | | | | | | | | | | | | | | | | | | | |
| Flower-bud formation . . . | .. | 112 | 156 | 189 | 276 | 368 | 341 | 317 | 282 | 238 | 128 | 38 | 29 | 4 | 2 | .. | .. | .. | .. | .. | .. | .. | .. | 2,430 |
| Flower-bud shedding . . . | .. | 7 | 20 | 32 | 24 | 61 | 87 | 160 | 172 | 168 | 218 | 306 | 188 | 79 | 9 | 8 | .. | .. | .. | .. | .. | .. | .. | 1,639 |
| Flower-opening . . . | .. | .. | .. | .. | 7 | 30 | 60 | 97 | 108 | 133 | 129 | 148 | 106 | 49 | 10 | 4 | .. | 1 | .. | .. | .. | .. | .. | 891 |
| UNCAGED PLANTS 20. | | | | | | | | | | | | | | | | | | | | | | | | |
| Flower-bud formation . . . | 29 | 55 | 110 | 284 | 452 | 511 | 614 | 676 | 596 | 677 | 544 | 407 | 225 | 254 | 130 | 119 | 86 | 45 | 28 | 8 | 12 | 15 | .. | 5,945 |
| Flower-bud shedding (Total) . . | .. | 6 | 34 | 31 | 96 | 148 | 274 | 326 | 441 | 414 | 455 | 628 | 476 | 324 | 251 | 136 | 161 | 183 | 194 | 157 | 68 | 18 | 15 | 4,766 |
| Flower-bud shed due to boll-worms | .. | .. | 6 | 3 | 14 | 40 | 109 | 157 | 260 | 247 | 267 | 323 | 276 | 189 | 162 | 67 | 62 | 49 | 39 | 11 | 2 | .. | .. | 2,288 |
| Flower-buds shed due to other causes | .. | 3 | 26 | 17 | 52 | 85 | 129 | 140 | 140 | 143 | 164 | 185 | 183 | 122 | 112 | 62 | 93 | 127 | 149 | 143 | 61 | 16 | 13 | 2,165 |
| Flower-opening . . . | .. | .. | .. | .. | 3 | 7 | 32 | 40 | 40 | 71 | 67 | 88 | 89 | 68 | 96 | 85 | 92 | 114 | 125 | 115 | 40 | 8 | .. | 1,180 |

TABLE XVI

Weekly flower-bud formation, flower-opening and shedding of flower-buds and bolls from the caged and control plants

(1927-28)

| | 9th September | 16th September | 23rd September | 30th September | 7th October | 14th October | 21st October | 28th October | 5th November | 12th November | 19th November | 26th November | 3rd December | 10th December | 17th December | 24th December | 31st December | 7th January | 14th January | 21st January | 28th January | 4th February | 11th February | 18th February | 25th February | TOTAL |
|---------------------------------------|---------------|----------------|----------------|----------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|-------|
| CAGED PLANTS 20. | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Flower-bud formation. | 11 | 117 | 220 | 478 | 478 | 375 | 647 | 717 | 1,219 | 1,021 | 508 | 743 | 802 | 1,085 | 360 | 149 | 28 | 1 | .. | .. | .. | .. | .. | .. | .. | 8,909 |
| Flower-bud shedding. | .. | 25 | 83 | 339 | 339 | 305 | 384 | 343 | 317 | 229 | 204 | 609 | 115 | 230 | 389 | 518 | 799 | 194 | 102 | 36 | .. | .. | .. | .. | .. | 5,005 |
| Flower-opening | .. | .. | .. | .. | .. | .. | .. | .. | .. | 23 | 111 | 205 | 304 | 448 | 450 | 823 | 570 | 409 | 202 | 63 | 19 | 6 | 1 | .. | .. | 3,334 |
| UNCAGED PLANTS 16. | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Flower-bud formation. | .. | 114 | 187 | 317 | 317 | 279 | 349 | 278 | 574 | 612 | 473 | 493 | 573 | 563 | 474 | 427 | 239 | 156 | 68 | 49 | 24 | 3 | .. | 1 | .. | 6,570 |
| Flower-bud shedding (Total). | .. | 26 | 98 | 225 | 225 | 180 | 352 | 239 | 239 | 299 | 319 | 310 | 302 | 351 | 219 | 187 | 290 | 276 | 301 | 243 | 195 | 70 | 54 | 27 | .. | 5,927 |
| Flower-buds shed due to boll-worms | .. | 3 | 24 | 12 | 12 | 18 | 75 | 55 | 88 | 131 | 95 | 116 | 158 | 163 | 91 | 65 | 46 | 69 | 37 | 26 | 13 | 9 | 2 | .. | .. | 1,308 |
| Flower-buds shed due to other causes. | .. | 14 | 52 | 121 | 121 | 124 | 234 | 143 | 78 | 84 | 64 | 117 | 69 | 130 | 102 | 105 | 207 | 167 | 210 | 191 | 154 | 51 | 39 | 17 | .. | 2,594 |
| Flower-opening | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | 6 | 12 | 31 | 67 | 67 | 116 | 142 | 108 | 279 | 223 | 221 | 109 | 46 | 22 | 2 | 1,542 |

TABLE XVII

Flower-opening, relative success from flowers to mature bolls and shedding of bolls with causes from caged plants and their controls

(1925-26, Surat)

| | 3rd October | 10th October | 17th October | 24th October | 31st October | 7th November | 14th November | 21st November | 28th November | 5th December | 12th December | 19th December | 26th December | 2nd January | 9th January | 16th January | 23rd January | 30th January | TOTAL |
|---|-------------|--------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|-------------|-------------|--------------|--------------|--------------|-------|
| CAGED PLANTS 8. | | | | | | | | | | | | | | | | | | | |
| Flowers opened . . . | .. | 1 | 16 | 46 | 133 | 178 | 188 | 113 | 28 | 3 | 3 | .. | .. | .. | .. | .. | .. | .. | 709 |
| Bolls shed (relative) . . . | .. | .. | 7 | 13 | 53 | 116 | 170 | 78 | 15 | 3 | 3 | .. | .. | .. | .. | .. | .. | .. | 458 |
| Flowers which developed into mature bolls (relative). | .. | 1 | 9 | 33 | 80 | 62 | 18 | 35 | 13 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 251 |
| Percentage of bolls to flowers . | .. | 100 | 56.3 | 71.7 | 60.1 | 34.8 | 9.6 | 31.0 | 46.4 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 35.4 |
| UNUSED PLANTS 19. | | | | | | | | | | | | | | | | | | | |
| Flowers opened . . . | .. | .. | 22 | 22 | 15 | 17 | 85 | 214 | 582 | 757 | 764 | 643 | 378 | 128 | 25 | 3 | .. | .. | 3,635 |
| Bolls shed (relative) . . . | .. | .. | 21 | 19 | 11 | 11 | 37 | 72 | 166 | 336 | 530 | 559 | 344 | 123 | 21 | 3 | .. | .. | 2,253 |
| Causes of boll shedding due to boll-worms. | .. | .. | 20 | 16 | 8 | 10 | 26 | 45 | 98 | 87 | 70 | 61 | 20 | 5 | .. | .. | .. | .. | 465 |
| Causes of boll-shedding due to other causes. | .. | .. | 1 | 3 | 3 | 1 | 11 | 27 | 68 | 249 | 460 | 498 | 324 | 118 | 21 | 3 | .. | .. | 1,787 |
| Flowers which developed into mature bolls (relative). | .. | .. | 1 | 3 | 4 | 6 | 48 | 142 | 396 | 421 | 234 | 84 | 34 | 5 | 4 | .. | .. | .. | 1,382 |
| Percentage of bolls to flowers . | .. | .. | 4.5 | 13.6 | 29.6 | 35.2 | 56.4 | 66.3 | 70.4 | 55.6 | 30.6 | 13.0 | 8.9 | 3.9 | 10.0 | .. | .. | .. | 35.0 |

TABLE XVIII

Flower-opening, relative success from flowers to mature bolls and shedding of bolls with causes of shedding from caged plants and their controls

(1926-27, Surat)

| | 23rd October | 30th October | 6th November | 13th November | 20th November | 27th November | 4th December | 11th December | 18th December | 25th December | 1st January | 8th January | 15th January | 22nd January | 29th January | 5th February | 12th February | 19th February | 26th February | Total |
|---|--------------|--------------|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|-------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|-------|
| CAGED PLANTS 10 | | | | | | | | | | | | | | | | | | | | |
| Flowers opened | .. | 7 | 30 | 60 | 97 | 108 | 133 | 129 | 148 | 106 | 49 | 19 | 4 | . | 1 | . | .. | .. | .. | 891 |
| Bolls shed (relative) | .. | 1 | 6 | 11 | 29 | 32 | 78 | 117 | 129 | 95 | 41 | 14 | 4 | . | 1 | . | .. | .. | .. | 556 |
| Flowers which developed into mature bolls (relative). | .. | 6 | 24 | 49 | 68 | 76 | 55 | 12 | 19 | 11 | 8 | 5 | .. | .. | . | .. | .. | .. | .. | 333 |
| Percentage of bolls to flowers | .. | 85.7 | 80.0 | 81.7 | 70.1 | 70.4 | 41.4 | 9.3 | 12.8 | 10.4 | 16.3 | 26.3 | . | . | . | . | . | . | . | 37.4 |
| UNCAGED PLANTS 20 | | | | | | | | | | | | | | | | | | | | |
| Flowers opened | . | 3 | 7 | 32 | 40 | 40 | 71 | 67 | 88 | 89 | 68 | 96 | 85 | 92 | 114 | 125 | 115 | 40 | 8 | 1,180 |
| Bolls shed (relative) | . | 1 | 4 | 23 | 23 | 29 | 61 | 58 | 77 | 72 | 53 | 59 | 40 | 40 | 44 | 86 | 51 | 80 | 5 | 706 |
| Bolls shed due to boll-worms | . | 1 | 4 | 17 | 21 | 24 | 50 | 50 | 74 | 65 | 48 | 44 | 27 | 16 | 20 | 15 | 3 | .. | .. | 479 |
| Bolls shed due to other causes | .. | .. | .. | 6 | 2 | 5 | 11 | 8 | 3 | 7 | 5 | 15 | 13 | 24 | 24 | 21 | 48 | 80 | 5 | 227 |
| Flowers which developed into mature bolls (relative). | .. | 2 | 3 | 9 | 17 | 11 | 10 | 9 | 11 | 17 | 15 | 37 | 45 | 52 | 70 | 89 | 64 | 10 | 3 | 474 |
| Percentage of bolls to flowers | .. | 66.6 | 42.8 | 28.1 | 42.5 | 27.5 | 14.0 | 13.4 | 12.5 | 19.1 | 22.0 | 38.5 | 35.9 | 56.5 | 61.4 | 71.2 | 55.6 | 25.0 | 37.5 | 40.1 |

TABLE XIX

Flower-opening, relative success from flowers to mature bolls and shedding of bolls with causes of shedding from the caged plants and their controls

(1927-28, Surat)

| | 5th November | 12th November | 19th November | 26th November | 3rd December | 10th December | 17th December | 24th December | 31st December | 7th January | 14th January | 21st January | 28th January | 4th February | 11th February | 18th February | 25th February | TOTAL |
|--|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|-------|
| CAGED PLANTS 20. | | | | | | | | | | | | | | | | | | |
| Flower-opening . . . | . | 23 | 111 | 205 | 304 | 448 | 450 | 523 | 570 | 409 | 202 | 63 | 19 | 6 | 1 | .. | .. | 3,334 |
| Bolls shed (relative) . . | . | 15 | 31 | 22 | 99 | 258 | 310 | 428 | 454 | 361 | 165 | 47 | 10 | 6 | 1 | .. | .. | 2,207 |
| Flowers which developed into mature bolls (relative) | . | 8 | 80 | 183 | 205 | 100 | 140 | 95 | 116 | 48 | 37 | 16 | 9 | .. | .. | .. | .. | 1,127 |
| Percentage of bolls to flowers . | . | 34.8 | 72.1 | 89.2 | 67.4 | 42.4 | 31.1 | 18.2 | 20.4 | 11.7 | 18.3 | 25.4 | 47.4 | . | .. | .. | .. | 33.8 |
| UN-CAGED PLANTS 16 | | | | | | | | | | | | | | | | | | |
| Flower-opening . . . | . | 1 | 6 | 12 | 31 | 67 | 67 | 116 | 142 | 108 | 279 | 223 | 221 | 109 | 46 | 22 | 2 | 1,542 |
| Bolls shedding (relative) . | . | 1 | 4 | 4 | 19 | 36 | 40 | 50 | 60 | 91 | 135 | 138 | 188 | 108 | 46 | 22 | 2 | 950 |
| Bolls shed due to boll-worms . | . | 1 | 4 | 3 | 16 | 23 | 23 | 11 | 25 | 24 | 24 | 10 | 6 | 2 | . | . | . | 192 |
| Bolls shed due to other causes . | .. | .. | . | 1 | 3 | 13 | 17 | 19 | 41 | 67 | 111 | 128 | 182 | 106 | 46 | 22 | 2 | 758 |
| Flowers which developed into mature bolls (relative) | .. | .. | 2 | 8 | 12 | 31 | 27 | 66 | 76 | 107 | 144 | 85 | 33 | 1 | . | .. | .. | 502 |
| Percentage of bolls to flowers . | .. | .. | 33 | 67 | 39 | 46 | 40 | 57 | 53 | 54 | 52 | 38 | 15 | 1 | . | .. | .. | 38.4 |

TABLE XX

Dates of picking of kapas from caged and uncaged plants and the number of bolls picked each time

| Pickings | 1928-29 | | | | | | 1929-30 | | | | 1930-31 | | | |
|-------------|-------------------------------|---------------------|------------------------|---------------------|------------------------|---------------------|------------------------|---------------------|------------------------|---------------------|------------------------|---------------------|-----------------|---------------------|
| | CAGED PLANTS (99) | | UNCAGED PLANTS (104) | | CAGED PLANTS (118) | | UNCAGED PLANTS (112) | | CAGED PLANTS (118) | | UNCAGED PLANTS (116) | | | |
| | Date of picking | No. of bolls picked | Date of picking | No. of bolls picked | Date of picking | No. of bolls picked | Date of picking | No. of bolls picked | Date of picking | No. of bolls picked | Date of picking | No. of bolls picked | Date of picking | No. of bolls picked |
| 1st picking | 7th to 9th January. | 683 | 8th to 10th February. | 1,055 | 1st to 2nd January. | 1,023 | 22nd to 24th February. | 1,809 | 5th to 8th January. | 148 | 20th to 28th February. | 5,563 | | |
| 2nd picking | 18th to 22nd January. | 1,943 | 22nd to 26th February. | 3,122 | 28th to 29th January. | 40 | 11th to 13th March. | 4,182 | 14th to 24th January. | 4,497 | 12th to 14th March. | 2,180 | | |
| 3rd picking | 31st January to 5th February. | 3,213 | 5th to 7th March. | 1,870 | 10th to 11th February. | 7,976 | 19th March. | 272 | 14th to 19th February. | 3,107 | 19th March. | 14 | | |
| 4th picking | 18th to 22nd February. | 4,078 | 19th to 20th March. | 561 | .. | .. | 24th March. | 29 | 10th March. | 771 | .. | .. | | |
| 5th picking | 28th to 28th February. | 1,192 | 22nd March. | 208 | .. | .. | .. | .. | .. | .. | .. | .. | | |
| 6th picking | 2nd to 7th March. | 110 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | | |
| TOTAL | .. | 11,169 | .. | 6,816 | .. | 9,039 | .. | 6,292 | .. | 8,623 | .. | 8,087 | | |

TABLE XXI

Classification of the locks collected per plant from the caged and uncaged plants during 3 years at Surat

| | No. of plants observed | Total locks | Sound locks | DAMAGED LOCKS. | | USELESS LOCKS. | |
|----------------|------------------------|-------------|-------------|-------------------|---------------------|-------------------|---------------------|
| | | | | Due to boll-worms | Due to other causes | Due to boll-worms | Due to other causes |
| 1928-29. | | | | | | | |
| Caged plants | 99 | 350 | 307 | 13 | 24 | 3 | 3 |
| Control plants | 104 | 202 | 144 | 38 | 9 | 7 | 4 |
| 1929-30. | | | | | | | |
| Caged plants | 113 | 244 | 236 | 4 | 2 | 2 | .. |
| Control plants | 112 | 171 | 156 | 9 | 2 | 3 | 1 |
| 1930-31. | | | | | | | |
| Caged plants | 118 | 223 | 206 | 14 | 2 | 1 | .. |
| Control plants | 116 | 212 | 191 | 18 | 1 | 2 | .. |

TABLE XXII
Population of Earias larvae in pods of 25 bhendi plants plucked as they became ripe on different varieties at Surat (plot un-manured)
 (1930-31)

| Date | SURAT 5 | | | SURAT 9 | | | KHANDESH LATE | | |
|--------------------|--------------------|----------------|------------------|--------------------|----------------|------------------|--------------------|----------------|------------------|
| | Weight in Tolas | No. of pods | Earias Larvae | Weight in Tolas | No. of pods | Earias Larvae | Weight in Tolas | No. of pods | Earias Larvae |
| 1st September 1930 | .. | .. | .. | 11 | 12 | 9 | .. | .. | .. |
| 5th September 1930 | .. | .. | .. | 3 | 8 | 2 | .. | .. | .. |
| 11th October 1930 | .. | .. | .. | .. | .. | .. | 7 | 4 | .. |
| 23rd October 1930 | .. | .. | .. | 10 | 16 | 31 | 35 | 26 | 12 |
| 28th October 1930 | 8 | 12 | 25 | 4 | 9 | 17 | 16 | 22 | 7 |
| 2nd November 1930 | 1 | 3 | 3 | 6 | 12 | 15 | 44 | 34 | 13 |
| 6th November 1930 | .. | .. | .. | 1 | 2 | .. | .. | .. | .. |
| 13th November 1930 | 3 | 2 | 2 | 2 | 5 | 3 | 22 | 24 | 8 |
| 22nd November 1930 | .. | .. | .. | 1 | 3 | 4 | 38 | 26 | 16 |
| 6th December 1930 | 11 | 5 | 1 | 8 | 14 | 4 | 17 | 17 | 2 |
| 13th December 1930 | 5 | 10 | 2 | 4 | 12 | 3 | 4 | 8 | 3 |
| 20th December 1930 | 10 | 14 | 7 | 2 | 5 | .. | 14 | 25 | 6 |
| 31st December 1930 | 8 | 20 | .. | 4 | 10 | 4 | 17 | 46 | 4 |
| 10th January 1931 | 26 | 50 | 5 | 10 | 37 | 5 | 36 | 97 | 4 |
| 22nd January 1931 | 16 | 24 | 7 | 17 | 45 | 5 | 26 | 46 | 5 |

TABLE XXIII

Population of Earias larvae in pods of 25 bhendi plants plucked as they became fully grown on different varieties at Surat (plot manured with castor-oil)

(1930-31)

| Date | SURAT 5 | | | SURAT 9 | | | AMERICAN LONG GREEN | | | AMERICAN DWARF GREEN | | | AMERICAN WHITE VELVET | | | KHANDER LATE | | | REMARKS |
|---------------------|---------|------|---------------|---------|------|---------------|---------------------|------|---------------|----------------------|------|---------------|-----------------------|------|---------------|--------------|------|---------------|---------|
| | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | |
| 23rd August 1930 | .. | .. | .. | 7 | 9 | 1 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 24th August 1930 | .. | .. | .. | 16 | 20 | 8 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 1st September 1930 | .. | .. | .. | 30 | 25 | 23 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 5th September 1930 | 8 | 6 | 3 | 21 | 20 | 11 | .. | .. | .. | 2 | 2 | .. | 1 | 1 | .. | .. | .. | .. | |
| 18th September 1930 | 25 | 10 | 12 | 27 | 24 | 21 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 4th October 1930 | 8 | 8 | 2 | .. | .. | 6 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 42 | 25 | .. | 4 |
| 11th October 1930 | .. | .. | .. | 3 | 9 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | 83 | 45 | 23 | |
| 23rd October 1930 | 54 | 50 | 94 | 22 | 35 | 50 | .. | .. | .. | 3 | 3 | 8 | .. | .. | .. | 17 | 14 | 9 | |
| 28th October 1930 | 24 | 35 | 51 | 9 | 16 | 23 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 54 | 38 | 18 | |
| 3rd November 1930 | 38 | 42 | 26 | 8 | 15 | 16 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 6 | 6 | 5 | |
| 6th November 1930 | 4 | 7 | 8 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | 17 | 13 | 4 | |
| 13th November 1930 | 16 | 11 | 12 | 3 | 4 | 1 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 42 | 31 | 10 | |
| 22nd November 1930 | 17 | 23 | 14 | 5 | 16 | 9 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 40 | 34 | 4 | |
| 6th December 1930 | 27 | 36 | 8 | 6 | 16 | 3 | 16 | 15 | 5 | 37 | 38 | 7 | 19 | 22 | 5 | 6 | 14 | 2 | |
| 13th December 1930 | 17 | 35 | 4 | 6 | 17 | 8 | 2 | 4 | 1 | 6 | 15 | 3 | 4 | 9 | 3 | 25 | 45 | 6 | |
| 20th December 1930 | 31 | 51 | 10 | 5 | 17 | 12 | 10 | 13 | 4 | 18 | 27 | 3 | 13 | 17 | 8 | 80 | 131 | 9 | |
| 31st December 1930 | 40 | 79 | 6 | 5 | 19 | 3 | 8 | 18 | 1 | 18 | 38 | 6 | 16 | 29 | 5 | 72 | 119 | 9 | |
| 10th January 1931 | 50 | 101 | 6 | 7 | 24 | 3 | 18 | 38 | 8 | 33 | 62 | 3 | 7 | 17 | 8 | 80 | 131 | 9 | |
| 22nd January 1931 | 60 | 98 | 18 | 22 | 64 | 9 | 28 | 44 | 6 | 32 | 63 | 10 | 29 | 55 | 8 | 72 | 119 | 9 | |

TABLE XXV

Population of Earias larvae on 25 plants of bhendi of different varieties in Khandesh, dibbled with the general crop of cotton on 20th June 1930

(1930-31, Jalgaon)

| Date | Nasik local | | | Katal local | | | Surat 5 | | | Surat 9 | | | American long green | | | American dwarf green | | | American white velvet | | | REMARKS |
|---------------------|-------------|------|---------------|-------------|------|---------------|---------|------|---------------|---------|------|---------------|---------------------|------|---------------|----------------------|------|---------------|-----------------------|------|---------------|---|
| | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | |
| 12th August 1930 | 74 | 40 | . | . | . | . | 46 | 27 | . | 46 | 27 | . | . | . | . | 7 | 3 | . | . | . | . | Excellent growth of plants throughout the season over 6 feet and the rest about 4 feet and over |
| 20th August 1930 | 106 | 44 | . | . | . | . | 80 | 38 | . | 80 | 38 | . | . | . | . | 10 | 5 | . | . | . | . | |
| 28th August 1930 | 125 | 48 | . | . | . | . | 130 | 46 | 1 | 130 | 46 | 1 | . | . | . | 16 | 8 | . | . | . | . | |
| 30th August 1930 | 57 | 52 | 1 | . | . | . | 30 | 49 | . | 30 | 49 | . | . | . | . | 16 | 8 | . | . | . | . | |
| 3rd September 1930 | 66 | 52 | 6 | 38 | 52 | 32 | 100 | 85 | 14 | 100 | 85 | 14 | . | . | . | 3 | 2 | 1 | 3 | 6 | 1 | |
| 13th September 1930 | 38 | 30 | 1 | 38 | 37 | 3 | 72 | 47 | 4 | 72 | 47 | 4 | . | . | . | 45 | 19 | 1 | 44 | 20 | . | |
| 22nd September 1930 | 12 | 9 | 2 | 46 | 31 | 3 | 28 | 10 | 1 | 3 | 2 | 2 | 6 | 1 | 16 | 7 | 1 | 21 | 8 | . | . | |
| 28th September 1930 | 12 | 18 | 3 | 83 | 31 | 3 | 27 | 7 | 3 | 11 | 11 | 2 | 9 | 1 | 8 | 8 | . | 15 | 6 | . | . | |
| 30th September 1930 | 28 | 32 | 15 | 127 | 48 | 6 | 3 | 1 | 30 | 30 | 5 | 18 | 7 | 1 | 12 | 8 | 2 | 10 | 5 | 1 | . | |
| 7th October 1930 | 64 | 81 | 53 | 210 | 86 | 0 | 34 | 18 | 5 | 88 | 68 | 32 | 77 | 37 | 5 | 50 | 17 | 4 | 36 | 20 | 2 | |
| 14th October 1930 | 14 | 21 | 34 | 44 | 34 | 5 | 39 | 25 | 11 | 71 | 15 | 5 | 63 | 37 | 6 | 43 | 20 | 5 | 27 | 14 | 1 | |
| 18th October 1930 | 12 | 15 | 11 | 107 | 80 | 20 | 80 | 56 | 32 | 7 | 10 | 5 | 44 | 22 | 7 | 43 | 21 | 11 | 66 | 37 | 7 | |
| 23rd October 1930 | 7 | 9 | 10 | 47 | 31 | 4 | 22 | 12 | 13 | 35 | 15 | 8 | 35 | 20 | 13 | 35 | 20 | 13 | 99 | 42 | 14 | |
| 26th October 1930 | 13 | 25 | 60 | 75 | 41 | 40 | 10 | 19 | 5 | 18 | 35 | 16 | 49 | 41 | 21 | 41 | 21 | 41 | 40 | 24 | 42 | |
| 13th November 1930 | . | . | . | 77 | 58 | 23 | 5 | 6 | 10 | 43 | 33 | 41 | 33 | 21 | 17 | 42 | 29 | 22 | 42 | 29 | 22 | |
| 25th November 1930 | . | . | . | 48 | 38 | 21 | 19 | 17 | 25 | 1 | 45 | 31 | 41 | 72 | 54 | 78 | 47 | 41 | 43 | 41 | 43 | |
| 1st December 1930 | . | . | . | 31 | 12 | 11 | 6 | 9 | 13 | . | 39 | 34 | 47 | 77 | 84 | 82 | 33 | 31 | 33 | 31 | 41 | |
| 7th December 1930 | . | . | . | 11 | 9 | 6 | 11 | 16 | 17 | . | 26 | 24 | 52 | 51 | 46 | 47 | 31 | 29 | 17 | 17 | 17 | |
| 17th December 1930 | . | . | . | 11 | 9 | 12 | 11 | 16 | 17 | . | 19 | 19 | 21 | 19 | 17 | 33 | 21 | 16 | 32 | 21 | 16 | |
| 25th December 1930 | . | . | . | 11 | 9 | 12 | 11 | 16 | 17 | . | 30 | 18 | 43 | 38 | 26 | 42 | 21 | 17 | 32 | 21 | 17 | |
| 6th January 1931 | . | . | . | 18 | 19 | 17 | 6 | 14 | 8 | . | 14 | 17 | 28 | 12 | 15 | 23 | 19 | 15 | 9 | 15 | 9 | |
| 15th January 1931 | . | . | . | 11 | 13 | 11 | . | 7 | 5 | . | 16 | 18 | 27 | 18 | 23 | 43 | . | . | . | . | . | |

TABLE XXVI
Population of Earias larvae on bhendi plants of different varieties, dibbled one month after the general crop (i.e., on 19th July 1930 at Jalgaon)
 (1930-31)

| Date | Nasik local | | | Katal local | | | Surat 5 | | | Surat 9 | | | American long green | | | American dwarf green | | | American white velvet | | | REMARKS |
|----------------------------------|-------------|------|---------------|-------------|------|---------------|---------|------|---------------|----------------|------|---------------|---------------------|------|---------------|----------------------|------|---------------|-----------------------|------|---------------|--|
| | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | Tolas | Pods | Earias larvae | |
| No. of plants observed each time | 25 plants | | | 4 plants. | | | 1 plant | | | Failed to grow | | | 3 plants. | | | 5 plants | | | 8 plants. | | | Poor growth of plants; most of them did not grow at all. |
| 16th September 1930 | 13 | 15 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 22nd September 1930 | 5 | 10 | 2 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 30th September 1930 | 11 | 10 | 1 | .. | .. | .. | 5 | 3 | 2 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 7th October 1930 | 18 | 13 | 11 | 6 | 3 | .. | 5 | 2 | .. | 2 | 1 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 13th October 1930 | 12 | 15 | 2 | .. | .. | .. | 1 | 1 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | |
| 20th October 1930 | 9 | 11 | 5 | 11 | 5 | 1 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 4 | 2 | .. | .. | .. | .. | |
| 26th October 1930 | 1 | 1 | 1 | 22 | 14 | 5 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | 1 | .. | 2 | 1 | 1 | |
| 12th November 1930 | 5 | 12 | 14 | 6 | 5 | 13 | 5 | 6 | 5 | .. | .. | .. | .. | .. | .. | 7 | 5 | 4 | 3 | 1 | 2 | |
| 22nd November 1930 | .. | .. | .. | 9 | 4 | 3 | 4 | 3 | 2 | .. | .. | .. | .. | .. | .. | 1 | 1 | 2 | 4 | 3 | 2 | |
| 5th December 1930 | 2 | 6 | 6 | 10 | 11 | 5 | 3 | 1 | 1 | .. | .. | .. | .. | .. | .. | 11 | 6 | 7 | 2 | 2 | 3 | |
| 10th December 1930 | 1 | 3 | 3 | 9 | 7 | 8 | 1 | 1 | .. | .. | .. | .. | .. | .. | .. | 5 | 3 | 2 | 5 | 2 | 5 | |
| 16th December 1930 | .. | .. | .. | .. | 4 | 3 | 5 | 2 | 5 | .. | .. | .. | .. | .. | .. | 11 | 7 | 7 | 12 | 8 | 6 | |
| 26th December 1930 | .. | .. | .. | 10 | 6 | 17 | 3 | 2 | 5 | .. | .. | .. | 2 | 1 | 2 | 5 | 3 | 5 | 8 | 5 | 9 | |
| 6th January 1931 | .. | .. | .. | 6 | 3 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | 12 | 17 | |
| 16th January 1931 | .. | .. | .. | 11 | 15 | 2 | 3 | 2 | 1 | .. | .. | .. | 6 | 3 | 8 | 6 | 5 | 8 | 12 | 8 | 5 | |

TABLE XXVIII

Boll-worm moths attracted to baits exposed in the cotton fields at Surat

(1927-28)

| Date | Cotton seed | Bhendi seed | Sesamum cake |
|----------------------------|-------------|------------------|--------------|
| 20th December 1927 | Started * | .. | .. |
| 23rd December 1927 | 8 | .. | .. |
| 24th December 1927 | 8 | .. | .. |
| 25th December 1927 | 1 | .. | .. |
| 30th December 1927 | ..* | .. | .. |
| 1st January 1928 | 1 | .. | .. |
| 2nd January 1928 | 17 | .. | .. |
| 6th January 1928 | 1* | Started * | .. |
| 7th January 1928 | .. | .. | .. |
| 8th January 1928 | 3 | 1 | .. |
| 9th January 1928 | 5 | .. | .. |
| 10th January 1928 | ..* | 1 ¹ | Started * |
| 14th January 1928 | ..* | .. ¹¹ | .. |
| 17th January 1928 | 1 | 2* | ..* |
| 19th January 1928 | ..* | ..* | ..* |
| 23rd January 1928 | ..* | 1* | 1* |
| 24th January 1928 | .. | .. | 1 |
| 25th January 1928 | ..* | ..* | ..* |
| 29th January 1928 | .. | 4 | 11 |
| 30th January 1928 | 1 | 1 | 1 |
| 31st January 1928 | 1 | 3 | 38 |
| 1st February 1928 | 1* | 1* | 2* |
| 2nd February 1928 | 3 | 2 | 8 |
| 4th February 1928 | .. | .. | 2 |
| 6th February 1928 | 1* | 1* | 1* |
| 8th February 1928 | .. | 2 | 5 |
| 9th February 1928 | .. | 6 | 3 |
| 11th February 1928 | .. | .. | 1 |
| 13th February 1928 | ..* | ..* | 1* |
| 14th February 1928 | 1 | 2 | 4 |
| 16th February 1928 | 2 | 1 | .. |
| 17th February 1928 | 1 | .. | .. |

* Indicates the days on which the baits were moistened with water.

TABLE XXIX

Boll-worm moths attracted to the bait pans at Surat
(1929-30)

| Week ending | Sesamum cake | | | Cotton seed cake | | | Groundnut cake | | |
|--------------------------|--------------|---------------------|-------------------------|------------------|---------------------|-------------------------|----------------|---------------------|-------------------------|
| | No. of pans | <i>Earias</i> moths | Moths of Pink boll-worm | No. of pans | <i>Earias</i> moths | Moths of Pink boll-worm | No. of pans | <i>Earias</i> moths | Moths of Pink boll-worm |
| 10th October 1929 . . . | 2 | 2 | .. | 4 | 2 | .. | .. | .. | .. |
| 26th October 1929 . . . | 3 | 1 | .. | 4 | 1 | .. | .. | .. | .. |
| 2nd November 1929 . . . | 3 | 1 | .. | 4 | . | 1 | .. | .. | .. |
| 9th November 1929 . . . | 3 | 6 | 6 | 4 | . | 28 | .. | .. | .. |
| 16th November 1929 . . . | 4 | 8 | 18 | 8 | 20 | 34 | .. | .. | .. |
| 23rd November 1929 . . . | 4 | 33 | 22 | 8 | 43 | 70 | .. | .. | .. |
| 30th November 1929 . . . | 4 | 95 | 9 | 8 | 56 | 39 | 2 | 21 | 6 |
| 7th December 1929 . . . | 4 | 16 | 10 | 8 | 52 | 25 | 2 | 15 | 1 |
| 14th December 1929 . . . | 4 | 23 | 10 | 8 | 23 | 19 | 2 | 3 | 4 |
| 21st December 1929 . . . | 4 | 50 | 7 | 8 | 46 | 13 | 2 | 36 | 7 |
| 28th December 1929 . . . | 4 | 53 | 7 | 8 | 26 | 14 | 2 | 2 | 1 |
| 4th January 1930 . . . | 4 | 14 | 11 | 8 | 14 | 23 | 2 | 5 | 1 |
| 11th January 1930 . . . | 4 | 43 | 25 | 8 | 67 | 97 | 2 | 7 | 22 |
| 18th January 1930 . . . | 4 | 1 | 4 | 8 | 6 | 10 | 2 | 2 | .. |
| 25th January 1930 . . . | 4 | 196 | 145 | 8 | 73 | 208 | 2 | 69 | 84 |
| 1st February 1930 . . . | 4 | 90 | 79 | 8 | 19 | 114 | 2 | 47 | 38 |
| 8th February 1930 . . . | 4 | 310 | 102 | 8 | 108 | 242 | 2 | 210 | 61 |
| 15th February 1930 . . . | 5 | 344 | 255 | 9 | 201 | 445 | 3 | 559 | 164 |
| 22nd February 1930 . . . | 5 | 189 | 407 | 9 | 96 | 620 | 3 | 63 | 213 |
| 1st March 1930 . . . | 5 | 14 | 23 | 9 | 7 | 75 | 3 | 21 | 15 |

TABLE XXX
Boll-worm moths attracted at the bait pans at Surat
(1930-31)

| Week ending. | Groundnut cake (2 pans) | | Sesamum cake (1 pan) | | Cotton seed-cake (1 pan) | | Mixture of groundnut, cotton seed-cakes (2 pans) | | Sandal Wood paste (1 pan) | | Nutmeg paste (1 pan) | | Mixture of sesamum and cotton seed-cakes (2 pans) | |
|--------------------|-------------------------|-------------------------|----------------------|-------------------------|--------------------------|-------------------------|--|-------------------------|---------------------------|-------------------------|----------------------|-------------------------|---|-------------------------|
| | Earias moths | Moths of Pink boll-worm | Earias moths | Moths of Pink boll-worm | Earias moths | Moths of Pink boll-worm | Earias moths | Moths of Pink boll-worm | Earias moths | Moths of Pink boll-worm | Earias moths | Moths of Pink boll-worm | Earias moths | Moths of Pink boll-worm |
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 |
| 11th October 1930 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 18th October 1930 | 3 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 25th October 1930 | 1 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 1st November 1930 | 7 | .. | 6 | .. | 1 | .. | .. | .. | 6 | .. | 16 | .. | .. | .. |
| 8th November 1930 | 65 | .. | 7 | .. | 2 | .. | .. | .. | 10 | .. | 8 | .. | .. | .. |
| 15th November 1930 | 4 | 17 | 3 | 5 | 1 | 13 | 9 | 13 | 3 | 1 | 6 | 5 | .. | .. |
| 22nd November 1930 | 10 | 65 | 5 | 14 | 3 | 42 | .. | 38 | 2 | .. | 6 | 5 | .. | .. |
| 29th November 1930 | 20 | 45 | 10 | 8 | 14 | 14 | 19 | 36 | 6 | 1 | 4 | .. | .. | .. |
| 6th December 1930 | 14 | 34 | 15 | 27 | 1 | 10 | 7 | 27 | 1 | 1 | 1 | .. | .. | .. |
| 13th December 1930 | 12 | 7 | 11 | 1 | .. | .. | 1 | 4 | 1 | 1 | 1 | .. | .. | .. |
| 20th December 1930 | 8 | 6 | 1 | 2 | .. | 2 | .. | 4 | * | .. | .. | .. | .. | .. |
| 27th December 1930 | 21 | 37 | 17 | 6 | 6 | 11 | 20 | 22 | .. | .. | .. | .. | .. | .. |
| 3rd January 1931 | 15 | 22 | 21 | 13 | .. | 15 | 8 | 35 | .. | .. | .. | .. | .. | .. |
| 10th January 1931 | 2 | 14 | .. | 4 | .. | 3 | 1 | 13 | .. | .. | .. | .. | .. | .. |
| 17th January 1931 | .. | 2 | .. | .. | .. | .. | .. | 2 | .. | .. | .. | .. | .. | .. |
| 24th January 1931 | 5 | 36 | 8 | 34 | 8 | 24 | 5 | 34 | .. | .. | .. | .. | .. | .. |
| 31st January 1931 | 8 | 74 | 3 | 17 | 1 | 42 | 21 | 77 | .. | .. | .. | .. | .. | .. |
| 7th February 1931 | 31 | 199 | 46 | 195 | .. | .. | .. | .. | .. | .. | .. | .. | 43 | 169 |
| 14th February 1931 | 18 | 162 | 31 | 151 | .. | .. | .. | .. | .. | .. | .. | .. | 23 | 163 |
| 21st February 1931 | 12 | 67 | 41 | 89 | .. | .. | .. | .. | .. | .. | .. | .. | 19 | 71 |
| 28th February 1931 | 2 | 8 | 4 | 21 | .. | .. | .. | .. | .. | .. | .. | .. | 8 | 28 |

* Stopped.

TABLE XXXI

Boll-worm moths attracted at the bait pans at Jalgaon

(1929-30)

| Fortnight ending | Sesamum cake (2 pans) | Cotton-seed cake (2 pans) | 4 pans (2 of sesamum cake and 2 of cotton- seed cake) |
|-------------------------------|--------------------------|------------------------------|--|
| | (<i>Earias</i> moths) | (<i>Earias</i> moths) | (Pink Boll- worm moths) |
| 14th September 1929 | 3 | .. | 4 |
| 28th September 1929 | 11 | 5 | 3 |
| 12th October 1929 | 75 | 43 | 126 |
| 26th October 1929 | 147 | 108 | 712 |
| 9th November 1929 | 128 | 124 | 460 |
| 23rd November 1929 | 201 | 346 | 1,347 |
| 7th December 1929 | 401 | 484 | 835 |
| 21st December 1929 | 109 | 49 | 269 |
| 4th January 1930 | 35 | 22 | 91 |
| 18th January 1930 | 13 | 17 | 55 |
| 1st February 1930 | 4 | 3 | 47 |
| 15th February 1930 | .. | 1 | 5 |
| 1st March 1930 | 1 | 1 | 75 |
| 15th March 1930 | .. | .. | 25 |
| 29th March 1930 | .. | .. | 4 |

TABLE XXXII

Boll-worm moths attracted to the bait pans at Jalgaon

(1930-31)

| Week ending | COTTON SEED CAKE | | GROUNDNUT CAKE FREELY DILUTED IN WATER OF THE WHOLE PAN | | COTTON SEED CAKE | | GROUNDNUT CAKE MOIS- TENED IN DISH | |
|--------------------|------------------|-----------|--|-----------|------------------|-----------|---------------------------------------|-----------|
| | Earias | Platyedra | Earias | Platyedra | Earias | Platyedra | Earias | Platyedra |
| 11th October 1930 | 2 | .. | 1 | .. | .. | .. | 1 | 1 |
| 18th October 1930 | 2 | .. | 3 | 1 | .. | .. | 11 | 5 |
| 25th October 1930 | 4 | 7 | 7 | 2 | 4 | 3 | 20 | 10 |
| 1st November 1930 | 2 | 5 | 2 | 32 | 12 | 26 | 11 | 23 |
| 8th November 1930 | 3 | 11 | 11 | 24 | 12 | 24 | 5 | 17 |
| 15th November 1930 | 5 | 19 | 11 | 117 | 11 | 51 | 15 | 41 |
| 22nd November 1930 | 6 | 47 | 31 | 367 | 9 | 157 | 16 | 128 |
| 29th November 1930 | 32 | 166 | 119 | 715 | 12 | 215 | 80 | 338 |
| 6th December 1930 | 20 | 245 | 197 | 1,161 | 66 | 993 | 190 | 210 |
| 13th December 1930 | 78 | 237 | 659 | 982 | 294 | 992 | 116 | 143 |
| 20th December 1930 | 21 | 206 | 191 | 390 | 181 | 530 | 64 | 127 |
| 27th December 1930 | 29 | 251 | 216 | 272 | 217 | 452 | 78 | 235 |
| 3rd January 1931 | 74 | 708 | 113 | 184 | 152 | 332 | 415 | 846 |
| 10th January 1931 | 49 | 770 | 152 | 129 | 299 | 313 | 590 | 843 |
| 15th January 1931 | 73 | 426 | 98 | 20 | 206 | 141 | 235 | 250 |

TABLE XXXIII

Weekly flower-opening, percentage of the bolls shed due to boll-worms out of the total bolls shed, number of flowers which developed into mature bolls and the percentage of the weekly flowers which succeeded into mature bolls in the control plot and in the plots dusted with calcium arsenate alone and calcium arsenate with lime.

(1926-27)

(Observations of 20 plants in each case)

| Weeks | CONTROL PLOT | | | | DUSTED WITH CALCIUM ARSENATE | | | | DUSTED WITH CALCIUM ARSENATE AND LIME 1:3 | | | |
|--------------------|----------------|--|--|---|------------------------------|---|--|---|---|--|--|---|
| | Flowers opened | Percent-age of bolls shed due to boll-worm out of total shed | No. of flowers developed into mature bolls | Percent-age of success of flowers into mature bolls | Flowers opened | Percent-age of bolls shed due to boll-worms out of total shed | No. of flowers developed into mature bolls | Percent-age of success of flowers into mature bolls | Flowers opened | Percent-age of bolls shed due to boll-worm out of total shed | No. of flowers developed into mature bolls | Percent-age of success of flowers into mature bolls |
| 4th December 1926 | 71 | 70.4 | 10 | 14.0 | 7 | 12.8 | 5 | 71.4 | 7 | 12.8 | 4 | 57.1 |
| 11th December 1926 | 67 | 74.6 | 9 | 13.4 | 44 | 20.4 | 28 | 63.6 | 41 | 46.3 | 16 | 39.1 |
| 18th December 1926 | 88 | 84.0 | 11 | 12.5 | 72 | 13.8 | 51 | 70.8 | 49 | 32.6 | 17 | 34.6 |
| 25th December 1926 | 89 | 73.0 | 17 | 19.1 | 193 | 15.0 | 118 | 61.5 | 107 | 32.7 | 49 | 45.7 |
| 1st January 1927 | 68 | 70.5 | 15 | 22.0 | 229 | 11.3 | 134 | 58.5 | 154 | 35.6 | 73 | 47.4 |
| 8th January 1927 | 96 | 45.8 | 37 | 38.5 | 296 | 8.4 | 98 | 31.0 | 211 | 26.5 | 112 | 53.4 |
| 15th January 1927 | 85 | 31.7 | 45 | 52.9 | 265 | 9.4 | 15 | 5.6 | 198 | 19.9 | 101 | 51.0 |
| 22nd January 1927 | 92 | 17.3 | 52 | 56.5 | 293 | 4.0 | 9 | 3.0 | 237 | 11.3 | 66 | 27.9 |
| 29th January 1927 | 114 | 17.5 | 70 | 61.4 | 209 | 1.0 | 6 | 2.8 | 196 | 7.3 | 31 | 15.8 |
| 5th February 1927 | 125 | 12.0 | 89 | 71.2 | 102 | 3.9 | 8 | 7.8 | 189 | 6.9 | 19 | 10.0 |
| 12th February 1927 | 115 | 2.6 | 64 | 55.6 | 33 | .. | 8 | 24.2 | 134 | 4.4 | 8 | 5.9 |

TABLE XXXIV

Weekly flower-opening and relative number of flowers which developed into mature bolls from the dusted and control plots

(1927-28)

| Treatments | 12th November | 19th November | 26th November | 3rd December | 10th December | 17th December | 24th December | 31st December | 7th January | 14th January | 21st January | 28th January | 4th February | 11th February | 18th February |
|---|---|---------------|---------------|--------------|---------------|---------------|---------------|---------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|
| (1) Dusting calcium arsenate at 9 days interval. | Flowers opened Bolls matured out of above flowers. | 17 10 | 62 37 | 177 123 | 344 195 | 287 60 | 408 28 | 397 5 | 207 1 | 77 .. | 6 .. | | | | |
| (2) Dusting as above and spraying fish oil rosin soap. | Flowers opened Bolls matured out of the above flowers. | | 16 10 | 50 35 | 103 64 | 283* 170 | 270 100 | 309 48 | 219 1 | 135 1 | 88 4 | 3 .. | | | |
| (3) Dusting calcium arsenate with gram flour. | Flowers opened Bolls matured out of the above flowers. | 1 .. | 44 20 | 131 62 | 290 138 | 194 80 | 251 70 | 246 31 | 258 12 | 115 11 | 42 3 | 7 1 | | | |
| (4) Dusting calcium arsenate twice at bud development and twice at flower-opening time. | Flowers opened Bolls matured out of the above flowers. | 3 1 | 81 14 | 89 53 | 190 115 | 137 58 | 196 42 | 236 27 | 192 8 | 117 1 | 43 2 | 9 .. | | | |
| (5) Dusting calcium arsenate twice at flowering time. | Flowers opened Bolls matured out of the above flowers. | | | 13 13 | 84 56 | 93 73 | 147 113 | 289 135 | * 335 36 | 421 20 | 286 11 | 97 8 | 11 .. | 2 1 | |
| (6) Untreated plot. | Flowers opened Bolls matured out of the above flowers. | | 14 4 | 37 13 | 61 22 | 108 45 | 77 37 | 112 54 | 159 76 | 165 63 | 137 68 | 116 47 | 55 10 | 22 2 | 8 1 |

* Spraying started.

TABLE XXXV

Analysis of calcium arsenate (through the courtesy of the Agricultural Chemist, College of Agriculture, Poona)

| Percentage of | Berger's calcium arsenate (from England) | Calcium arsenate from Grasselli & Co. (from America) |
|---|--|--|
| Moisture | 1.70 | traces. |
| Sand | 0.25 | 0.20 |
| Soluble arsenic oxide (As_2O_3) | 1.26 | 0.22 |
| Total arsenic oxide (As_2O_3) | 45.43 | 40.11 |

TABLE XXXVI

Dates of dusting in the different plots in the year 1928-29

| Plot No. | Treatment given | Date of 1st dusting | Date of 2nd dusting | Date of 3rd dusting | Date of 4th dusting |
|----------|--|---------------------|---------------------|---------------------|---------------------|
| 1 | Dusting with blackened calcium arsenate when the boll-worms are feeding in shoots. | 8th October. | 18th October. | 29th October. | .. |
| 2 | Dusting with blackened calcium arsenate when the flower-bud development is vigorous. | 6th November. | 15th November. | 24th November. | 3rd December. |
| 3 | Dusting with blackened calcium arsenate when the boll-development has commenced. | 19th November. | 28th November. | 7th December. | 17th December. |
| 4 | Dusting with white calcium arsenate alone when vigorous flower-bud development is in progress. | 5th November. | 16th November. | 25th November. | 3rd December. |
| 5 | Dusting with sodium silico-fluoride alone when the flower-bud development is in progress. | 7th November. | 16th November. | 26th November. | 5th December. |
| 6 | Dusting with blackened sodium silico-fluoride when flower-bud development is in progress. | 8th November. | 17th November. | 27th November. | 7th December. |

TABLE XXXVII

Weekly flower-opening and success of flowers to mature bolls from 20 plants in each case from plots dusted with insecticides and their controls

(1928-29)

| Plot No | Treatment | 3rd November | 10th November | 17th November | 24th November | 1st December | 8th December | 15th December | 22nd December | 29th December | 5th January | 15th January | 22nd January | 29th January | 2nd February |
|---------|--|--|---------------|---------------|---------------|--------------|--------------|---------------|---------------|---------------|-------------|--------------|--------------|--------------|--------------|
| 2 | Blackened calcium arsenate at bud development. | Flowers opened Bolls matured out of the above flowers | .. | .. | 79 51 | 156 85 | 240 99 | 322 125 | 477 142 | 379 59 | 193 4 | 73 | 21 | 1 | .. |
| 2-A | Control for No. 2 | Flowers opened Bolls matured out of the above flowers | .. | .. | 35 8 | 77 16 | 98 26 | 171 56 | 296 130 | 281 142 | 225 83 | 194 34 | 167 26 | 89 5 | 48 4 |
| 4 | White calcium arsenate at bud development. | Flowers opened Bolls matured out of the above flowers | .. | .. | 106 58 | 212 121 | 294 161 | 484 214 | 644 225 | 446 51 | 269 11 | 116 2 | 35 | .. | .. |
| 4-A | Control for No. 4 | Flowers opened Bolls matured out of the above flowers | .. | .. | 141 38 | 174 47 | 162 48 | 196 83 | 269 150 | 252 123 | 185 76 | 179 48 | 160 29 | 81 13 | 14 1 |
| 6 | Blackened sodium silico-fluoride at bud development. | Flowers opened Bolls matured out of the above flowers | .. | .. | 85 49 | 233 111 | 306 110 | 464 182 | 560 221 | 272 53 | 88 4 | 28 | 4 | 1 | .. |
| 6-A | Control for No. 6 | Flowers opened Bolls matured out of the above flowers | .. | .. | 51 14 | 97 28 | 109 31 | 154 71 | 228 116 | 244 133 | 178 96 | 135 43 | 108 21 | 86 8 | 34 2 |
| 5 | White sodium silico-fluoride at bud development. | Flowers opened Bolls matured out of the above flowers | 1 | .. | 77 49 | 154 55 | 219 28 | 396 176 | 282 184 | 87 31 | 28 8 | 28 3 | 19 3 | 8 1 | 19 3 |
| 5-A | Control for No. 5 | Flowers opened Bolls matured out of the above flowers | .. | .. | 76 27 | 159 56 | 137 56 | 260 112 | 375 202 | 369 179 | 298 104 | 271 66 | 210 41 | 136 8 | 43 5 |

TABLE XXXVIII

Weekly flower-opening and success of flowers to mature bolls from 20 plants in each case from plots dusted with insecticides and their controls

(1928-29)

| Plot No. | Treatment | | 3rd November | 10th November | 17th November | 24th November | 1st December | 8th December | 15th December | 22nd December | 29th December | 5th January | 12th January | 19th January | 26th January | 2nd February |
|----------|---|--|--------------|---------------|---------------|---------------|--------------|--------------|---------------|---------------|---------------|-------------|--------------|--------------|--------------|--------------|
| 1 | Blackened calcium arsenate at shoot attack | Flowers opened | 7 | 51 | 262 | 102 | 157 | 135 | 107 | 182 | 137 | 125 | 116 | 48 | 18 | 4 |
| | | Bolls matured out of the above flowers | 4 | 9 | 78 | 38 | 42 | 40 | 68 | 106 | 59 | 44 | 39 | 9 | 3 | .. |
| 1-A | Control for No. 1 | Flowers opened | | 3 | 60 | 120 | 135 | 237 | 269 | 317 | 215 | 175 | 168 | 72 | 27 | 5 |
| | | Bolls matured out of the above flowers | | 21 | 20 | 99 | 42 | 83 | 120 | 150 | 75 | 30 | 24 | 7 | 5 | 1 |
| 3 | Blackened calcium arsenate at boll development. | Flowers opened | | 1 | 85 | 149 | 170 | 295 | 462 | 472 | 293 | 173 | 69 | 13 | .. | .. |
| | | Bolls matured out of the above flowers | | 1 | 39 | 88 | 75 | 159 | 212 | 116 | 17 | 3 | 1 | 2 | .. | .. |
| 3 A | Control for No. 3 | Flowers opened | | | 75 | 110 | 145 | 226 | 321 | 269 | 184 | 139 | 106 | 58 | 31 | .. |
| | | Bolls matured out of the above flowers | | | 25 | 49 | 46 | 70 | 147 | 119 | 62 | 21 | 17 | 14 | 3 | .. |

TABLE XXXIX

Dates of dusting insecticides in different plots and the total quantity of the insecticide used with the approximate cost of the poisons required

(1929-30)

| Plot No. | Treatment | Date of 1st dusting | Date of 2nd dusting | Date of 3rd dusting | Date of 4th dusting | Date of 5th dusting | Total dust required calculated per acre (main insecticide only) | Approximate cost of the total insecticides used per acre |
|----------|--|---------------------|---------------------|---------------------|---------------------|---------------------|---|--|
| 2 | Blackened calcium arsenate at 20 lbs. mixture per acre per turn. | 25th Novem-ber. | 9th Decem-ber. | 19th Decem-ber. | 24th Decem-ber. | 3rd January . | 52 | Rs. A. P. 13 0 0 |
| 3 | White calcium arsenate at 20 lbs. per acre per turn and nicotine sulphate when aphid appear. | Do. | Do. | Do. | Do. | Do.* | 101 | 25 4 0 |
| 4 | White calcium arsenate dusting at 10 lbs. per acre per dusting. | Do. | Do. | Do. | Do. | Do. | 54 | 13 8 0 |
| 5 | Blackened sodium silico-fluoride, dusting at 40 lbs. of mixture per acre per dusting. | 29th Novem-ber | 16th Decem-ber. | 25th Decem-ber. | 4th January . | .. | 111 | 27 12 0 |
| 6 | Blackened sodium silico-fluoride at 40 lbs. mixture per acre per turn and nicotine sulphate when aphid appear. | Do. | Do. | Do. | Do.* | .. | 110 | 27 8 0 |
| 7 | White sodium silico-fluoride at 20 lbs. per acre per dusting. | Do. | Do. | Do. | Do. | .. | 94 | 23 8 0 |

* With the addition of nicotine sulphate.

TABLE XL

Weekly flower-opening and success of flowers into mature bolls from 20 plants from plots dusted with insecticides and their controls

(1929-30)

| Plot No. | Treatment | | 18th October | 25th October | 2nd November | 9th November | 16th November | 23rd November | 30th November | 7th December | 14th December | 21st December | 28th December | 4th January | 11th January | 18th January | 25th January | 1st February | 8th February | 15th February |
|----------|--|---|--------------|--------------|--------------|--------------|---------------|---------------|---------------|--------------|---------------|---------------|---------------|-------------|--------------|--------------|--------------|--------------|--------------|---------------|
| 2 | Blackened calcium arsenate at 20 lbs an acre. | Flowers opened | .. | 4 | 5 | 7 | 17 | 32 | 32 | 84 | 123 | 140 | 165 | 236 | 246 | 239 | 153 | 78 | 8 | 2 |
| | | Bolls matured out of the above flowers. | .. | .. | .. | .. | .. | 3 | 9 | 44 | 78 | 84 | 101 | 148 | 87 | 11 | 4 | .. | .. | .. |
| | Control for No. 2 | Flowers opened | .. | 1 | 5 | 7 | 6 | 7 | 9 | 44 | 102 | 139 | 131 | 248 | 279 | 237 | 235 | 134 | 35 | .. |
| | | Bolls matured out of the above flowers | .. | .. | .. | .. | .. | .. | 4 | 17 | 49 | 74 | 82 | 175 | 151 | 53 | 9 | 10 | 1 | .. |
| 3 | White calcium arsenate at 20 lbs to an acre and nicotine sulphate when aphid appear. | Flowers opened | .. | .. | 2 | 9 | 23 | 27 | 37 | 76 | 117 | 145 | 211 | 285 | 252 | 211 | 105 | 13 | .. | .. |
| | | Bolls matured out of the above flowers. | .. | .. | .. | .. | 3 | 4 | 15 | 47 | 79 | 100 | 127 | 117 | 40 | 5 | .. | 1 | .. | .. |
| | Control for No 3 | Flowers opened | .. | 4 | 7 | 17 | 23 | 27 | 30 | 80 | 113 | 133 | 132 | 170 | 241 | 222 | 203 | 92 | 24 | 2 |
| | | Bolls matured out of the above flowers | .. | .. | 1 | 3 | .. | 2 | 8 | 36 | 49 | 64 | 80 | 120 | 135 | 46 | 9 | 3 | 2 | .. |
| 4 | White calcium arsenate at 10 lbs. to an acre. | Flowers opened | .. | 5 | 3 | 5 | 19 | 33 | 27 | 41 | 114 | 121 | 166 | 212 | 307 | 190 | 178 | 58 | 8 | .. |
| | | Bolls matured out of the above flowers. | .. | .. | .. | .. | .. | 6 | 8 | 28 | 85 | 69 | 90 | 94 | 99 | 18 | 1 | .. | .. | .. |
| | Control for plot No. 4 | Flowers opened | .. | .. | 1 | 6 | 24 | 21 | 33 | 64 | 101 | 118 | 156 | 223 | 237 | 208 | 126 | 44 | 10 | 3 |
| | | Bolls matured out of the above flowers. | .. | .. | .. | .. | 1 | 3 | 6 | 30 | 65 | 63 | 103 | 125 | 90 | 24 | 2 | .. | 1 | .. |

TABLE XLI

Population of Earias larvae on weeds at Surat in 1928-29

| Week ending | <i>Hibiscus rugosus</i> | | <i>Hibiscus manihot</i> | | <i>Abutilon graveolens</i> | | <i>Abutilon indicum</i> | |
|-------------------------|-------------------------|---------------|-------------------------|---------------|----------------------------|---------------|-------------------------|---------------|
| | No. of plants examined | No. of larvae | No. of plants examined | No. of larvae | No. of plants examined | No. of larvae | No. of plants examined | No. of larvae |
| 14th July 1928 . . | 300 | .. | .. | .. | 30 | .. | .. | .. |
| 21st July 1928 . . | 500 | .. | .. | .. | 125 | .. | .. | .. |
| 28th July 1928 . . | 1,000 | 1 | 275 | .. | 140 | .. | . | .. |
| 4th August 1928 . . | 850 | | 200 | .. | 150 | .. | . | .. |
| 11th August 1928 . . | 1,100 | | 250 | . | 215 | 1 | 60 | .. |
| 18th August 1928 . . | .. | . | 110 | . | 90 | 5 | 45 | .. |
| 25th August 1928 . . | 600 | 3 | 20 | .. | 40 | .. | .. | .. |
| 1st September 1928 . . | 200 | 1 | .. | .. | 50 | 7 | .. | .. |
| 8th September 1928 . . | 150 | .. | .. | | 50 | .. | .. | .. |
| 15th September 1928 . . | 200 | .. | .. | .. | .. | .. | 45 | .. |
| 22nd September 1928 . . | 100 | .. | 40 | .. | .. | .. | .. | .. |
| 29th September 1928 . . | 150 | .. | 50 | .. | .. | .. | 24 | 29 |
| 6th October 1928 . . | 50 | .. | 50 | 40 | .. | .. | 7 | 36 |
| 13th October 1928 . . | 60 | .. | 55 | 65 | .. | .. | 5 | 75 |
| 20th October 1928 .. | 80 | 12 | 9 | 13 | . | .. | 2 | 14 |
| 27th October 1928 . . | .. | . | 5 | 35 | .. | .. | 2 | 9 |
| 3rd November 1928 . . | 200 pods. | 11 | 10 | 21 | .. | .. | .. | |
| 10th November 1928 . . | 100 pods. | 7 | 10 | 20 | .. | .. | .. | . |

TABLE XLII

Population of Earias larvae on weeds at Surat in 1929-30

| Week ending | <i>Hibiscus rugosus</i> | | <i>Hibiscus mansuetor</i> | | <i>Hibiscus ficulneus</i> | | <i>Hibiscus pandurus</i> — forms | | <i>Abutilon graveolens</i> | | <i>Abutilon indicum</i> | |
|-----------------------|-------------------------|-----------------|---------------------------|------------------|---------------------------|-----------------|-------------------------------------|------------------|----------------------------|------------------|-------------------------|------------------|
| | Plants examined | No of larvae | Plants examined | No. of larvae | Plants examined | No of larvae | Plants examined | No. of larvae | Plants examined | No. of larvae | Plants examined | No. of larvae |
| 4th July 1929 . | .. | . | .. | .. | . | .. | .. | .. | 25 | .. | .. | .. |
| 13th July 1929 . | 450 | . | .. | .. | .. | .. | .. | .. | 25 | .. | .. | .. |
| 20th July 1929 . | 700 | .. | .. | .. | .. | .. | .. | .. | 25 | .. | .. | .. |
| 27th July 1929 . | 500 | .. | 100 | .. | .. | .. | .. | .. | 44 | .. | 20 | .. |
| 3rd August 1929 . | 500 | .. | .. | .. | .. | .. | .. | .. | 20 | .. | .. | .. |
| 10th August 1929 . | 500 | .. | 50 | .. | .. | .. | .. | .. | .. | .. | 25 | .. |
| 17th August 1929 . | 500 | .. | 25 | .. | 15 | .. | 25 | .. | 25 | .. | 25 | .. |
| 24th August 1929 . | 500 | .. | 25 | .. | 8 | . | 30 | .. | 20 | .. | 5 | .. |
| 31st August 1929 . | 450 | .. | 25 | .. | 5 | .. | 25 | .. | 11 | .. | 5 | .. |
| 7th September 1929 . | 250 | 3 | 50 | .. | 3 | .. | 10 | .. | 8 | 1 | 8 | 3 |
| 14th September 1929 . | 100 | 2 | 50 | .. | .. | .. | 15 | .. | 10 | 3 | 3 | .. |
| 21st September 1929 . | 100 | .. | 50 | 2 | .. | .. | 10 | .. | 10 | 17 | 10 | 27 |
| 28th September 1929 . | 100 | 1 | 33 | 13 | 5 | .. | 10 | 8 | 11 | 7 | 3 | 20 |
| 5th October 1929 . | 75 | .. | 25 | 3 | 2 | .. | 10 | 4 | 10 | .. | 3 | .. |
| 12th October 1929 . | 60 | .. | 25 | 15 | .. | .. | 10 | 9 | 5 | .. | 4 | 2 |
| 19th October 1929 . | 75 | 11 | 16 | 7 | .. | .. | 10 | 21 | 7 | 12 | 3 | 2 |
| 26th October 1929 . | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 2nd November 1929 . | 50 | 8 | 10 | 25 | 1 | .. | 10 | 6 | 5 | 13 | 3 | 3 |
| 9th November 1929 . | 10 | 4 | 7 | 2 | .. | .. | .. | .. | 4 | 16 | 3 | 25 |
| 16th November 1929 . | 10 | 4 | 6 | .. | .. | .. | 10 | 3 | .. | .. | 3 | .. |

TABLE XLIII
Population of *Earias* larvae on weeds at Surat in 1930-31

| Week ending | <i>Hibiscus rugosus</i> | | <i>Hibiscus mantali</i> | | <i>Hibiscus scaberrimus</i> | | <i>Hibiscus grandifolius</i> —formis | | <i>Abutilon graveolens</i> | | <i>Abutilon indicum</i> | |
|---------------------|-------------------------|---------------|-------------------------|---------------|-----------------------------|---------------|---|--------|----------------------------|--------|-------------------------|--------|
| | Plants examined | No. of larvae | Plants examined | No. of larvae | Pods | No. of larvae | Pods | Larvae | Pods | Larvae | Pods | Larvae |
| 3rd July 1930 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 10th July 1930 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 17th July 1930 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 24th July 1930 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 31st July 1930 | 50 | .. | 25 | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 7th August 1930 | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 15th August 1930 | 50 | .. | 13 | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 22nd August 1930 | 75 | .. | 30 | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 29th August 1930 | 110 | .. | 30 | .. | .. | .. | .. | .. | .. | .. | .. | .. |
| 5th September 1930 | 103 | .. | 25 | .. | .. | .. | .. | .. | .. | .. | 100 | 19 |
| 12th September 1930 | 50 | .. | 30 | .. | .. | .. | .. | .. | .. | .. | 16 | 1 |
| 19th September 1930 | 47 | .. | 10 | .. | .. | .. | .. | .. | .. | .. | 67 | 13 |
| 26th September 1930 | 123 pods. | 2 | 10 | 4 | .. | .. | .. | .. | 14 | 4 | .. | .. |
| 3rd October 1930 | 181 | 5 | 10 | 10 | .. | .. | 20 | 1 | .. | .. | 104 | 25 |
| 10th October 1930 | 139 | .. | 10 | 49 | 28 | 1 | 5 | 2 | 50 | 27 | 54 | 16 |
| 17th October 1930 | 225 | 2 | 10 | 4 | 55 | 6 | 25 | 6 | 250 | 48 | 150 | 18 |
| 24th October 1930 | 118 | 4 | 15 | 21 | 105 | 20 | 19 | 1 | 231 | 30 | 106 | 24 |
| 31st October 1930 | 50 | 1 | 25 pods. | 12 | 30 | 5 | 58 | 6 | 100 | 29 | 60 | 11 |
| 7th November 1930 | 63 | 6 | 14 | 22 | 96 | 11 | 63 | 13 | 123 | 26 | 90 | 9 |
| 14th November 1930 | 20 | .. | 25 | 10 | 67 | 18 | 76 | 7 | 200 | 100 | 63 | 4 |
| 21st November 1930 | .. | .. | .. | .. | 10 | .. | 126 | 5 | 163 | 10 | 66 | 7 |
| 28th November 1930 | .. | .. | 32 | 12 | 50 | .. | 86 | 17 | 125 | 16 | 90 | 4 |

TABLE XLIV

Population of Earias larvae on plants of Abutilon indicum examined on the 15th of October 1927

(A case of very heavy attack)

| No. of affected pods | Larvae found in each form | Total larvae found |
|----------------------|---------------------------|--------------------|
| 78 | 1 | 78 |
| 21 | 2 | 42 |
| 7 | 3 | 21 |
| 1 | 4 | 4 |
| Total . 107 | .. | 145 |

TABLE XLV

Population of Earias larvae on Abutilon during summer

(Plants found in gardens of Surat)

| Date | No. of plants | No. of pods | Total Earias |
|---------------------------|---------------|-------------|--------------|
| 20th April 1929 | 7 | 36 | 5 |
| 27th April 1929 | 28 | 122 | 19 |
| 4th May 1929 | 12 | 9 | 2 |
| 11th May 1929 | .. | .. | .. |
| 18th May 1929 | 25 | 34 | 9 |
| 25th May 1929 | 25 | 107 | 23 |
| 1st June 1929 | 15 | 11 | 2 |
| 8th June 1929 | 25 | 175 | 64 |
| 15th June 1929 | 25 | 67 | 28 |
| 22nd June 1929 | 25 | 71 | 8 |
| 29th June 1929 | 25 | 33 | 4 |
| 6th July 1929 | 25 | .. | .. |
| 13th July 1929 | 25 | 41 | .. |
| 20th July 1929 | 25 | .. | .. |
| 27th July 1929 | 64 | 2 | .. |

TABLE XLVI

*Population of Earias larvae on bhendi plants (Hibiscus esculentus) in summer
(1928-29)*

| Date | No. of plants | Earias worms |
|---------------------------|---------------|--------------|
| 25th April 1928 | 100 | 36 |
| 3rd May 1928 | 100 | 81 |
| 11th May 1928 | 100 | 74 |
| 16th May 1928 | 100 | 54 |
| 22nd May 1928 | 100 | 136 |
| 2nd June 1928 | 100 | 303 |
| 5th June 1928 | 100 | 126 |
| 12th June 1928 | 100 | 116 |
| 19th June 1928 | 100 | 95 |
| 27th June 1928 | 75 | 19 |
| 30th June 1928 | 50 | 50 |
| 4th July 1928 | 100 | 33 |
| 10th July 1928 | 100 | 23 |
| 17th July 1928 | 100 | 8 |
| 23rd July 1928 | 200 | 7 |
| 30th July 1928 | 200 | 65 |

TABLE XLVII

Population of Earias larvae on sprouts of standing cotton sticks at Surat

(50 plants each time)

| Weeks | 1928-29 SEASON | | | Weeks | 1929-30 SEASON | | |
|-------------------|----------------|--------------|--------------|-------------------|----------------|--------------|--------------|
| | No. of buds | No. of bolls | Total Earias | | No. of buds | No. of bolls | Total Earias |
| 2nd March 1929 . | .. | 832 | 112 | 8th March 1930 . | .. | 668 | 26 |
| 9th March 1929 . | .. | 350 | 56 | 15th March 1930 . | .. | 134 | 8 |
| 16th March 1929 . | 10 | 70 | 34 | 22nd March 1930 . | .. | 26 | 2 |
| 23rd March 1929 . | 34 | 20 | 8 | 29th March 1930 . | .. | 6 | .. |
| 30th March 1929 . | 75 | 2 | 2 | 5th April 1930 . | 16 | .. | 1 |
| 6th April 1929 . | 203 | 4 | 14 | 12th April 1930 . | 43 | 1 | 6 |
| 13th April 1929 . | 319 | 16 | 27 | 19th April 1930 . | 198 | 3 | 9 |
| 20th April 1929 . | 554 | 26 | 33 | 26th April 1930 . | 263 | 6 | 16 |
| 27th April 1929 . | 469 | 39 | 57 | 3rd May 1930 . | 323 | 10 | 27 |
| 4th May 1929 . | 310 | 44 | 76 | 10th May 1930 . | 395 | 19 | 11 |
| 11th May 1929 . | 337 | 26 | 41 | 17th May 1930 . | 591 | 49 | 30 |
| 18th May 1929 . | 354 | 21 | 36 | 24th May 1930 . | 628 | 86 | 8 |
| 25th May 1929 . | 231 | 17 | 25 | 31st May 1930 . | 620 | 243 | 22 |
| 1st June 1929 . | 467 | 54 | 48 | | | | |

TABLE XLVIII

Weekly flower-opening, success of weekly flowers into bolls, and shedding of bolls due to boll-worms from the ratoon crop and also from the new crop of cotton

(1928-29)

(10 plants in each case)

| | 6th October | 13th October | 20th October | 27th October | 3rd November | 10th November | 17th November | 24th November | 1st December | 8th December | 15th December | 22nd December | 29th December | 5th January | 12th January | 19th January | 26th January | 2nd February | 9th February | 16th February | 23rd February | Total |
|---------------------------------------|-------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|--------------|--------------|---------------|---------------|---------------|-------------|--------------|--------------|--------------|--------------|--------------|---------------|---------------|-------|
| <i>Ratoon crop.</i> | | | | | | | | | | | | | | | | | | | | | | |
| Flower opening . . | 48 | 50 | .. | .. | .. | .. | 9 | 3 | 1 | 5 | 14 | 19 | 17 | 15 | 15 | 8 | 17 | 5 | 3 | 2 | 3 | 234 |
| Bolls matured from the above flowers. | 1 | .. | .. | .. | .. | .. | 1 | .. | .. | 1 | 3 | 3 | 8 | 4 | 3 | 3 | 5 | 3 | 2 | 2 | 3 | 42 |
| Boll shedding (Total) . | .. | 74 | 15 | 2 | .. | 1 | 4 | 6 | 5 | 3 | 4 | 10 | 10 | 10 | 12 | 11 | 6 | 5 | 5 | 9 | .. | 192 |
| Bolls shed due to boll-worms. | .. | 70 | 15 | 2 | .. | 1 | 4 | 6 | 5 | 1 | 3 | 8 | 4 | 6 | 1 | .. | 2 | 2 | 3 | 3 | .. | 136 |
| <i>New crop of 1928-29.</i> | | | | | | | | | | | | | | | | | | | | | | |
| Flower opening . . | .. | .. | .. | .. | .. | .. | 26 | 39 | 59 | 67 | 108 | 97 | 69 | 40 | 27 | 11 | 3 | 2 | .. | .. | .. | 548 |
| Bolls matured from the above flowers. | .. | .. | .. | .. | .. | .. | 13 | 13 | 22 | 37 | 64 | 48 | 19 | 4 | 4 | 3 | 1 | .. | .. | .. | .. | 228 |
| Boll shedding (Total) . | .. | .. | .. | .. | .. | .. | 2 | 13 | 15 | 30 | 40 | 31 | 33 | 38 | 50 | 36 | 23 | 6 | 2 | 1 | .. | 320 |
| Bolls shed due to boll-worms. | .. | .. | .. | .. | .. | .. | 2 | 10 | 13 | 24 | 22 | 12 | 6 | 6 | 3 | 3 | 1 | .. | .. | .. | .. | 102 |

TABLE XLIX

Population of Earias larvae on the sprouts from cotton stumps from the Government Farm and cultivators' area till the sowing of the new crops of 1928-29

| Date | CULTIVATORS' FIELDS | | FARM AREA | |
|------------------------------|---------------------|---------------|------------------|---------------|
| | Sprouts examined | No. of larvae | Sprouts examined | No. of larvae |
| 10th May 1928 | 75 | <i>Ntl</i> | .. | .. |
| 19th May 1928 | 75 | 1 | .. | .. |
| 25th May 1928 | 75 | 1 | .. | .. |
| 1st June 1928 | 100 | <i>Ntl</i> | 100 | 2 |
| 7th June 1928 | 200 | 1 | 200 | 1 |
| 15th June 1928 | 200 | 3 | 200 | <i>Ntl</i> |
| 22nd June 1928 | 200 | 2 | 200 | 4 |
| 30th June 1928 | 200 | 1 | 200 | 9 |
| 9th July 1928 | 200 | <i>Ntl</i> | 200 | 2 |
| 11th July 1928 | 250 | 2 | .. | .. |
| 20th July 1928 | 200 | 8 | 500 | 14 |
| 31st July 1928 | 200 | 6 | 200 | 6 |
| 11th August 1928 | 200 | 3 | 400 | 1 |
| 15th August 1928 | 200 | 15 | .. | .. |
| 23rd August 1928 | 200 | 29 | .. | .. |
| 31st August 1928 | 200 | 13 | .. | .. |
| 5th September 1928 | 200 | 13 | .. | .. |

TABLE L

Population of Earias larvae on the sprouts from the cotton stumps from the cultivators' area till the sowing of the new crops of 1929-30

| KOSAD AREA | | | PIPLOD AREA | | |
|---------------------|---|-----------------------------|---------------------|---|-----------------------------|
| Date | No. of stumps from which sprouts were collected | No. of <i>Earias</i> larvae | Date | No. of stumps from which sprouts were collected | No. of <i>Earias</i> larvae |
| 16th May 1929 | 100 | 11 | 4th May 1929 | 100 | .. |
| 21st May 1929 | 100 | . | 11th May 1929 | 100 | 2 |
| 30th May 1929 | 75 | 1 | 18th May 1929 | 100 | .. |
| 1st June 1929 | 100 | 8 | 25th May 1929 | 200 | 2 |
| 8th June 1929 | 55 | .. | 1st June 1929 | 200 | 5 |
| 14th June 1929 | 100 | .. | 12th June 1929 | 200 | 1 |
| 28th June 1929 | 150 | .. | 22nd June 1929 | 200 | .. |
| 6th July 1929 | 200 | 1 | 27th June 1929 | 200 | .. |
| 27th July 1929 | 100 | .. | 6th July 1929 | 200 | 3 |
| 7th September 1929 | 100 | 7 | 18th July 1929 | 200 | 2 |
| 20th September 1929 | 100 | .. | 25th July 1929 | 200 | 1 |
| | | | 3rd August 1929 | 200 | .. |
| | | | 10th August 1929 | 200 | 2 |
| | | | 17th August 1929 | 150 | 2 |
| | | | 25th August 1929 | 100 | 3 |
| | | | 6th September 1929 | 100 | 3 |
| | | | 14th September 1929 | 50 | 5 |

APPENDIX II

INDIAN CENTRAL COTTON COMMITTEE, TECHNOLOGICAL LABORATORY

SPINNING TEST REPORTS

1929-30 crop

Spinning Test Report No. 151.—Two Samples of Surat 1027 A. L. F. submitted by the Cotton Entomologist, Surat

| Sample No. | Cotton | Season | Weight |
|------------|-----------------------------------|---------|--------|
| | | | Lb. |
| 774 | 1027 A. L. F., sample A | 1929-30 | 10 |
| 775 | 1027 A. L. F., sample B | 1929-30 | 8½ |

I. Grader's Report

| | CAGED PLANTS | CONTROL PLANTS |
|---|---------------------------|------------------|
| | Sample A | Sample B |
| Contract valued under | Broach | Broach. |
| Class | Extra superfine | Extra superfine. |
| Colour | White | White. |
| Staple length | Full 1½ inch | 1½ inch. |
| Staple strength | Good | Good. |
| Regularity | Good | Poor. |
| Valued above or below contract rate | Rs. 150 on | Rs. 120 on. |
| Basis | Rs. 200 | Rs. 200. |
| Date of valuation | 11th July 1930 | 11th July 1930. |
| Remarks | | |

II. Fibre Particulars

| Mean group-length in eighth of an inch | PERCENTAGE | |
|--|------------|----------|
| | Sample A | Sample B |
| 1. Fibre-length Distribution (Balls Sorter)— | | |
| 3 | 0 | 0.1 |
| 4 | 1.4 | 1.8 |
| 5 | 2.4 | 3.6 |
| 6 | 4.8 | 7.1 |
| 7 | 8.9 | 17.9 |
| 8 | 17.2 | 32.1 |
| 9 | 30.9 | 25.7 |
| 10 | 24.4 | 10.0 |
| 11 | 7.6 | 0.7 |
| 12 | 2.4 | 0 |
| 2. Fibre-length (inch)— | | |
| (a) By Balls Sorter | 1.10 | 0.99 |
| (b) By Baer Sorter | 1.07 | 1.03 |
| 3. Fibre-weight per inch (Millionth of an ounce) | 0.202 | 0.167 |

III. Spinning Tests

1. *Treatment*.—These cottons were passed through the Porcupine Crighton (once only), Hopper, Scutcher (3 times), Card, Drawing (2 heads), Slubber, Inter, Rover, and spun from single hank roving in Ring Frame No. 1.

2. *Spinning Master's Report*—

Sample A.—The cotton is white; bright; perfectly clean; has a good feel; fairly well-ginned (seems to be 'Charkha' ginned). The staple is fine, fairly regular and strong.

Sample B.—The cotton is white; perfectly clean; bright; well-ginned; has a good soft feel. The staple is fine, regular and strong.

3. *Spinning Test details and results*.—See table appended.

IV. Remarks

Fibre.—Sample A is 7 per cent longer, but 21 per cent coarser than Sample B.

Waste.—Both samples gave low blow-room losses. Their card room losses were about normal.

Breakages.—Yarn breakages in the ring frame were comparatively few in all cases.

Yarns.—Both samples gave “even” 20’s and “fairly even” 34’s. All the yarns are fairly free from nep. From the yarn-strength results these cottons are adjudged suitable for the following highest standard warp counts:—

| Sample No. | Cotton | Highest standard warp counts |
|------------|----------------------------------|------------------------------|
| 774 | 1027 A. L. F. Sample A | 33’s |
| 775 | 1027 A. L. F. Sample B | 37’s |

Conclusion.—The lint from the caged plants, though possessing a somewhat longer staple, is decidedly coarser and rather inferior in spinning performance to the lint from the control plants.

(Sd.) A. JAMES TURNER,

Director, Technological Laboratory.

Dated the 30th August 1930.

INDIAN CENTRAL COTTON COMMITTEE, TECHNOLOGICAL LABORATORY

Spinning Test Report No. 236. Samples of Surat cotton submitted by the Cotton Physiologist, Surat

| Sample No. | Cotton | Season | Gross Weight |
|------------|------------------------------|---------|--------------|
| | | | Lbs. |
| 1099 | Protected plants | 1930-31 | 14½ |
| 1100 | Unprotected plants | 1930-31 | 15½ |

I. Grader's Report

| | Protected plants | Unprotected plants |
|--|------------------------------|--------------------|
| Contract valued under | Broach | Broach. |
| Class | Extra superfine | Extra superfine |
| Colour | Creamy | Creamy |
| Staple length | 1 $\frac{1}{16}$ " | 1 $\frac{1}{16}$ " |
| Staple strength | Good | Good. |
| Regularity | Fair | Fair. |
| Value above or below contract rate | Rs 70 on | Rs 75 on. |
| Basis | Rs 185 | Rs 185 |
| Date of valuation | 25th July 1931 | 25th July 1931. |
| Remarks | | |

II. Fibre Particulars

| Mean group-length in eighth of an inch | PERCENTAGE. | |
|--|------------------|--------------------|
| | Protected plants | Unprotected plants |
| 1. Fibre-length Distribution (Balls Sorter)— | | |
| 3 | 1.5 | 0.6 |
| 4 | 2.6 | 1.4 |
| 5 | 3.3 | 2.8 |
| 6 | 7.0 | 6.3 |
| 7 | 15.1 | 13.5 |
| 8 | 28.1 | 31.8 |
| 9 | 25.8 | 32.9 |
| 10 | 12.9 | 9.3 |
| 11 | 3.7 | 1.4 |
| 2. Fibre-length (inch)— | | |
| (a) By Balls Sorter | 1.01 | 1.02 |
| (b) By Baer Sorter | 1.02 | 0.99 |
| 3. Fibre-weight per inch (millionth of an ounce) | 0.192 | 0.184 |

III. Spinning Tests

1. *Treatment*.—All these cottons were passed through the Porcupine, Crighton (once only), Hopper, Scutcher (3 times), Card, Drawing (2 heads), Slubber, Inter, Rover, and Spun from single hank roving in ring frame No. 1.

2. *Spinning Master's Report*.—

Protected plants.—The cotton is white in colour; bright; perfectly clean; has a soft silky feel; very well-ginned. The card sliver is clean; web good and ten flat strips weigh 9.7 grams.

Unprotected plants.—The cotton is white in colour; bright; perfectly clean; has good soft feel; well-ginned. The card sliver is clean; web even; nep-free; and of a nice tension and ten flat strips weigh 11.2 grams.

3. *Spinning Test Details and Results*.—See table appended.

IV. Remarks

Fibre.—Sample Nos. 1099 and 1100 are slightly longer than 1027 A. L. F. (spaced 3'×3': Physiological section). The fibre weight per inch of these two samples is the same as that of 1027 A. L. F.

Waste.—The cottons were picked in a very clean condition and gave a low blow-room loss. The card-room loss was normal in each case.

Breakages

Yarns.—The cottons gave "even to fairly even" 30's, but the yarns of lint from protected plants are slightly less even in the higher count. The yarns given by the lint from the protected and unprotected plants are slightly neppy. From the yarn-strength results, these cottons are adjudged suitable for the following highest warp standard counts:

| Sample No. | Cotton | Counts |
|------------|------------------------------|--------|
| 1099 | Protected plants | 33's |
| 1100 | Unprotected plants | 35's |

Conclusions.—It is noteworthy that the lint from the unprotected plants gave yarns which are stronger than those from the protected plants. There is very little variation in the spinning value of these two samples as compared with the last year's results.


(Sd.) N. AHMAD,

Director, Technological Laboratory.


Dated, 13th August 1931.

MULTIPLE TEMPERATURE INCUBATOR.

SECTION



PLAN



Plan and section of a multiple temperature incubator

Scale :- $\frac{3}{8}$ " = 1'

APPENDIX III

DESCRIPTION OF THE MULTIPLE TEMPERATURE INCUBATOR

A device for securing different ranges of fixed temperature, higher as well as lower than the surrounding air, is described in the Technical Bulletin No. 38 of the Ministry of Agriculture, Egypt. The principle underlined in the device is, that when the two ends of a column of metal are maintained at two constant temperatures, various intermediate temperatures can be recorded at different points in the intervening space of the metal.

We felt the necessity of an instrument which would enable us to study the life of the boll-worms under varying temperatures, which are experienced during the different parts of the season. We, therefore, utilised the suggestion from the above Bulletin and prepared a Multiple Temperature Incubator, the construction and the working of which has been described below :—

CONSTRUCTION

Copper sheets $\frac{1}{8}$ inch in thickness were purchased and a trough (A-B), 16 feet in length 6 inches in height and 4 inches in breadth was prepared. One end of this trough was inserted into the copper tank (C), projecting 9 inches into it. The copper tank was one foot square. The other end projected two inches into another copper tank (D), which was two inches more in height than the above one (C). The whole length of the trough and the tank were placed in a rectangle wooden box one foot six inches in height and two feet four inches in width. Insulation was provided for the trough by placing an inch of layer of *kapas* round it and the remaining part of the box was filled in with saw dust. (Plate XXI).

The trough was divided into fifteen compartments (with wooden partitions) of one foot each with a separate cover for each. The covers were made up of four layers, two of which consisted of wooden planks $\frac{1}{4}$ inch in thickness and the other two of asbestos and felt respectively. Asbestos was sandwiched between the two wooden planks and the felt was struck to the bottom of the lower one. Holes were bored in the covers to accommodate the thermometers for reading the temperature in each one of the compartments.

WORKING

The tank (D) is filled with water which is continuously kept boiling. The tank C at the other end contains ice-cold water, temperature being maintained by continuous addition of ice. Thus the two ends of the trough (A and B) are kept at constant temperatures. A temperature gradient is thus set up between the two ends, giving different intermediate temperatures from one end to the other.

It takes about six hours to get regulated temperatures after the heating of the water is started and ice is placed in the other tank.

Absolute constant temperatures are not available but the diurnal variation in each of the compartment does not generally exceed four degrees Fahrenheit and is good enough for all purposes where greater regulation of the temperature is not necessary.

Ranges of temperature available in the twelve compartments beginning from the cold end are given below :—

| Compartments | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
|----------------------|----|----|----|----|----|----|----|----|----|-----|-----|-----|
| Range of temp. in F. | 58 | 66 | 73 | 79 | 80 | 84 | 89 | 90 | 96 | 100 | 110 | 122 |
| | to | to | to | to | to | to | to | to | to | to | to | to |
| | 64 | 70 | 75 | 81 | 83 | 86 | 91 | 93 | 98 | 102 | 112 | 125 |

APPENDIX IV

SPOTTED BOLL-WORMS IN KHANDESH

A considerable amount of information, which was collected mostly during 1928 and 1929 in Gujarat, showed that the Spotted Boll-worms did not aestivate and that they passed their period between two cotton-growing seasons (roughly from April to August) on the fresh growth from the cotton plants and their stumps in the general cotton area, and also on *bhendi* (*Hibiscus esculentus*), which was found growing in stray irrigated plots.

As regards Khandesh, it was known that the cotton crop of that district also was affected by the Spotted Boll-worms, but exact information about the incidence of this pest in that locality was not available, and knowledge about the means of carry-over of this pest in that tract, from one season to another was particularly lacking.

In Khandesh, the cotton crop is sown in June or in July with the earliest showers of rain, and the crop is ready in October or November in normal years. During exceptional seasons, however, the crop extends up to January, if heavy late rains are received. Thus there is a span of at least six months between the two cotton crops and the summer climate being rigorous and the soil not being very highly retentive of moisture, the conditions are not very favourable for resprouting of cotton plants after the end of the season. It was, therefore, necessary to study the sources of food of this pest in that tract during the off season, and to locate the source of fresh infection to the new crop of cotton. A few observations were, therefore, made from August 1929 to January 1931 in Khandesh. Mr. Nadkarni, the junior author, was in sole charge of this work.

INCIDENCE OF SPOTTED BOLL-WORMS DURING 1929 AND 1930

During 1929 the sowing of cotton on the Government farm at Jalgaon (East Khandesh) was done in the second week of June. The flower-bud formation commenced by the beginning of August, and it was about that time that the presence of Spotted Boll-worms was first noticed in the cotton fields during that season. The flower-bud formation increased very rapidly during the month of August and flower opening began by the last week of the same month. In spite of the large number of flower-buds and bolls, the population of these worms continued to be meagre in the cotton fields, till the beginning of October, when it suddenly increased and by the second week of October as many as 84 larvae were recorded on 100 plants of cotton. By this time, however, the first picking of cotton was ready and hence the cotton crop of that season practically escaped a serious damage.

TABLE I

Weekly population of Earias larvae on 100 plants of cotton from Government Farm at Jalgaon, 1929

| Week ending dates | Flower-buds | Bolls | Larvae in flower-buds | Larvae in bolls | Total larvae | REMARKS |
|---------------------------|-------------|-------|-----------------------|-----------------|--------------|-------------------------|
| 3rd August 1929 . . . | 263 | .. | 1 | .. | 1 | |
| 10th August 1929 . . . | 564 | .. | 1 | .. | 1 | |
| 17th August 1929 . . . | 990 | .. | 1 | .. | 1 | |
| 24th August 1929 . . . | 1,312 | 49 | 2 | 1 | 3 | |
| 31st August 1929 . . . | 1,487 | 149 | 5 | 1 | 6 | |
| 7th September 1929 . . . | 1,099 | 270 | .. | 1 | 1 | |
| 14th September 1929 . . . | 1,055 | 305 | 1 | 2 | 3 | |
| 21st September 1929 . . . | 718 | 622 | 3 | 3 | 6 | |
| 28th September 1929 . . . | 423 | 774 | 4 | 14 | 18 | |
| 5th October 1929 . . . | 102 | 734 | 2 | 69 | 71 | |
| 12th October 1929 . . . | 30 | 488 | 6 | 78 | 84 | First picking of kapas. |
| 19th October 1929 . . . | .. | 210 | .. | 42 | 42 | |
| 26th October 1929 . . . | 2 | 100 | .. | 11 | 11 | Second picking. |
| 2nd November 1929 . . . | .. | 55 | .. | 7 | 7 | |
| 9th November 1929 . . . | .. | 23 | .. | 3 | 3 | |
| 16th November 1929 . . . | 9 | 6 | .. | 1 | 1 | Third picking. |

It has been a practice in Khandesh to grow a few rows of *bhendi* (*Hibiscus esculentus*) along with cotton. Population of *Earias* larvae on these plants also was recorded by examining 15 plants of *bhendi*, collected from the cotton fields, once in a week.



TABLE II

Earias population on 15 plants of bhendi at Jalgaon

1929

| Weeks. | No. of flower-buds | No. of pods | Total No. of larvae | REMARKS |
|-------------------------------|--------------------|-------------|---------------------|-------------------------|
| 17th August 1929 | 181 | 36 | 12 | |
| 24th August 1929 | 135 | 82 | 7 | |
| 31st August 1929 | 40 | 104 | 10 | |
| 7th September 1929 | 8 | 59 | 29 | |
| 14th September 1929 | .. | 55 | 79 | |
| 21st September 1929 | .. | 39 | 51 | |
| 28th September 1929 | 25 | 58 | 41 | Plants began to dry up. |

PLATE XXII.



Showing weekly rainfall and periodical population of *Earias* larvae on cotton and *bhendi* plants at Jalgaon during 1929 and 1930

It is seen that comparatively a very much larger population of *Earias* larvae was present on 15 plants of *bhendi* than what was found on 100 cotton plants during the months of August and September. However, as soon as the *bhendi* plants dried up, the population on cotton suddenly increased, indicating thereby that the Spotted Boll-worms from *bhendi* plants diverted their attention to the flower-buds and bolls of cotton as soon as they were liberated from *bhendi* plants. (Plate XXII).

The vegetative shoots of cotton are not destroyed by this pest in Khandesh because the flower-buds on these plants are available very early; and besides the shoots of neglectum cottons are thin and woody and are not therefore as inviting to these worms as the succulent shoots of herbacium cottons.

During the next season both cotton and *bhendi* were sown at Jalgaon in the third week of June. The earliest population of *Earias* larvae was noticed in the shoots of *bhendi* plants by the middle of July. It did not, however, increase on these plants during the month of September in this season, chiefly because it was controlled by heavy rains during the first fortnight of that month. Consequently a sudden increase of *Earias* larvae on cotton plants at the end of the *bhendi* crop was not noticed.

TABLE III

Population of Earias larvae on 15 plants of bhendi from Government Farm, Jalgaon, 1930

| Weeks | No. of pods | No. of larvae | REMARKS |
|-------------------------------|-------------|---------------|-----------|
| 5th July 1930 | .. | .. | |
| 12th July 1930 | .. | 1 | In shoot. |
| 19th July 1930 | .. | .. | |
| 26th July 1930 | .. | 1 | In shoot. |
| 2nd August 1930 | .. | 1 | Do. |
| 9th August 1930 | 54 | 1 | |
| 16th August 1930 | 51 | Nil | |
| 23rd August 1930 | 73 | 3 | |
| 30th August 1930 | 61 | 4 | |
| 6th September 1930 | 44 | 4 | |
| 13th September 1930 | 27 | 3 | |
| 20th September 1930 | 81 | 6 | |
| 27th September 1930 | 19 | 5 | |

The population of Spotted Boll-worms on cotton gradually increased and reached its maximum at the end of October. Due to heavy rains in the month of September (Table XVII of this appendix) the cotton crop of this season was delayed and therefore a good number of flower-buds and bolls were available on these plants till the month of January. The damage caused by the Spotted Boll-worms to these forms continued during all this period and hence the cotton crop of this year suffered a heavier loss than the crop of the previous season.

TABLE IV

Weekly population of Earias larvae on 100 plants of cotton from the Government Farm at Jalgaon

1930-31

| Week ending dates | Total flower-buds | Total bolls | Larvae in flower-buds | Larvae in bolls | Total larvae |
|---------------------------|-------------------|-------------|-----------------------|-----------------|--------------|
| 9th August 1930 . . . | 24 | .. | .. | .. | .. |
| 16th August 1930 . . . | 50 | .. | .. | .. | .. |
| 23rd August 1930 . . . | 80 | .. | .. | .. | .. |
| 30th August 1930 . . . | 378 | .. | .. | .. | .. |
| 6th September 1930 . . . | 367 | 5 | .. | .. | .. |
| 13th September 1930 . . . | 1,171 | 90 | 8 | 1 | 9 |
| 20th September 1930 . . . | 540 | 170 | 8 | 2 | 10 |
| 27th September 1930 . . . | 445 | 685 | 4 | 9 | 13 |
| 4th October 1930 . . . | 289 | 499 | 6 | 7 | 13 |
| 11th October 1930 . . . | 300 | 585 | 12 | 29 | 41 |
| 18th October 1930 . . . | 497 | 458 | 27 | 38 | 65 |
| 25th October 1930 . . . | 213 | 576 | 21 | 87 | 108 |
| 1st November 1930 . . . | 205 | 458 | 29 | 84 | 113 |
| 8th November 1930 . . . | 199 | 315 | 19 | 58 | 77 |
| 15th November 1930 . . . | 336 | 154 | 20 | 31 | 51 |
| 22nd November 1930 . . . | 360 | 170 | 19 | 16 | 35 |
| 29th November 1930 . . . | 538 | 154 | 46 | 32 | 78 |
| 6th December 1930 . . . | 430 | 145 | 58 | 32 | 90 |
| 13th December 1930 . . . | 823 | 131 | 97 | 43 | 140 |
| 20th December 1930 . . . | 503 | 103 | 67 | 33 | 100 |
| 27th December 1930 . . . | 515 | 141 | 47 | 23 | 70 |
| 3rd January 1931 . . . | 456 | 80 | 7 | 6 | 13 |
| 10th January 1931 . . . | 622 | 140 | 47 | 13 | 60 |
| 17th January 1931 . . . | 800 | 344 | 69 | 72 | 141 |
| 24th January 1931 . . . | 926 | 338 | 44 | 31 | 75 |
| 31st January 1931 . . . | 781 | 196 | 26 | 55 | 81 |

The first picking of cotton was delayed till the end of October and the last was finished by the middle of December.

These observations show that the population of the Spotted Boll-worms in Khandesh first appears on the monsoon crop of *bhendi* and it rapidly increases on these plants during the months of August and September, if it is not checked by rains. As soon as the *bhendi* plants finish their life, all the moths concentrate on the cotton plants and damage the crop until it is over. This crop escapes a severe damage if it matures quickly before the pest increases on it. The pickings of cotton in Khandesh usually begin sometime during October or November and the last picking is over before the end of December but in some years it is delayed till the month of February.

TABLE V

Dates of picking kapas on Government Farm at Jalgaon

| Year | Dates of first picking | Dates of second picking | Dates of third picking |
|----------------|------------------------|-------------------------|------------------------|
| 1921 | 29th October . | 23rd November . | 16th December. |
| 1922 | 3rd November . | 16th December . | 9th February. |
| 1923 | 18th November . | 13th December . | 6th January. |
| 1924 | 20th October . | 20th November . | 26th December. |
| 1925 | 23rd October . | 6th November . | 11th December. |
| 1926 | 4th November . | 9th December . | 2nd January. |
| 1927 | 21st October . | 18th November . | 16th January. |
| 1928 | 5th November . | 19th November . | . |
| 1929 | 11th October . | 25th October . | 15th November. |

ESTIMATE OF DAMAGE TO THE COTTON CROP

Fifty cotton plants were kept under observation both during 1929 and 1930 for recording the flower opening and for examining the shed bolls.

TABLE VI

Flower opening and shedding of bolls from 50 plants at Jalgaon during 1929 and 1930

| Year | Total flowers opened | Total bolls shed | Bolls shed due to boll-worms | Percentage bolls shed due to boll-worms |
|----------------|----------------------|------------------|------------------------------|---|
| 1929 | 651 | 353 | 54 | 15.3 |
| 1930 | 621 | 277 | 147 | 53.0 |

It was found that in 1929 only about 15 per cent of the shed bolls showed injury caused by the Spotted Boll-worms, whereas during 1930 fifty-three per cent of the shed bolls indicated boll-worm damage (Plate XXIII).

In addition to these observations, 100 plants of cotton were grown by night-caging at Jalgaon during 1930. The caging work at Jalgaon was found to be rather difficult because the plants had to be spaced very close according to the local method and there could not be sufficient room between the plants and the curtains. It was particularly difficult to keep off the Pink Boll-worm moths, which remained hidden by the side of clods or in partially opened bolls and laid their eggs on the caged plants.

TABLE VII

Yield of kapas from the caged plants and their controls at Jalgaon in 1930



(In each case total figures from 100 plants are given)

| | Good kapas in tolas | Damaged kapas in tolas | | Total kapas in tolas |
|--------------------------------------|------------------------|------------------------|------------------------|-------------------------|
| | | Due to boll-worms | Due to other causes | |
| Caged plants | 84.7 | 10.5 | 1.4 | 96.7 |
| Control plants (group No. 1) | 62.5 | 11.5 | 2.0 | 76.0 |
| Control plants (group No. 2) | 63.7 | 10.7 | 1.0 | 75.5 |

The caged plants yielded about 26 per cent more *kapas* than the control plants, and probably this difference would have been greater if the developed bolls could have been effectively saved from damage by Pink Boll-worms.

SPOTTED BOLL-WORMS DURING THE OFF-SEASON

The cotton crop during 1929 was over at Jalgaon by the middle of November. After this the cotton plants began to put up fresh flower-buds and bolls, and therefore a plot of about ten *gunthas* was reserved on the Jalgaon Farm for noting the *Earias* population. One hundred fresh shoots were collected every week for examination.



Showing the weekly flower-opening, shedding of bolls and relative success of bolls from weekly flowers on cotton plants at Jalgaon.

TABLE VIII

Population of Earias kurvae from 100 shoots of cotton collected from the Jalgaon farm after the pickings were over

1929-30

| Weeks | Flower-buds | Bolls | Total <i>Earias</i> larvae |
|------------------------------|-------------|-------|----------------------------|
| 23rd November 1929. | 239 | 54 | 22 |
| 30th November 1929 | 329 | 62 | 56 |
| 7th December 1929 | 359 | 97 | 77 |
| 14th December 1929 | 497 | 101 | 83 |
| 21st December 1929 | 576 | 92 | 72 |
| 28th December 1929 | 385 | 49 | 34 |
| 4th January 1930 | 260 | 38 | 22 |
| 11th January 1930 | 260 | 66 | 20 |
| 18th January 1930 | 283 | 52 | 21 |
| 25th January 1930 | 198 | 60 | 12 |
| 1st February 1930 | 211 | 66 | 20 |
| 8th February 1930 | 252 | 67 | 35 |
| 15th February 1930 | 220 | 57 | 16 |
| 22nd February 1930 | 159 | 61 | 9 |
| 1st March 1930 | 174 | 100 | 6 |
| 8th March 1930 | 123 | 96 | 10 |
| 15th March 1930 | 197 | 116 | 14 |
| 22nd March 1930 | 151 | 111 | 10 |
| 29th March 1930 | 170 | 85 | 10 |

Spotted Boll-worms continued to breed in this plot till the end of March ; but the general cotton area in the District was almost completely cleared up by the end of January, as can be seen from the information collected regarding the standing cotton plots from two villages near Jalgaon. The total area under oservation was 150 acres at Mamrabad and 130 acres at Asoda.

TABLE IX

Approximate estimate of the area of the standing cotton plots at Mamrabad and Asoda after the harvest of cotton crop of 1929

| Dates of observation | Area with standing cotton stalks at Mamrabad | Area with standing cotton stalks at Asoda |
|-----------------------------|--|---|
| 8th December 1929 | 110 | 75 |
| 4th January 1930 | 40 | 40 |
| 11th January 1930 | 2 | 1 |
| 24th January 1930 | 0.5 | Nil |
| 1st February 1930 | 0.25 | Nil |

This must be considered to be a very early cleaning of the cotton fields, because in certain seasons the pickings of *kapas* continue till the end of January and consequently the removal of the cotton stalks continues for a couple of months more. It is unlikely, however, to find standing cotton fields in Khandesh after the month of March even if the season is late.

The cotton plants in Khandesh are removed by repeatedly working the blade harrow across the fields. Most of the plants come out with the roots and a few stumps that remain in the soil do not sprout up. The fresh growth from stumps which is so common in Gujarat is practically absent in Khandesh.

Hibiscus rugosus, *Hibiscus ficulneous*, *Hibiscus manihot* and plants of *Abutilon* sp. are chiefly the wild food plants of *Earias* in Khandesh; but none of them can be seen in green condition in the general cotton areas after the month of December. It was only at Kusumba in West Khandesh that a large number of plants of *Abutilon* sp. were found growing during summer in the irrigated plots.

Earias larvae, therefore cannot get any food in the general cotton fields in Khandesh after the month of March. There is, however, a large number of wells in several villages and there are small irrigated plots round many of these wells, which can supply water during the summer. It is in these plots that some of the food plants of *Earias* can thrive during summer.

There are about 12,165 acres in West Khandesh and 18,973 acres in East Khandesh which are irrigated by water from wells. As an instance we might state that 33 out of 111 villages of Jalgaon Taluka have more than 10 wells each, which supply water for irrigation, and of these there are 4 villages which have each more than 100 such wells.

It was decided to keep some of these plots under observation for recording the presence of the food plants of the Spotted Boll-worms in these plots. Three representative villages, viz. Nagardevala, Kusumba and Nimbhora, were, therefore, selected in addition to Jalgaon, for this purpose.

Some of the food plants, which were found in these plots and which could supply food to the *Earias* larvae during the period between the two cotton-growing seasons, were the following:—

- (1) *Abutilon* sp. (2) Standing cotton plants of the previous season. (3) *Bhendi* (*Hibiscus esculentus*) plants.

Abutilon sp.

This is the only weed, which sometimes continues to thrive during the summer months, in the irrigated plots in Khandesh. It is not, however, very common and it was seen in large numbers in the gardens at Kusumba only.

TABLE X

Pods from plants of Abutilon sp. from Kusumba

| Month of examination | No. of green capsules | No. of <i>Earias</i> larvae | REMARKS |
|----------------------|-----------------------|-----------------------------|--|
| January | 200 | 44 | Most of the larvae from these capsules belonged to the species, <i>Earias insulana</i> . |
| February | 150 | 33 | |
| March | 90 | 14 | |
| April | 30 | 13 | |
| May | .. | 1 (in flower-bud). | |
| June | 15 | 1 | |
| July | 12 | .. | |
| August | 100 | 2 | |

(2) STANDING COTTON PLANTS

The general cotton area is completely free from living cotton plants within a couple of months after the harvest of *kupas*. A few of these plants, however, could be found

growing along the borders and hedges of the irrigated plots in several places, and *Earias* larvae could be traced in flower-buds and bolls collected from these plants.

TABLE XI

Earias larvae on flower-buds and bolls collected from cotton plants growing in irrigated plots

| Month of examination | Nagardevala | | Kusumba | | Nimbhora | |
|----------------------|-----------------------------|----------------------|--------------------------------|----------------------|-----------------------------|----------------------|
| | Total flower-buds and bolls | <i>Earias</i> larvae | Total flower-buds and bolls | <i>Earias</i> larvae | Total flower-buds and bolls | <i>Earias</i> larvae |
| January | 414 | 83 | 200 | 27 | 211 | 36 |
| February | 85 | 5 | 153 | 17 | 314 | 22 |
| March | 196 | 27 | 134 | 6 | 258 | 16 |
| April | 115 | 26 | .. | .. | 30 | .. |
| May | 45 | Nil | Forms not available hereafter. | | | |
| June | 39 | 4 | .. | .. | .. | .. |
| July | 41 | .. | .. | .. | .. | .. |
| August | 2,000 | 1 | .. | .. | .. | .. |

These forms were available at all the places till the end of March but it was only at Nagardevala that they could be collected till the commencement of the next crop.

Bhendi (Hibiscus esculentus)

The monsoon crop of *bhendi* is over by the month of October or November. A few stray plants of this crop, however, continue to grow in the irrigated plots. *Earias* larvae can be collected from them till February.

TABLE XII

Earias larvae in bhendi pods from the remnant of the monsoon crop of bhendi

| Month | JALGAON | | NAGARDEVALA | |
|-------------------------|---------------|----------------------|---------------|----------------------|
| | Pods examined | <i>Earias</i> larvae | Pods examined | <i>Earias</i> larvae |
| January 1930 | 10 | 9 | 20 | 20 |
| February 1930 | 44 | 14 | 4 | 4 |

A second crop of *bhendi* is generally sown in March or April in the irrigated plots in several villages, and these plants are helpful for supplying food to the Spotted Boll-worms after March till the monsoon.

TABLE XIII

Earias larvae in bhendi pods from the summer crop 1930

| Month of examination | JALGAON | | NAGARDEVALA | | KUSUMBA | | NIMBHORA | |
|----------------------|---------|--------|-------------|--------|---------|--------|----------|--------|
| | Pods | larvae | Pods | Larvae | Pods | Larvae | Pods | Larvae |
| March | 88 | 15 | .. | .. | .. | .. | .. | .. |
| April | 83 | 68 | 38 | 21 | 25 | 28 | .. | .. |
| May | 73 | 56 | 26 | 15 | .. | 2* | .. | .. |
| June | 22 | 14 | 25 | 15 | .. | 3* | 10 | 2 |
| July | .. | .. | 25 | 22 | 15 | 1 | 30 | 2 |

* In shoots.

The pods from the plots mentioned above are available in the local market after the middle of May and *Earias* larvae can often be collected from them.

TABLE XIV

Earias larvae in bhendi pods, secured from the Jalgaon market from the local crop of bhendi, 1930

| Weeks | No. of pods | <i>Earias</i> larvae |
|---------------------------|-------------|----------------------|
| 22nd May 1930 | 65 | 5 |
| 30th May 1930 | 82 | 6 |
| 7th June 1930 | 56 | 2 |
| 15th June 1930 | 65 | 5 |
| 21st June 1930 | 53 | 3 |
| 30th June 1930 | 45 | 13 |
| 7th July 1930 | 37 | 5 |
| 14th July 1930 | 59 | 2 |
| 21st July 1930 | 42 | .. |
| 28th July 1930 | 46 | 3 |
| 2nd August 1930 | 71 | 2 |

Before this local supply of pods is available they are imported in Jalgaon market from Nasik district, during the months of April and May.

TABLE XV

Earias larvae from market pods of bhendi from Nasik district

| Dates | No. of pods | No. of <i>Earias</i> larvae |
|---------------------------|-------------|-----------------------------|
| 14th April 1930 | 41 | 27 |
| 16th April 1930 | 52 | 33 |
| 26th April 1930 | 24 | 12 |
| 11th May 1930 | 32 | 10 |
| 17th May 1930 | 65 | 22 |

POSSIBILITIES OF AESTIVATION OF THE SPOTTED BOLL-WORMS IN KHANDESH

There has been already enough evidence to show that in Khandesh also the Spotted Boll-worms are active throughout the year and their larvae can be collected from the irrigated plots even during the summer months. We wanted, however, to clear up the question about the likelihood of their aestivating in the cotton areas of Khandesh, as pupae in the soil or resting as moths. Small blocks of cotton area were, therefore, selected, when the cotton plants were removed after the harvest of *kapas*, and the soil from them was thoroughly examined to a depth of three to five feet (according to the depth of the cracks) in order to locate the pupae in the soil.

TABLE XVI

Results of soil examination for searching out Earias pupae

| Time of soil examination | Area examined in square feet | Cocoons found | Live pupae |
|--|------------------------------|---------------|------------|
| Two days after the plants were removed . | 15×10 | 32 | <i>nil</i> |
| 22 days after the plants were removed . | 15×5 | 66 | <i>nil</i> |
| 30 days after the plants were removed . | 15×10 | 23 | <i>nil</i> |
| 45 days after the plants were removed . | 10×10 | 11 | <i>nil</i> |

Bait pans with cotton and sesamum cakes, which usually attracted a large number of *Earias* moths (Appendix I, Tables X, XI, XXI and XXII) were continuously exposed in the cotton areas at Jalgaon, during summer after the cotton plants were removed. Not a single moth, however, could be collected from these pans during the period mentioned above.

In Khandesh also, therefore, we could not get any indications about the possibility of aestivation of the Spotted Boll-worms, during the period between two cotton-growing seasons.

SUMMARY

The cotton crop in Khandesh escapes serious damage from the Spotted Boll-worms if it matures very early, before this pest gets an opportunity to increase in numbers. However, it cannot be considered to be a minor pest in years when the crop is delayed. In such seasons an increase of about 25 per cent in the yield of *kapas* might be expected if this pest can be controlled.

No indications were obtained of the possibilities of aestivation of this pest in Khandesh; and it was found to be active throughout the year.

The cotton areas are cleared up before the end of March and the stumps of cotton do not sprout, as they do in Gujarat; there is, therefore, no food available for these larvae in the general cotton areas.

Irrigated plots around the wells are found to be chiefly responsible for helping the carry-over of these insects from season to season, and the presence of *bhendi* plants in

these plots during the months of April, May and June appear to be most harmful, though stray plants of cotton and *Abutilon* weeds are also found to harbour *Earias* larvae in these plots to a certain extent. (Plate XXIV).

It appears that Spotted Boll-worms will no longer be a troublesome pest in Khandesh if the irrigated plots are kept free from plants, which afford food to these insects during the interval between two cotton-growing seasons.

It would be rather difficult to dissuade the cultivators from growing *bhendi* in these plots in summer, because it is considered to be a profitable crop.


Measures will also be necessary to prevent the importation of *bhendi* pods from other Districts to avoid fresh infection.

A few trials were made at Jalgaon, during 1930, of using *bhendi* as a trap crop. The results have been discussed in pages 45-48.

TABLE XVII

Weekly rainfall at Jalgaon. (Records from Government Farm)

| Months | RAINFALL IN INCHES | | |
|---------------------|--------------------|-------|-------|
| | Weeks | 1929 | 1930 |
| June | first . . . | 0.90 | .. |
| | second . . . | .. | 0.45 |
| | third . . . | 6.23 | 4.22 |
| | fourth . . . | 2.54 | 1.62 |
| July | first . . . | 2.59 | 4.50 |
| | second . . . | 2.03 | .. |
| | third . . . | 0.58 | 1.68 |
| | fourth . . . | 0.16 | 0.53 |
| August | first . . . | 0.60 | 3.02 |
| | second . . . | 0.13 | 0.34 |
| | third . . . | .. | .. |
| | fourth . . . | 0.55 | 0.35 |
| September | first . . . | 0.36 | 2.79 |
| | second . . . | .. | 15.92 |
| | third . . . | 0.90 | 0.11 |
| | fourth . . . | 0.31 | .. |
| October | first . . . | 0.86 | .. |
| | second . . . | .. | .. |
| | third . . . | .. | .. |
| | fourth . . . | .. | 0.18 |
| TOTAL | .. | 18.74 | 36.31 |



Showing sources of food for the Spotted Boll-worms during the period between two cotton-growing seasons in Khandesh

APPENDIX V

PHYCITA INFUSELLA MYER. (COTTON SHOOT-ROLLER) AT SURAT

This is another pest which is well established on cotton in South Gujarat. It appeared at Surat every year during the course of about eight years of the investigations on Spotted Boll-worms. A certain amount of information was incidentally collected about this pest.

The caterpillars of the cotton shoot-roller bind together the tender leaves of the cotton shoots and destroy the tiny growing buds. A part of one of the leaves is rolled up and the larva hides itself inside that. As a result of the damage of this pest, the further growth of the shoots is arrested. Periodical observations during 1925-26 showed that 62 per cent of the shoots of the main stems were damaged by these larvae.

POPULATION OF *Phycita* LARVAE ON COTTON CROP

The incidence of this pest at Surat during four years commencing from 1927-28 can be seen from tables II to V of this Appendix. The attack of these larvae on the cotton plants begins when the seedlings are from 4 to 6 inches in height. Their population generally increases till the end of October or the beginning of November, when it begins to decline until it completely disappears from the cotton plants, or becomes almost insignificant, in March.

HIBERNATION

Of the four tables mentioned above, tables II and III show the classification of the larvae into three arbitrary classes, viz., small, medium and big. This classification will show that during 1928, after the end of December, small-sized larvae could not be found, and again during 1929, the population of the small-sized larvae declined appreciably from the middle of November; so that after the middle of December, most of the larvae collected were of the big size. This indicated that these larvae hibernate either in November or December, and thereafter fresh oviposition practically ceases. These caterpillars roll up one or two leaves, which dry up, and the resting stage is passed in these leaves. As the season advances, these dry leaves drop away from the plants, along with the resting larvae, and therefore their number on the plants declines until they can no longer be found on the cotton plants during February and March. From the beginning of April fresh small-sized larvae begin to appear again on the cotton plants, and this, therefore, seems to be their time for coming out from hibernation.

Phycita LARVAE ON STANDING COTTON PLANTS AFTER HARVEST

By the time the pickings of *kapas* are over, the plants begin to put up fresh growth. It is, therefore, these new shoots, which are attacked by these larvae when they reappear

in the cotton fields. Table I will show that the population of these larvae on the standing cotton sticks increases very rapidly during April, May and June.

Phycita LARVAE ON SPROUTS FROM STUMPS

Periodical examination of the fresh shoots growing from the stumps of the cotton plants, which are usually removed by hacking with a light pick-axe, showed that *Phycita* larvae could easily be collected from these sprouts during May, June, July and August in 1928 and 1929. (Table I at the end of this appendix.)

It is thus seen that the caterpillars of *Phycita infusella* Meyr. feed on the shoots of the cotton plants from August till November or December, when they enter into a resting stage. They emerge from this stage at the commencement of April and sprouts from standing cotton plants and stumps constitute their source of food from April till the seedlings of the next crop are ready for their attack.

It appears that the large carry-over of this pest to the next crop can effectively be stopped if the cotton plants are uprooted immediately after the pickings are over, so that there are no standing cotton plants or their stumps to provide food for these worms. The clean-up measures against the Spotted Boll-worms (discussed on pages 91 and 92) are, therefore, very likely to be beneficial from the point of view of controlling this pest also.

TABLE I

Cotton shoot-rollers (Phycita infusella Meyr.) on sprouts from stumps of cotton, Surat, 1928-29

| Date of examination | Sprouts from No. of stumps | No. of larvae |
|--------------------------|----------------------------------|---------------|
| 10th May 1928 | 75 | 8 |
| 19th May 1928 | 75 | 19 |
| 25th May 1928 | 75 | 75 |
| 1st June 1928 | 100 | 5 |
| 7th June 1928 | 200 | 57 |
| 15th June 1928 | 200 | 22 |
| 22nd June 1928 | 200 | 33 |
| 30th June 1928 | 200 | 49 |

TABLE I—*contd.*

Cotton shoot-rollers (Phycita infusella Meyr.) on sprouts from stumps of cotton, Surat, 1928-29—contd.

| Date of examination | Sprouts from No. of stumps | No. of larvae |
|----------------------------|----------------------------------|---------------|
| 9th July 1928 | 200 | 17 |
| 20th July 1928 | 200 | 8 |
| 31st July 1928 | 200 | 5 |
| 11th August 1928 | 200 | 10 |
| 15th August 1928 | 200 | 23 |
| 23rd August 1928 | 200 | 69 |
| 31st August 1928 | 200 | 53 |
| 4th May 1929 | 100 | 6 |
| 11th May 1929 | 100 | 10 |
| 18th May 1929 | 100 | 15 |
| 25th May 1929 | 200 | 17 |
| 1st June 1929 | 200 | 32 |
| 12th June 1929 | 200 | 26 |
| 22nd June 1929 | 200 | 4 |
| 27th June 1929 | 200 | 8 |
| 6th July 1929 | 200 | 5 |
| 18th July 1929 | 200 | 7 |
| 25th July 1929 | 200 | 7 |
| 3rd August 1929 | 200 | 19 |
| 10th August 1929 | 200 | 24 |
| 17th August 1929 | 150 | 16 |
| 25th August 1929 | 100 | 12 |

TABLE II

Larval population of Phycita infusella Meyr. on cotton plants at Surat in 1927-28

| Week ending dates | No. of plants | No. of larvae | Week ending dates | No. of plants | No. of larvae |
|-----------------------|---------------|---------------|----------------------|---------------|---------------|
| 1 | 2 | 3 | 1 | 2 | 3 |
| 17th September 1927 . | 25 | 9 | 3rd March 1928 . . | 25 | nil |
| 24th September 1927 . | 25 | 15 | 10th March 1928 . . | 25 | 1 |
| 1st October 1927 . | 25 | 41 | 17th March 1928 . . | 25 | 1 |
| 8th October 1927 . | 25 | 26 | 24th March 1928 . . | 25 | nil |
| 15th October 1927 . | 25 | 53 | 31st March 1928 . . | 50 | 10 |
| 22nd October 1927 . | 25 | 62 | 7th April 1928 . . | 50 | 17 |
| 29th October 1927 . . | 25 | 65 | 14th April 1928 . . | 50 | 18 |
| 5th November 1927 . | 25 | 56 | 21st April 1928 . . | 50 | 31 |
| 12th November 1927 . | 25 | 57 | 28th April 1928 . . | 50 | 16 |
| 19th November 1927 . | 25 | 64 | 5th May 1928 . . | 50 | 27 |
| 26th November 1927 . | 25 | 58 | 12th May 1928 . . | 50 | 33 |
| 3rd December 1927 . | 25 | 28 | 19th May 1928 . . | 50 | 82 |
| 10th December 1927 . | 25 | 27 | 26th May 1928 . . | 50 | 44 |
| 17th December 1927 . | 25 | 28 | 2nd June 1928 . . | 50 | 54 |
| 24th December 1927 . | 25 | 25 | 9th June 1928 . . | 50 | 102 |
| 31st December 1927 . | 25 | 25 | 16th June 1928 . . | 50 | 142 |
| 7th January 1928 . . | 25 | 20 | 23rd June 1928 . . | 50 | 127 |
| 14th January 1928 . . | 25 | 14 | 30th June 1928 . . | 50 | 99 |
| 21st January 1928 . . | 25 | 9 | 7th July 1928 . . | 50 | 129 |
| 28th January 1928 . . | 25 | 10 | 14th July 1928 . . | 50 | 78 |
| 4th February 1928 . | 25 | 12 | 21st July 1928 . . | 50 | 70 |
| 11th February 1928 . | 25 | 6 | 28th July 1928 . . | 50 | 68 |
| 18th February 1928 . | 25 | 3 | 4th August 1928 . . | 50 | 66 |
| 25th February 1928 . | 25 | 1 | 11th August 1928 . . | 50 | 26 |

TABLE III

Larval population of Phycita infusella Meyr. on cotton plants, with classification of the sizes of larvae, at Surat, 1928-29

| Week ending dates | No. of plants | SIZES OF LARVAE | | | Total larvae |
|-------------------------------|---------------|----------------------|--------|-----|--------------|
| | | Small | Medium | Big | |
| 1 | 2 | 3 | 4 | 5 | 6 |
| 28th July 1928 | 1,350 | 0 | 1 | 0 | 1 |
| 4th August 1928 | 1,000 | .. | .. | .. | .. |
| 11th August 1928 | 1,000 | 3 | 6 | 1 | 10 |
| 18th August 1928 | 1,000 | 6 | 16 | 4 | 26 |
| 25th August 1928 | 750 | 7 | 10 | 13 | 30 |
| 1st September 1928 | 750 | 11 | 23 | 29 | 63 |
| 8th September 1928 | 250 | 2 | 8 | 4 | 14 |
| 15th September 1928 | 125 | 1 | 4 | 5 | 10 |
| 22nd September 1928 | | Plants not examined. | | | |
| 29th September 1928 | 100 | 7 | 12 | 3 | 22 |
| 6th October 1928 | 50 | 5 | 8 | 3 | 16 |
| 13th October 1928 | 25 | 5 | 7 | 2 | 14 |
| 20th October 1928 | 25 | 5 | 3 | 2 | 10 |
| 27th October 1928 | 25 | 6 | 5 | 1 | 12 |
| 3rd November 1928 | 25 | 7 | 11 | 1 | 19 |
| 10th November 1928 | 25 | 3 | 10 | 1 | 14 |
| 17th November 1928 | 25 | 4 | 7 | 2 | 13 |
| 24th November 1928 | 25 | 3 | 6 | 4 | 13 |
| 1st December 1928 | 25 | Sizes not noted | | | 3 |
| 8th December 1928 | 25 | Sizes not noted | | | 4 |
| 15th December 1928 | 25 | 1 | 1 | 6 | 8 |
| 22nd December 1928 | 25 | 1 | 1 | 3 | 5 |
| 29th December 1928 | 25 | 1 | 3 | 1 | 5 |

TABLE III—contd

Larval population of Phycita infusella Meyr. on cotton plants with classification of the sizes of larvae, at Surat, 1928-29—contd

| Week ending dates | No. of plants | SIZES OF LARVAE | | | Total larvae |
|------------------------------|---------------|-----------------|--------|-----|--------------|
| | | Small | Medium | Big | |
| 1 | 2 | 3 | 4 | 5 | 6 |
| 5th January 1929 | 25 | .. | 1 | 4 | 5 |
| 12th January 1929 | 25 | .. | 1 | 1 | 2 |
| 19th January 1929 | 25 | .. | 1 | 2 | 3 |
| 26th January 1929 | 25 | .. | 1 | .. | 1 |
| 2nd February 1929 | 25 | .. | .. | 2 | 2 |
| 9th February 1929 | 25 | .. | .. | 1 | 1 |
| 16th February 1929 | 25 | .. | .. | 1 | 1 |
| 23rd February 1929 | 25 | .. | .. | .. | .. |
| 2nd March 1929 | 25 | .. | .. | .. | .. |
| 9th March 1929 | 25 | .. | .. | .. | .. |
| 16th March 1929 | 25 | .. | .. | .. | .. |
| 23rd March 1929 | 25 | .. | .. | .. | .. |
| 30th March 1929 | 50 | .. | .. | .. | .. |
| Cotton pickings were over. | | | | | |
| 6th April 1929 | 50 | 5 | 1 | .. | 6 |
| 13th April 1929 | 50 | 4 | 4 | 1 | 9 |
| 20th April 1929 | 50 | 4 | 1 | 1 | 6 |
| 27th April 1929 | 50 | 10 | 5 | 1 | 16 |
| 2nd May 1929 | 50 | 27 | 11 | .. | 38 |
| 9th May 1929 | 50 | 25 | 17 | 10 | 52 |
| 16th May 1929 | 50 | 18 | 21 | 14 | 53 |
| 23rd May 1929 | 50 | 13 | 11 | 18 | 42 |
| 30th May 1929 | 50 | 28 | 13 | 19 | 60 |

TABLE IV

Larval population of Phycita infusella Meyr. on cotton plants at Surat

1929-30

| Week ending dates | No. of plants examined | SIZES OF LARVAE | | | Total larvae |
|-----------------------------|------------------------|-----------------|--------|-----|--------------|
| | | Small | Medium | Big | |
| 1 | 2 | 3 | 4 | 5 | 6 |
| 3rd August 1929 | 1,300 | .. | .. | .. | . |
| 10th August 1929 | 887 | 14 | 4 | .. | 18 |
| 17th August 1929 | 1,162 | 7 | 9 | 7 | 23 |
| 24th August 1929 | 396 | 4 | 4 | 7 | 15 |
| 31st August 1929 | 860 | 9 | 7 | 8 | 24 |
| 7th September 1929 | 750 | 2 | 2 | 4 | 8 |
| 14th September 1929 | 500 | 11 | 4 | 7 | 25 |
| 21st September 1929 | 250 | 8 | 6 | 4 | 18 |
| 28th September 1929 | 125 | 19 | 10 | 17 | 46 |
| 2nd October 1929 | 50 | 9 | 12 | 8 | 29 |
| 11th October 1929 | 25 | 22 | 9 | 6 | 37 |
| 16th October 1929 | 25 | 38 | 26 | 12 | 76 |
| 23rd October 1929 | 25 | 50 | 35 | 10 | 95 |
| 30th October 1929 | 25 | 54 | 38 | 19 | 111 |
| 8th November 1929 | 25 | 52 | 19 | 37 | 138 |
| 14th November 1929 | 25 | 21 | 18 | 60 | 136 |
| 21st November 1929 | 25 | 6 | 36 | 67 | 109 |
| 28th November 1929 | 25 | 3 | 25 | 75 | 103 |
| 5th December 1929 | 25 | 1 | 15 | 52 | 68 |
| 12th December 1929 | 25 | .. | 5 | 57 | 62 |

TABLE IV—*contd**Larval population of Phycita infusella* Meyr. on cotton plants at Surat—*contd*1929-30—*contd*

| Week ending dates | No. of plants examined | SIZES OF LARVAE. | | | Total larvae |
|------------------------------|------------------------|------------------|--------|-----|--------------|
| | | Small | Medium | Big | |
| 1 | 2 | 3 | 4 | 5 | 6 |
| 19th December 1929 | 25 | 1 | .. | 42 | 43 |
| 26th December 1929 | 25 | .. | .. | 32 | 32 |
| 2nd January 1930 | 25 | .. | .. | 39 | 39 |
| 9th January 1930 | 25 | 1 | .. | 14 | 15 |
| 16th January 1930 | 25 | .. | .. | 7 | 7 |
| 23rd January 1930 | 25 | .. | .. | 4 | 4 |
| 30th January 1930 | 25 | .. | .. | 3 | 3 |
| 6th February 1930 | 25 | .. | .. | 2 | 2 |
| 13th February 1930 | 25 | .. | .. | .. | .. |
| 20th February 1930 | 25 | .. | .. | .. | .. |
| 27th February 1930 | 25 | .. | .. | .. | .. |
| 6th March 1930 | 25 | .. | .. | .. | .. |
| 13th March 1930 | 25 | .. | .. | .. | .. |
| 20th March 1930 | 25 | .. | .. | .. | .. |
| 27th March 1930 | 25 | .. | .. | .. | .. |

Cotton pickings were over.

| | | | | | |
|---------------------------|----|---|----|----|---|
| 5th April 1930 | 50 | 1 | .. | .. | 1 |
| 11th April 1930 | 50 | 2 | .. | .. | 2 |
| 16th April 1930 | 50 | 2 | 1 | 1 | 4 |

TABLE V

Larval population of Phycita infusella on cotton plants at Surat

1930-31

| Weeks of examination | No. of plants examined | Total larvae |
|-------------------------------|------------------------|--------------|
| 13th August 1930 | 1,000 | .. |
| 21st August 1930 | 1,000 | 10 |
| 26th August 1930 | 1,000 | 21 |
| 3rd September 1930 | 500 | 12 |
| 11th September 1930 | 200 | 9 |
| 18th September 1930 | 100 | 3 |
| 25th September 1930 | 50 | 11 |
| 3rd October 1930 | 25 | 10 |
| 10th October 1930 | 25 | 11 |
| 17th October 1930 | 25 | 23 |
| 24th October 1930 | 25 | 10 |
| 31st October 1930 | 25 | 14 |
| 7th November 1930 | 25 | 9 |
| 14th November 1930 | 25 | 6 |
| 21st November 1930 | 25 | 8 |
| 28th November 1930 | 25 | 4 |
| 5th December 1930 | 25 | 2 |
| 12th December 1930 | 25 | 5 |
| 19th December 1930 | 25 | 5 |
| 26th December 1930 | 25 | 3 |
| 2nd January 1931 | 25 | 5 |
| 9th January 1931 | 25 | 3 |
| 16th January 1931 | 25 | 1 |
| 23rd January 1931 | 25 | .. |
| 30th January 1931 | 25 | .. |

APPENDIX VI.

PINK BOLL-WORM (*PLATYEDRA GOSSYPIELLA* SAUND.) IN SOUTH GUJARAT

During the period of investigation on the Spotted Boll-worms at Surat, it was possible to make a few observations about the Pink Boll-worm also.

Twenty-five cotton plants were examined every week during the cotton-growing season, for recording the population of the Spotted Boll-worms (page No. 26). When these plants were examined, records of the Pink Boll-worm larvae were maintained. The results of these observations, made at Surat for seven years from 1924 to 1931, are included in Tables VII to XIII of this appendix. Table XIV shows the population of the Pink Boll-worm larvae during 1927-28 on the cotton crop on the Government Farm at Broach.

It will be seen from these tables that Pink Boll-worm is a well established pest in South Gujarat and that its earliest presence in the cotton fields can be noticed during October. There are no bolls on the cotton plants at that time, and hence their earliest appearance happens to be in flower-buds. In one case, a Pink Boll-worm larva was found in the vegetative shoot of the cotton plant. This was, however, an exceptional case. Though there are some instances of the presence of these larvae on the cotton plants during October, their number does not increase rapidly until well developed bolls are available on the cotton plants during December or January.

The figures further indicate that Pink Boll-worm cannot be normally considered to be a very serious pest in South Gujarat. This will probably be more clearly seen from statement No. XV of this appendix, which gives the number of green bolls examined every week from 25 cotton plants, and the number of Pink Boll-worm larvae calculated for 100 green bolls.

In four out of seven years, the population practically never increased to a level of even 10 larvae for every 100 green bolls. It was only during 1927-28 and 1928-29 it rose to about 25 larvae per 100 bolls, and the damage due to this pest to the cotton crop at Surat in 1928-29 might be considered to be the highest during all these years.

The population of the Spotted Boll-worms declines to a very low level by the end of December or the beginning of January, and it is about this time that the Pink Boll-worm larvae increase in numbers. This happens to be the period of the development of bolls, and hence the boll-worm damage which can be seen in the open bolls at Surat can almost completely be attributed to the Pink Boll-worms. In order to estimate this damage to the grown-up bolls more than 100 plants were kept under observation during 1928-29, 1929-30 and 1930-31.

At the time of collecting the *kapas*, all the bolls were carefully examined, and the *kapas* from sound and damaged locks was collected separately. The number of damaged and useless locks was also recorded.

TABLE I

Proportion of locks damaged by boll-worms, out of the total locks collected per plant during the above three years at Surat

| Year | No. of plants observed | Total No. of locks picked | No. of sound locks | Damaged locks showing boll-worm injury | Useless locks showing boll-worm injury | Total locks with boll-worm injury | Percentage of locks with boll-worm injury |
|-------------|------------------------|---------------------------|--------------------|--|--|-----------------------------------|---|
| 1928-29 . . | 104 | 202 | 144 | 38 | 7 | 45 | 22 |
| 1929-30 . . | 112 | 171 | 156 | 9 | 3 | 12 | 7 |
| 1930-31 . . | 116 | 212 | 191 | 18 | 2 | 20 | 9 |

TABLE II

Proportion of kapas from damaged locks out of the total saleable kapas, per plant, during the above three years at Surat

| Year | No. of plants observed | Total saleable kapas in grms. | Kapas from sound locks in grms. | Kapas from locks damaged by boll-worm in grms. | Percentage of damaged kapas |
|-----------------|------------------------|-------------------------------|---------------------------------|--|-----------------------------|
| 1928-29 | 104 | 130.6 | 105.5 | 25.1 | 19 |
| 1929-30 | 112 | 113.2 | 107.9 | 5.3 | 5 |
| 1930-31 | 116 | 150.1 | 140.7 | 9.4 | 6 |

From these statements it can be seen that during 1928-29, which appeared to be the worst Pink Boll-worm season at Surat, a fifth of the *kapas* came from the damaged locks. During the other two years the proportion of damaged *kapas* was only 5 and 6 per cent respectively.

RESTING STAGE OF PINK BOLL-WORM LARVAE

In the Punjab as well as in Egypt, the caterpillars of Pink Boll-worm pass the winter in hibernation [Bindra 1928 and Gough 1919]. In the Punjab the larvae enter into the resting stage during the months of October, November and December. These larvae are known as the long-cycle larvae, to differentiate them from the short-cycle larvae which pupate as soon as they are fully fed, and emerge as moths after a normal period

of pupation. The emergence from the long-cycle larvae commences by the beginning of April. In South Gujarat, however, the period of boll development generally falls in the months of December and January, which are the coldest months; and it is during these months that the population of the caterpillars of Pink Boll-worm increases rapidly. In order, therefore, to ascertain the presence or absence of the long-cycle larvae in South Gujarat, observations were undertaken on (1) green bolls, (2) the bolls rejected on the cotton sticks after harvest and (3) the *kapas* and seed.

OBSERVATIONS ON GREEN BOLLS

In the first week of January 1926 (about 4 weeks before the 1st picking of *kapas*), 800 fully developed green bolls were collected at random, and were allowed to dry in open cages; because, if kept in closed jars, they began to rot within a short time. They were afterwards kept in 3 receptacles for noting the emergence of moths. Before the middle of March, 30 Pink Boll-worm moths emerged from these bolls, and then there was no emergence till the middle of April.

During the last fortnight of April, all these bolls were carefully examined and 63 caterpillars of Pink Boll-worm were found in them as mentioned below:—

30 larvae were found in flimsy lint cocoons.

25 larvae were found in single seeds.

7 larvae were found in double seeds.

1 larva was found free in the cage.

—

63 Total.

Another lot of 600 grown-up green bolls was collected on the 4th January 1926 and these bolls, after drying, were kept in earthen pots, which were covered with thin muslin cloth on the 4th February 1926. Emergence of moths from these bolls was recorded.

TABLE III

| Time of emergence | No. of Pink Boll-worm moths |
|---|-----------------------------|
| Before the 15th of March 1926 | 29 |
| In July 1926 | 27 |
| In August 1926 | 9 |
| In September 1926 | 3 |
| In October 1926 | 1 |
| In November 1926 | .. |
| In December 1926 | .. |
| In January 1927 | 1 |

These observations in 1926 showed that long-cycle larvae did exist in South Gujarat also.

During the next season, 1,300 fully developed green bolls were collected when the second picking was in progress. They were kept in four cages, after drying, on the 29th of April 1927 for noting the emergence of the Pink Boll-worm moths from them.

TABLE IV

| Month of emergence | No. of Pink Boll-worm moths |
|--------------------------|-----------------------------|
| May 1927 | 29 |
| June 1927 | 43 |
| July 1927 | 259 |
| August 1927 | 84 |
| September 1927 | 3 |
| October 1927 | 1 |

Another lot of 300 fully grown green bolls was collected on 8th April 1927, and was kept under observation as in the above case till the 21st July 1927, when all the bolls were removed for examination. Before this date, 49 Pink Boll-worm moths had emerged from these bolls : 11 in May, 6 in June, and 32 in July. Examination of these bolls disclosed 16 Pink Boll-worm larvae and 7 pupae as stated below :—

5 larvae in lint cocoons.
 3 larvae in single seeds.
 4 larvae in double seeds.
 3 larvae in treble seeds.
 1 larva in group of seeds.

—
 16 Total larvae.

2 pupae in lint cocoons.
 1 pupa in single seed.
 1 pupa in double seed.
 3 pupae free in the lint.

—
 7 Total pupae,

These observations in 1927 show that it was not the cold which stimulated the larvae to display the long-cycle response because a large number of the long-cycle larvae were obtainable from the green bolls collected during April.

During 1927-28, samples of full grown bolls (green) each consisting of 250 bolls, were collected both at Surat and at Broach during the boll-development period. Four samples were collected at Broach and two at Surat. The dates of collecting these samples and the emergence of Pink Boll-worm moths from these bolls are shown in the table below.

TABLE V

Emergence of Pink Boll-worm moths from green bolls collected at intervals at Broach and Surat

| Time of emergence | Broach sample No. I collected on 27th December 1927 | Broach sample No. II collected on 14th January 1928 | Broach sample No. III collected on 30th January 1928 | Broach sample No. IV collected on 28th February 1928 | Surat sample No. I collected on 16th January 1928 | Surat sample No. II collected on 23rd February 1928 |
|------------------------|---|---|--|--|---|---|
| Before 30th April 1928 | 85 | 20 | .. | 5 | 11 | .. |
| In June 1928 | .. | .. | 1 | 2 | 1 | 1 |
| In July 1928 | 8 | 1 | 7 | 24 | .. | 17 |
| In August 1928 | 3 | 3 | .. | 9 | .. | 6 |
| In September 1928 | .. | .. | .. | 2 | .. | 1 |

The bolls collected during February both at Surat and Broach have given the largest number of long-cycle larvae. Probably the number of the resting caterpillars would have been still greater if the samples had been collected even later, i.e., in March or April.

OBSERVATIONS ON DRY BOLLS REJECTED ON THE PLANTS

There are always some damaged bolls which are left on the plants after the pickings are over. A few of these drop down to the soil, and the rest are carried along with the cotton sticks, which are stored in heaps. On the 27th of May 1926, 501 such dry bolls

were collected from 194 cotton plants from a stack. Examination of these bolls showed the presence of 44 Pink Boll-worm larvae in them. Fifteen of these larvae were found in single seeds, 27 were found between the compact locks of lint and the carpels and two were found parasitised.

In the next season, two lots of similar dry bolls were collected on the 27th of May 1927. The first lot consisting of 247 bolls was collected from 105 plants, and the second lot of 302 bolls was collected from 109 plants.

The first lot was examined after the 15th of July 1927, and one pupa of Pink Boll-worm and five full grown Pink Boll-worm caterpillars were obtained from these 247 bolls. The bolls from the second lot were kept under observation for noting the emergence of Pink Boll-worm moths. Only one moth emerged from these bolls on the 16th of August 1927.

There is thus a possibility of a small number of the long-cycle larvae resting in the dry bolls left out on the cotton plants after the harvest of *kapas*.

OBSERVATIONS ON *kapas* AND SEED

Examination of kapas. Two pounds of *kapas* collected from the general lot at Surat was examined from 8th to 23rd June and 12 Pink Boll-worm caterpillars were found in it.

7 larvae in single seeds.

3 larvae in lint cocoons.

2 larvae free in the lint.

—

12 Total.

Another sample of two lbs. of *kapas* from the second picking was examined from 26th June to 5th July and only one Pink Boll-worm larva was found in a single seed.

Besides this, two samples of *kapas*, of one pound each from the first and the second pickings respectively, were kept in jars for noting the emergence of Pink Boll-worm moths. From one pound of *kapas* of the first picking, only one Pink Boll-worm moth emerged on the 11th of August 1928; and from the sample from the second, one moth emerged in the 3rd week of July and another in the last week of July.

Examination of seed. Small quantities of cotton seed from the *kapas* of the first and second pickings were examined by splitting open every seed during July and August 1928. The cotton seed was obtained by ginning the *kapas* with hand gins. All the seeds which showed damage caused by the boll-worms, were kept separate and the total quantity was weighed at the end of each examination. The number of Pink Boll-worm larvae found in the seeds can be seen from the next table. All the caterpillars were found in single seeds,

TABLE VI

Examination of cotton seed

1928

| Quantity of seed examined | No. of Pink boll-worm larvae found | Quantity of damaged seed | Time of examination |
|-----------------------------------|------------------------------------|--------------------------|---------------------|
| | | Tolas | |
| 2 lbs. from 1st Picking | 5 | 7 | July. |
| 2 lbs. from 1st Picking | 1 | 7.75 | August. |
| 1 lb. from 2nd picking | .. | 2.5 | July end. |
| 1½ lb. from 2nd picking | 3 | 5.25 | August. |

It is thus seen that the *kapas* as well as the cotton seed both have a small number of resting Pink Boll-worm caterpillars in them.

Some of the baits which were tried for collecting the moths of Spotted Boll-worms attracted the moths of Pink Boll-worm also as can be seen from tables VIII to XI of appendix No II.

CONCLUSIONS

Pink boll-worm appears every year in the cotton fields in South Gujarat. It increases in number during December and January, but normally it does not cause a very serious damage to the cotton crop. About twenty-five larvae for every 100 bolls were recorded only in two out of the seven seasons during the worst period of the attack.

During 1928-29, which was the worst Pink Boll-worm year, 19 per cent of the *kapas* was picked from damaged locks, whereas during the next two seasons the proportion of the damaged *kapas* was 6 and 5 per cent respectively.

The long-cycle larvae are found in this tract, and this pest is carried over from one season to another by these larvae resting in *kapas*, in cotton seed and in the dry bolls rejected in the cotton fields. The exact responsibility of each of these in the carry-over of the pest could not, however, be worked out.

It was further seen that the maximum emergence of the Pink Boll-worm moths from the resting larvae takes place in July. The emergence declines in August, and after

the end of August, there is only a very occasional emergence in September and October. During July and August there is practically no food available for these worms on the cotton crop, as it is sown at the end of June or at the beginning of July. The plants are consequently very small, and even flower-buds are generally absent. Probably from the stray emergence of moths during September and October, a small infestation continues on the flower-buds and young bolls until developed bolls are available. These developed bolls are usually available in December and the pest has, therefore, to carry on its mere existence till that time. From December onwards, the population begins to rise rapidly.

The long interval between the period of the emergence of the moths from the resting stage and the period of the availability of the developed bolls (which is the principal food of these larvae) seems to be the chief factor, which prevents this pest from assuming alarming proportions in this important tract of cotton.

TABLE VII

Weekly population of Pink Boll-worm larvae on 25 cotton plants at Surat (1924-25)

| Week ending | No. of flower-buds | Larvae in flower-buds | No. of bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|------------------------|--------------------|-----------------------|--------------|-----------------|-----------------------------|
| 12th December 1924 . . | 2,260 | 7 | 367 | 7 | 14 |
| 19th December 1924 . . | 2,095 | 3 | 644 | 11 | 14 |
| 26th December 1924 . . | 1,652 | 8 | 876 | 8 | 16 |
| 2nd January 1925 . . | 971 | nil. | 1,005 | 59 | 59 |
| 9th January 1925 . . | 461 | 1 | 1,000 | 29 | 30 |
| 16th January 1925 . . | 205 | nil. | 1,110 | 46 | 46 |
| 23rd January 1925 . . | 172 | nil. | 865 | 14 | 14 |
| 30th January 1925 . . | 102 | nil. | 979 | 16 | 16 |
| 6th February 1925 . . | 14 | nil. | 962 | 28 | 22 |
| 13th February 1925 . . | .. | .. | 583 | 14 | 14 |
| 20th February 1925 . . | .. | .. | 625 | 39 | 39 |
| 27th February 1925 . . | .. | .. | 357 | 27 | 27 |
| 6th March 1925 . . | .. | .. | 116 | 5 | 5 |

TABLE VIII

Weekly population of Pink Boll-worm larvae on 25 cotton plants at Surat (1925-26)

| Week ending | No. of flower-buds | Larvae in flower-buds | No. of bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|--------------------------|-----------------------|--------------------------|-----------------|--------------------|-----------------------------------|
| 18th December 1925 . . . | 323 | .. | 151 | 1 | 1 |
| 25th December 1925 . . . | 927 | .. | 1,186 | 29 | 29 |
| 1st January 1926 . . . | 401 | 1 | 1,127 | 61 | 62 |
| 8th January 1926 . . . | 97 | .. | 966 | 68 | 68 |
| 15th January 1926 . . . | 49 | .. | 821 | 85 | 85 |
| 22nd January 1926 . . . | 24 | .. | 838 | 99 | 99 |
| 29th January 1926 . . . | .. | .. | 795 | 131 | 131 |

TABLE IX

Weekly population of Pink Boll-worm larvae on 25 cotton plants at Surat (1926-27)

| Week ending | No. of flower-buds | Larvae in flower-buds | No. of bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|--------------------------|-----------------------|--------------------------|-----------------|--------------------|-----------------------------------|
| 28th January 1927 . . . | 2,090 | .. | 229 | 2 | 2 |
| 4th February 1927 . . . | 1,890 | .. | 439 | .. | .. |
| 11th February 1927 . . . | 1,991 | .. | 794 | 12 | 12 |
| 18th February 1927 . . . | 1,271 | .. | 1,081 | 12 | 12 |
| 25th February 1927 . . . | 682 | .. | 1,071 | 14 | 14 |
| 4th March 1927 . . . | 195 | .. | 888 | 46 | 46 |
| 11th March 1927 . . . | 9 | .. | 801 | 69 | 69 |
| 18th March 1927 . . . | .. | .. | 731 | 51 | 51 |
| 25th March 1927 . . . | .. | .. | 690 | 32 | 32 |
| 1st April 1927 . . . | .. | .. | 286 | 11 | 11 |
| 8th April 1927 . . . | .. | .. | 116 | 3 | 3 |

TABLE X

Weekly population of Pink Boll-worm larvae on cotton plants at Surat (1927-28)

| No. of plants | Dates | Flower-buds | Larvae in flower-buds | Bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|---------------|-------------------------|-------------|-----------------------|-------|-----------------|-----------------------------|
| 25 | 16th September 1927 . . | .. | .. | .. | .. | 1 in shoot. |
| 25 | 28th October 1927 . . | 1,158 | 2 | .. | .. | 2 |
| 25 | 4th November 1927 . . | 1,263 | 2 | .. | .. | 2 |
| 25 | 11th November 1927 . . | 1,976 | .. | .. | .. | .. |
| 25 | 18th November 1927 . . | 2,392 | 4 | 2 | .. | 4 |
| 25 | 25th November 1927 . | 2,421 | .. | 13 | 2 | 2 |
| 25 | 2nd December 1927 . . | 2,302 | 5 | 2 | 2 | 7 |
| 25 | 9th December 1927 . . | 2,496 | 6 | 107 | 9 | 15 |
| 25 | 16th December 1927 . . | 3,501 | 3 | 170 | 6 | 9 |
| 25 | 23rd December 1927 . . | 3,541 | 6 | 189 | 7 | 13 |
| 25 | 30th December 1927 . . | 3,530 | 4 | 406 | 28 | 32 |
| 25 | 7th January 1928 . . | 3,204 | .. | 487 | 17 | 17 |
| 25 | 14th January 1928 . . | 2,763 | 3 | 1,156 | 60 | 63 |
| 25 | 21st January 1928 . . | 1,849 | .. | 1,164 | 18 | 18 |
| 25 | 28th January 1928 . . | 968 | .. | 1,357 | 33 | 33 |
| 25 | 4th February 1928 . . | 607 | .. | 1,629 | 35 | 35 |
| 25 | 11th February 1928 . . | 105 | .. | 1,270 | 63 | 63 |
| 25 | 18th February 1928 . . | 56 | .. | 1,001 | 87 | 87 |
| 25 | 25th February 1928 . . | 14 | .. | 857 | 123 | 123 |
| 25 | 3rd March 1928 . . | 4 | .. | 527 | 67 | 67 |
| 25 | 10th March 1928 . . | .. | .. | 300 | 18 | 18 |

TABLE X—*contd**Weekly population of Pink Boll-worm larvae on cotton plants at Surat (1927-28)—contd*

| No. of plants | Dates | Flower-buds | Larvae in flower-buds | Bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|---------------|-----------------------|-------------|-----------------------|-------|-----------------|-----------------------------|
| 25 | 17th March 1928 . . | .. | .. | 183 | 19 | 19 |
| 25 | 24th March 1928 . . | .. | .. | 72 | 8 | 8 |
| 50 | 31st March 1928 . . . | 58 | .. | 51 | 11 | 11 |
| 50 | 5th April 1928 . . . | 107 | .. | 16 | 4 | 4 |
| 50 | 13th April 1928 . . . | 346 | .. | 1 | .. | .. |
| 50 | 20th April 1928 . . . | 350 | .. | 2 | .. | .. |
| 50 | 27th April 1928 . . . | 386 | .. | 1 | .. | .. |
| 50 | 4th May 1928 . . . | 471 | .. | 21 | .. | .. |
| 50 | 9th May 1928 . . . | 665 | .. | 30 | .. | .. |
| 50 | 18th May 1928 . . . | 488 | .. | 43 | .. | .. |
| 50 | 23rd May 1928 . . . | 362 | .. | 58 | .. | .. |
| 50 | 31st May 1928 . . . | 346 | .. | 74 | 3 | 3 |
| 50 | 8th June 1928 . . . | 278 | .. | 48 | 1 | 1 |
| 50 | 13th June 1928 . . . | 132 | .. | 18 | 5 | 5 |
| 50 | 20th June 1928 . . . | 69 | . | 8 | 1 | 1 |
| 50 | 27th June 1928 . . . | 83 | .. | 4 | .. | .. |
| 50 | 4th July 1928 . . . | 280 | .. | 6 | .. | .. |
| 50 | 11th July 1928 . . . | 136 | . | 8 | 1 | 1 |
| 50 | 17th July 1928 . . . | 130 | .. | 8 | 5 | 5 |
| 50 | 24th July 1928 . . . | 54 | .. | 11 | 7 | 7 |
| 50 | 3rd August 1928 . . | 93 | 1 | 4 | 1 | 4 |
| 50 | 8th August 1928 . . | 37 | .. | 7 | 2 | 2 in shoot. 2 |

TABLE XI

Weekly population of Pink Boll-worm larvae on cotton plants at Surat (1928-29)

| No. of plants | Dates | No. of flower-buds | Larvae in flower-buds | No. of green bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|---------------|------------------------|--------------------|-----------------------|--------------------|-----------------|-----------------------------|
| 25 | 9th November 1928 . . | 2,758 | 1 | 4 | .. | 1 |
| 25 | 16th November 1928 . . | 2,767 | 4 | 50 | .. | 4 |
| 25 | 23rd November 1928 . . | 3,006 | 4 | 119 | 4 | 8 |
| 25 | 30th November 1928 . . | 3,235 | .. | 210 | 2 | 2 |
| 25 | 8th December 1928 . . | 3,201 | 3 | 314 | 1 | 4 |
| 25 | 14th December 1928 . . | 2,797 | 2 | 338 | 13 | 15 |
| 25 | 21st December 1928 . . | 2,915 | 1 | 610 | 7 | 8 |
| 25 | 29th December 1928 . . | 3,162 | .. | 863 | 28 | 28 |
| 25 | 4th January 1929 . . | 1,744 | .. | 863 | 22 | 22 |
| 25 | 11th January 1929 . . | 1,025 | .. | 851 | 6 | 6 |
| 25 | 18th January 1929 . . | 804 | .. | 1,062 | 12 | 12 |
| 25 | 25th January 1929 . . | 681 | .. | 1,005 | 32 | 32 |
| 25 | 1st February 1929 . . | 188 | .. | 1,027 | 71 | 71 |
| 25 | 8th February 1929 . . | 30 | .. | 794 | 107 | 107 |
| 25 | 15th February 1929 . . | 3 | .. | 761 | 200 | 200 |
| 25 | 22nd February 1929 . . | 1 | .. | 586 | 143 | 143 |
| 25 | 20th February 1929 . . | .. | .. | 416 | 100 | 100 |
| 25 | 8th March 1929 . . | .. | .. | 175 | 21 | 21 |
| 25 | 15th March 1929 . . | 5 | .. | 35 | 4 | 4 |
| 25 | 22nd March 1929 . . | 17 | .. | 10 | .. | .. |
| 50 | 28th March 1929 . . | 75 | .. | 2 | .. | .. |
| 50 | 6th April 1929 . . | 293 | .. | 4 | .. | .. |
| 50 | 12th April 1929 . . | 310 | .. | 16 | .. | .. |
| 50 | 19th April 1929 . . | 554 | .. | 26 | .. | .. |
| 50 | 25th April 1929 . . | 469 | .. | 39 | 1 | 1 |
| 50 | 2nd May 1929 . . | 310 | .. | 44 | .. | .. |
| 50 | 10th May 1929 . . | 337 | .. | 26 | .. | .. |
| 50 | 16th May 1929 . . | 354 | .. | 21 | .. | .. |
| 50 | 24th May 1929 . . | 231 | .. | 17 | 1 | 1 |
| 50 | 29th May 1929 . . | 467 | .. | 54 | 6 | 6 |

TABLE XII

Weekly population of Pink Boll-worm larvae on cotton plants at Surat (1929-30)

| No. of plants | Dates | No. of flower-buds | Larvae in flower-buds | No. of green bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|---------------|------------------------|--------------------|-----------------------|--------------------|-----------------|-----------------------------|
| 25 | 30th October 1929 . . | 1,667 | 1 | 8 | .. | 1 |
| 25 | 8th November 1929 . . | 1,993 | .. | 5 | .. | .. |
| 25 | 14th November 1929 . . | 2,627 | .. | 17 | .. | .. |
| 25 | 21st November 1929 . . | 2,529 | .. | 20 | .. | .. |
| 25 | 28th November 1929 . . | 3,381 | .. | 32 | .. | .. |
| 25 | 5th December 1929 . . | 3,637 | .. | 72 | 2 | 2 |
| 25 | 12th December 1929 . . | 3,829 | .. | 149 | 2 | 2 |
| 25 | 19th December 1929 . . | 2,673 | .. | 338 | 1 | 1 |
| 25 | 26th December 1929 . . | 2,831 | .. | 493 | .. | .. |
| 25 | 2nd January 1930 . . | 2,537 | .. | 866 | 3 | 3 |
| 25 | 9th January 1930 . . | 1,721 | .. | 1,350 | .. | .. |
| 25 | 16th January 1930 . . | 940 | .. | 1,814 | 6 | 6 |
| 25 | 23rd January 1930 . . | 288 | .. | 1,468 | 19 | 19 |
| 25 | 30th January 1930 . . | 67 | .. | 1,102 | 18 | 18 |
| 25 | 6th February 1930 . . | 20 | .. | 1,170 | 31 | 31 |
| 25 | 13th February 1930 . . | 5 | .. | 947 | 22 | 22 |
| 25 | 20th February 1930 . . | .. | .. | 882 | 14 | 14 |
| 25 | 27th February 1930 . . | .. | .. | 693 | 10 | 10 |
| 25 | 6th March 1930 . . | .. | .. | 334 | 2 | 2 |
| 25 | 13th March 1930 . . | .. | .. | 67 | 2 | 2 |
| 25 | 20th March 1930 . . | .. | .. | 13 | 1 | 1 |
| 25 | 27th March 1930 . . | .. | .. | 3 | .. | .. |
| 50 | 5th April 1930 . . | 16 | .. | .. | .. | .. |
| 50 | 11th April 1930 . . | 43 | .. | 1 | .. | .. |
| 50 | 16th April 1930 . . | 198 | .. | 3 | .. | .. |
| 50 | 23rd April 1930 . . | 263 | .. | 6 | .. | .. |
| 50 | 29th April 1930 . . | 323 | .. | 10 | .. | .. |
| 50 | 6th May 1930 . . | 395 | .. | 19 | 1 | 1 |
| 50 | 14th May 1930 . . | 591 | .. | 49 | 1 | 1 |
| 50 | 22nd May 1930 . . | 628 | .. | 86 | 1 | 1 |
| 50 | 27th May 1930 . . | 620 | .. | 243 | 2 | 2 |

TABLE XIII

Weekly population of Pink Boll-worm larvae on 25 plants at Surat (1930-31)

| Dates | Buds | Larvae in buds | Bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|------------------------|-------|----------------|-------|-----------------|-----------------------------|
| 21st November 1930 . . | 2,688 | 1 | 32 | 1 | 2 |
| 28th November 1930 . . | 3,998 | 2 | 75 | 11 | 13 |
| 5th December 1930 . . | 5,192 | 5 | 208 | 8 | 13 |
| 12th December 1930 . . | 4,224 | 15 | 288 | 17 | 32 |
| 19th December 1930 . . | 4,207 | 21 | 868 | 28 | 49 |
| 26th December 1930 . . | 3,919 | 25 | 1,707 | 87 | 112 |
| 2nd January 1931 . . | 2,198 | 4 | 1,895 | 65 | 69 |
| 9th January 1931 . . | 927 | 2 | 2,167 | 57 | 59 |
| 16th January 1931 . . | 225 | 3 | 1,692 | 48 | 51 |
| 23rd January 1931 . . | 18 | .. | 1,248 | 51 | 51 |
| 30th January 1931 . . | 1 | .. | 1,165 | 81 | 81 |
| 6th February 1931 . . | .. | .. | 1,287 | 84 | 84 |
| 13th February 1931 . . | .. | .. | 894 | 56 | 56 |
| 20th February 1931 . . | .. | .. | 470 | 27 | 27 |
| 27th February 1931 . . | .. | .. | 102 | 9 | 9 |

TABLE XIV

Weekly population of Pink Boll-worm larvae on 25 plants each week

1927-28 (Broach)

| Dates | Buds | Larvae in buds | Bolls | Larvae in bolls | Total Pink Boll-worm larvae |
|------------------------|-------|----------------|-------|-----------------|-----------------------------|
| 10th December 1927 . . | 2,520 | .. | 379 . | 2 | 2 |
| 17th December 1927 . . | 2,449 | 1 | 836 | 19 | 20 |
| 24th December 1927 . . | 1,081 | .. | 676 | 19 | 19 |
| 31st December 1927 . . | 632 | .. | 939 | 17 | 17 |
| 7th January 1928 . . . | 429 | .. | 755 | 56 | 56 |
| 14th January 1928 . . | 366 | .. | 592 | 36 | 36 |
| 21st January 1928 . . | 95 | .. | 642 | 45 | 45 |
| 28th January 1928 . . | 38 | .. | 671 | 38 | 38 |
| 4th February 1928 . . | 8 | .. | 609 | 38 | 38 |
| 11th February 1928 . . | .. | .. | 580 | 42 | 42 |
| 18th February 1928 . . | .. | .. | 457 | 40 | 40 |
| 25th February 1928 . . | .. | .. | 349 | 15 | 15 |
| 3rd March 1928 . . . | .. | .. | 464 | 7 | 7 |

TABLE XV

Pink Boll-worm larvae for every 100 bolls examined

[illegible]

APPENDIX VII

PINK BOLL-WORM (*PLATYEDRA GOSSYPIELL* 1 SAUND.) IN
KHANDESH

A few observations about the Pink Boll-worm were maintained during the period of work upon Spotted Boll-worms of nearly 18 months in Khandesh. The incidence of the larval population of this pest in the cotton fields during 1929 and 1930 can be seen from the following tables :—

TABLE I

Larval population of Pink Boll-worm on 100 cotton plants from Government Farm, Julgaon

1929

| Weeks | No. of green bolls | No. of Pink Boll-worm larvae | REMARKS |
|-------------------------------|--------------------|------------------------------|----------------------|
| 24th August 1929 | 49 | 1 | |
| 31st August 1929 | 148 | .. | |
| 7th September 1929 | 270 | 3 | |
| 14th September 1929 | 305 | .. | Bolls began to open. |
| 21st September 1929 | 622 | 3 | |
| 28th September 1929 | 774 | 5 | |
| 5th October 1929 | 734 | 12 | |
| 12th October 1929 | 488 | 14 | 1st picking. |
| 19th October 1929 | 210 | 38 | |
| 26th October 1929 | 100 | 47 | 2nd picking. |
| 2nd November 1929 | 55 | 36 | |
| 9th November 1929 | 23 | 6 | |
| 16th November 1929 | 6 | .. | Last picking. |

TABLE II

Pink Boll-worm larvae from 50 full-grown green bolls collected from the standing cotton stalks on Jalgaon Farm

1929-30

| Weeks | No. of Pink Boll-worm larvae |
|------------------------------|------------------------------|
| 23rd November 1929 | 28 |
| 30th November 1929 | 90 |
| 7th December 1929 | 69 |
| 14th December 1929 | 59 |
| 21st December 1929 | 4 |
| 28th December 1929 | 47 |
| 4th January 1930 | 71 |
| 11th January 1930 | 38 |
| 18th January 1930 | 44 |
| 25th January 1930 | 44 |
| 1st February 1930 | 9 |
| 8th February 1930 | 15 |
| 15th February 1930 | 18 |
| 22nd February 1930 | 13 |
| 1st March 1930 | 9 |
| 8th March 1930 | 20 |
| 15th March 1930 | 13 |
| 22nd March 1930 | 5 |
| 29th March 1930 | 4 |

TABLE III

Pink Boll-worm larvae on 100 cotton plants from Jalgaon Farm

1930

| Weeks | No. of green bolls | No. of larvae | REMARKS |
|------------------------------|-----------------------|---------------|---------------|
| 4th October 1930 | 499 | .. . | |
| 11th October 1930 | 585 | 2 | |
| 18th October 1930 | 458 | 19 | |
| 25th October 1930 | 576 | 73 | 1st picking. |
| 1st November 1930 | 458 | 75 | |
| 8th November 1930 | 315 | 48 | |
| 15th November 1930 | 154 | 18 | |
| 22nd November 1930 | 170 | 147 | 2nd picking. |
| 29th November 1930 | 154 | 160 | |
| 6th December 1930 | 145 | 156 | |
| 13th December 1930 | 181 | 294 | Last picking. |
| 20th December 1930 | 103 | 156 | |
| 27th December 1930 | 141 | 138 | |
| 3rd January 1931 | 80 | 25 | |
| 10th January 1931 | 140 | 109 | |
| 17th January 1931 | 344 | 228 | |
| 24th January 1931 | 338 | 209 | |
| 31st January 1931 | 496 | 237 | |

TABLE IV

Larval population of Pink Boll-worm on 50 cotton plants from villages near Jalgaon during 1930-31

| Fortnights of examination | | Mamurabad | | Pimrala | | Nagardevala | |
|---------------------------|-------------------|-------------|--------|-------------|--------|-------------|--------|
| | | Green bolls | Larvae | Green bolls | Larvae | Green bolls | Larvae |
| September | { 1st Fortnight . | 79 | .. | 155 | .. | 626 | 18 |
| | { 2nd Fortnight . | 241 | 5 | 147 | .. | .. | .. |
| October | { 1st Fortnight . | 291 | 34 | 152 | 43 | 318 | 105 |
| | { 2nd Fortnight . | 446 | 166 | 169 | 34 | .. | .. |
| November | { 1st Fortnight . | 338 | 75 | 103 | 55 | 155 | 157 |
| | { 2nd Fortnight . | 75 | 52 | 78 | 110 | .. | .. |
| December | { 1st Fortnight . | 203 | 335 | 57 | 42 | 150 | 149 |
| | { 2nd Fortnight . | 65 | 55 | 58 | 54 | .. | .. |
| January | { 1st Fortnight . | 99 | 85 | 86 | 68 | .. | .. |
| | { 2nd Fortnight . | 100 | 72 | 73 | 17 | .. | .. |

The cotton season of 1929 was very early and practically all the crop was over before the population of Pink Boll-worm could rise. During 1930, however, the rapid multiplication of these insects had begun at about the time when the first picking of cotton was in progress. From the last week of November till the last week of December there was found a larger number of these larvae than the total number of bolls which were examined. Even after the last picking was over a large number of green bolls could be collected from the cotton fields but no *kapas* was available from them because most of them were destroyed by this pest.

In order to determine the period when the caterpillars hibernated in Khandesh, 200 full grown green bolls were collected, three times during the season of 1929 when the first, second and the third pickings respectively were in progress. The emergence from these bolls is given in the next table.

TABLE V

Emergence of Pink Boll-worm moths from developed green bolls collected on different dates

| Month of emergence | Number of Pink Boll-worm moths which emerged | | |
|-------------------------|--|---|---|
| | 200 bolls collected on 4th October 1929 | 200 bolls collected on 15th November 1929 | 200 bolls collected on 18th November 1929 |
| October 1929 | 2 | 7 | .. |
| November 1929 | 1 | 15 | .. |
| December 1929 | .. | 1 | 78 |
| January 1930 | .. | 1 | 8 |
| February 1930 | .. | 1 | 5 |
| March 1930 | .. | .. | 5 |
| April 1930 | .. | .. | 2 |
| May 1930 | .. | .. | .. |
| June 1930 | .. | .. | 4 |
| July 1930 | 1 | 6 | 28 |
| August 1930 | .. | .. | 2 |

After the pickings are over there are always some dry bolls, which remain on the plants, being rejected as useless bolls at the time of collecting *kapas*. There is a possibility of the long-cycle larvae resting in these bolls. These bolls either drop away in the fields or are carried along with the cotton sticks, which are generally heaped up for fuel purposes. Two hundred such bolls were collected on 17th January 1930 from one of the heaps and were kept under observation.

From these bolls, two Pink Boll-worm moths emerged in July 1930 and three more moths came out in August 1930.

From the information upon the emergence of these moths from green bolls collected at different times of the season, it appears that the largest number of larvae hibernate when the crop is about to be over. Some of these long-cycle larvae can also be found in the dry bolls which are rejected on the cotton plants.

A few moths emerged practically every month till August 1930, from the bolls which were collected in November 1929; but the largest number of moths emerged during July 1930. There was no emergence after August.

Some of the moths which emerged from the resting stage were kept under observation to note their longevity. Seventeen of them were kept without any food, nine of them were given only water, and the remaining two were offered diluted molasses, (Table VI App. VII.)

The average life of the moths which were kept without any food was 8 days although one of them lived for 24 days. The moths which were supplied with water only, lived for 11 days on an average, and the maximum period of life was 19 days. The moths which were given diluted molasses lived for 25 and 30 days respectively.

It will be seen from Table II (App. VII) that the short-cycle larvae continued to breed in the cotton fields on the fresh growth from the standing plants till the end of March 1930.

The information collected about this pest in Khandesh is scanty and refers practically to only one complete season. However, it appears that Pink Boll-worm is a more serious pest in Khandesh than the Spotted Boll-worms, and that the damage due to this pest is greater in years when the crop is delayed due to heavy rains during September, October or November. The damage is not restricted to the dirty *kapas* which finds its way to the market, but there is, in addition, a large number of bolls which are entirely valueless and have to be rejected at the time of collecting the *kapas*.

TABLE VI

Longevity of life of the Pink Boll-worm moths emerging from resting stage at Jalgaon

1930

| Serial No. of moths | Date of emergence | Date on which the moth was dead | Total life in days |
|---|---------------------------|---------------------------------|--------------------------|
| <i>Moths kept unfed.</i> | | | |
| 1 | 30th June 1930 | 7th July 1930 | 7 |
| 2 | 11th July 1930 | 20th July 1930 | 9 |
| 3 | 19th July 1930 | 23rd July 1930 | 4 |
| 4 | 19th July 1930 | 28th July 1930 | 9 |
| 5 | 21st July 1930 | 28th July 1930 | 7 |
| 6 | 22nd July 1930 | 23rd July 1930 | 1 |
| 7 | 22nd July 1930 | 23rd July 1930 | 1 |
| 8 | 22nd July 1930 | 26th July 1930 | 4 |
| 9 | 22nd July 1930 | 27th July 1930 | 5 |
| 10 | 22nd July 1930 | 1st August 1930 | 10 |
| 11 | 22nd July 1930 | 15th August 1930 | 24 |
| 12 | 25th July 1930 | 5th August 1930 | 12 |
| 13 | 26th July 1930 | 4th August 1930 | 9 |
| 14 | 26th July 1930 | 5th August 1930 | 8 |
| 15 | 28th July 1930 | 31st July 1930 | 3 |
| 16 | 4th August 1930 | 16th August 1930 | 12 |
| 17 | 8th August 1930 | 27th August 1930 | 19 |
| | | Average life | 8 |
| <i>Moths which were given water only.</i> | | | |
| 1 | 4th July 1930 | 8th July 1930 | 4 |
| 2 | 14th July 1930 | 21st July 1930 | 7 |
| 3 | 19th July 1930 | 30th July 1930 | 11 |
| 4 | 19th July 1930 | 1st August 1930 | 13 |
| 5 | 19th July 1930 | 4th August 1930 | 16 |
| 6 | 19th July 1930 | 30th July 1930 | 11 |
| 7 | 23rd July 1930 | 26th July 1930 | 3 |
| 8 | 23rd July 1930 | 4th August 1930 | 12 |
| 9 | 23rd July 1930 | 11th August 1930 | 19 |
| | | Average life | 11 |
| <i>Moths which were given diluted molasses.</i> | | | |
| 1 | 28th July 1930 | 22nd August 1930 | 25 |
| 2 | 28th July 1930 | 27th August 1930 | 30 |

APPENDIX VIII

CAPACITY OF THE COTTON PLANT TO REPLACE ITS FORMS SHED DUE TO THE BOLL-WORM INJURY

It has been established that the Spotted Boll-worms appear in the cotton field as soon as vigorous growth of the cotton plant commences, and continue to feed on them throughout the season. The flower-buds and the young bolls which are attacked by the worms drop down from the plant, resulting in a continuous wastage of the reproductive organs of the cotton plant. It is, of course, inconceivable that the cotton plant should mature all its flower-buds and flowers into bolls, but it was not very clear if all the shedding due to the insects was a net loss, or if the plant possessed a capacity to recover any loss from such shedding by stronger development of the forms which would normally appear later, or by augmenting their number by additional bud formation.

A small beginning for ascertaining the possibilities was, therefore, made during 1925, when a group of only five plants was selected, and the flowers that appeared on them were pruned continuously up to a fixed date in the case of each plant. Thus the pruning of flowers from the first plant was stopped on the 30th of November; and that from the others on the 10th December, 20th December, 30th December, and 10th of January respectively. The flowers that appeared after these dates were not disturbed. Daily flowering was recorded from the commencement of the experiment and the shed bolls were collected to examine if they were attacked by the boll-worms. The results are tabulated below :—

TABLE I

| Dates up to which the flowers were pruned | Plant numbers | | | | |
|---|---------------|---------------|---------------|---------------|--------------|
| | 1 | 2 | 3 | 4 | 5 |
| | 30th November | 10th December | 20th December | 30th December | 10th January |
| Number of flowers pruned | 6 | 49 | 236 | 75 | 118 |
| Total number of flowers opened. | 96 | 120 | 287 | 81 | 118 |
| Number of flowers opened after pruning was stopped. | 90 | 71 | 51 | 6 | <i>Nil</i> |
| Number of mature bolls | 33 | 37 | 40 | 4 | <i>Nil</i> |
| Number of bolls shed after pruning was stopped. | 57 | 34 | 11 | 2 | <i>Nil</i> |
| Bolls shed due to boll-worm | 10 | 7 | 7 | <i>Nil</i> | <i>Nil</i> |
| Bolls shed due to other causes | 47 | 27 | 4 | 2 | <i>Nil</i> |

The results of these observations on single plants could not be taken as conclusive evidence, but indicated that the plants had responded, in their effort to make up the number of mature bolls, by developing a large proportion of flowers immediately after the removing of flowers was stopped. The performance of the plant No. 3 especially deserves admiration in as much as it yielded 40 mature bolls from 51 late flowers, after undergoing the pruning of 236 of its first flowers.

The continuation of the removal of flowers from plants Nos. 4 and 5 ultimately proved disastrous to the plants, as only six flowers could open on plant No. 4 after the pruning was stopped; and there was no more flowering on the 5th plant after the tenth of January. It was thus indicated that the plant cannot stand an indefinite destruction of its flowers, and that any unusual and continuous damage to the flowers would exhaust the capacity of the plant to replace the lost forms.

Experiments on a larger scale were undertaken in 1926. A total number of forty plants was grown free from boll-worms by "night caging", to avoid the disturbing factor of the shedding due to boll-worms in addition to the artificial removal of the forms. Out of these, 15 plants were used for the pruning of the flower-buds, and the remaining 25 for the trials of the flower pruning.

The first 15 plants were divided into 3 groups of 5 plants each. The first group was reserved as a control. From the second, all the flower-buds formed up to the 30th of October were pruned. From the third, the flower-buds were similarly pruned on the 13th of November. Care was taken to remove only such of the flower-buds as could be nipped off without injuring the growing points, and therefore the smallest flower-buds were left alone. Thus 236 flower-buds were removed of the total of 358 existing on the plants of the 2nd set on the 30th of October; and 202 buds out of 336 present on the plants of the 3rd set on the 13th of November. Records of the weekly flower-bud formation, flower opening and the relative success of flowers to bolls were maintained. (Tables III and IV of this Appendix.)

The result of the flower-bud pruning (as can be seen from the tables) was that the flowering was delayed by four weeks in the second set, and by seven weeks in the third one. The flowering, however, when it appeared showed a much higher intensity and a larger proportion of the first flowers was retained than in the control plants. Attention is drawn to another important fact, that after the pruning of the flower-buds, the fresh flower-bud formation in both pruned sets, continued at a higher level for longer periods than that in the control set. This indicated that not only does the plant respond to the loss of its flower-buds by retaining its later forms, but that it also augments bud formation by new development.

FLOWER-PRUNING

The remaining 25 plants were divided into 5 sets of 5 plants each. Four sets were used for the removal of the flowers and the fifth was retained as a control. The flowers that appeared in all the sets (except the control) were removed daily up to the 13th November, 27th November, 11th December, and the 25th of December respectively.

Records of daily flower-opening, the number of flowers removed, number of flowers that opened after the removal of flowers was stopped and the relative percentage of flowers that developed into mature bolls were maintained. (Table V of this appendix.)

Examination of the curves of flower opening (Plate XXV) shows that the peak of flowering is reached in all the sets at about the same time but that, where the removal of flowers is prolonged, the intensity of flower opening continues to be higher, instead of declining as in the control plants. The prolongation of the high rate of flower opening, following continued flower-pruning, naturally increases the average number of flowers per plant, in proportion to the duration of the period of the removal. The average performance of the plants from the above five sets is shown in the following table.

TABLE II

Results of flower-pruning


(Average per plant of 5 plants in each case)

| Treatment | Flowers per plant | Flowers pruned | Flowers opened after pruning was stopped | Mature bolls per plant |
|--|-------------------|----------------|--|------------------------|
| Control | 75 | <i>Nil</i> | 75 | 30 |
| 1st set (flowers pruned up to 13th November) . . | 70 | 5 | 65 | 25 |
| 2nd set (flowers pruned up to 28th November) . | 81 | 20 | 61 | 22.8 |
| 3rd set (flowers pruned up to 11th December) . . | 107 | 42 | 65 | 23.6 |
| 4th set (flowers pruned up to 25th December) . . | 126 | 71 | 55 | 24.2 |

It will further be seen that the number of mature bolls per plant is not substantially different, and curiously enough the number of bolls shed in every case is almost similar except in the last case; showing that the plants had an adequate supply of flowers to develop into bolls and even a few more to shed in the normal course, even after the pruning of the first flowers.

These experiments have, therefore, shown that the cotton plant under Surat conditions evinces a large capacity for replacing the lost flower-buds or flowers, by development of its later forms, which might otherwise have shed. It further indicates a certain degree of auto-regulation of the formation of flower-buds and flowers, thus assuring enough supply of these forms for the requirement of the plant.

It should, however, be remembered that the preliminary experiment indicated that this capacity for replacing forms is limited, and that the plant is unable to make good the destruction of its forms indefinitely.



Pruning of Flowers 1926-27.

Showing weekly flowering, proportion of flowers pruned and the number of successful bolls from weekly flowers in the flower-pruning experiments of 1926-27

TABLE III

Flower-bud formation, flower-opening and the success of flowers into mature bolls from five plants where 226 flower-buds were removed on the 30th of October, 1926

(5 plants in each case)

| Weeks | Control plot | | | 226 flower-buds removed | | |
|---------------------|----------------------|----------------|-----------------------------|-------------------------|----------------|-----------------------------|
| | Flower-bud formation | Flower opening | Success of flowers to bolls | Flower-bud formation | Flower opening | Success of flowers to bolls |
| 3rd October . . . | .. | .. | .. | 20 | .. | .. |
| 10th October . . . | 40 | .. | .. | 31 | .. | .. |
| 17th October . . . | 63 | .. | .. | 73 | .. | .. |
| 24th October . . . | 85 | .. | .. | 97 | .. | .. |
| 30th October . . . | 116 | 4 | 4 | 137* | .. | .. |
| 6th November . . . | 201 | 15 | 12 | 192 | .. | .. |
| 13th November . . . | 147 | 26 | 19 | 194 | .. | .. |
| 20th November . . . | 163 | 37 | 21 | 230 | .. | .. |
| 27th November . . . | 132 | 48 | 38 | 217 | 13 | 12 |
| 4th December . . . | 133 | 57 | 36 | 138 | 96 | 81 |
| 11th December . . . | 100 | 60 | 9 | 118 | 107 | 47 |
| 18th December . . . | 28 | 79 | 16 | 61 | 110 | 16 |
| 25th December . . . | 19 | 69 | 10 | 23 | 112 | 20 |
| 1st January . . . | 4 | 35 | 7 | 5 | 82 | 25 |
| 8th January . . . | 2 | 17 | 4 | 4 | 39 | 8 |
| 15th January . . . | .. | 4 | .. | .. | 16 | 1 |
| 22nd January . . . | .. | 1 | .. | .. | 1 | .. |

* 226 flower-buds removed on the 30th of October 1926.

TABLE IV

Flower-bud formation, flower-opening and the relative success of flowers into mature bolls from five plants where 202 flower-buds were removed on the 13th November 1926

(5 night-caged plants in each case)

| Weeks | Control plants | | | Plants with 202 flower-buds removed | | |
|---------------------|----------------------|----------------|----------------------------|-------------------------------------|----------------|-----------------------------|
| | Flower-bud formation | Flower opening | Success of flower to bolls | Flower-bud formation | Flower opening | Success of flowers to bolls |
| 10th October . . . | 40 | .. | .. | 8 | .. | .. |
| 17th October . . . | 63 | .. | .. | 27 | .. | .. |
| 24th October . . . | 85 | .. | .. | 43 | .. | .. |
| 30th October . . . | 116 | 4 | 4 | 64 | .. | .. |
| 6th November . . . | 201 | 15 | 12 | 98 | .. | .. |
| 13th November . . . | 147 | 26 | 19 | 96* | .. | .. |
| 20th November . . . | 163 | 37 | 21 | 140 | .. | .. |
| 27th November . . . | 132 | 48 | 38 | 171 | .. | .. |
| 4th December . . . | 133 | 57 | 36 | 132 | .. | .. |
| 11th December . . . | 100 | 60 | 9 | 108 | .. | .. |
| 18th December . . . | 28 | 79 | 16 | 100 | 47 | 46 |
| 25th December . . . | 19 | 69 | 10 | 49 | 97 | 70 |
| 1st January . . . | 4 | 35 | 7 | 44 | 82 | 12 |
| 8th January . . . | 2 | 17 | 4 | 15 | 85 | 3 |
| 15th January . . . | .. | 4 | .. | 5 | 69 | 4 |
| 22nd January . . . | .. | 1 | .. | 1 | 24 | 2 |
| 29th January . . . | .. | .. | .. | .. | 4 | .. |

* 202 flower-buds removed on the 13th November 1926.

TABLE V

Statement showing daily flower opening, flower pruning, boll shedding and the percentage success of bolls from the different sets of the flower pruning experiment.

1926-27

| Week ending | CHECK — (10 PLANTS) | | | 1ST SET :— (5 PLANTS) | | | 2ND SET — (5 PLANTS) | | | 3RD SET — (5 PLANTS) | | | 4TH SET — (5 PLANTS) | | |
|--------------------|---------------------|------------|-----------|-----------------------|------------|-----------|----------------------|------------|-----------|----------------------|------------|-----------|----------------------|------------|-----------|
| | Flowers | Bolls Shed | % success | Flowers | Bolls Shed | % success | Flowers | Bolls Shed | % success | Flowers | Bolls Shed | % success | Flowers | Bolls Shed | % success |
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 |
| 30th October 1926 | 3 | . | 100 | 1 | Pruned | .. | 1 | Pruned | .. | 2 | Pruned | .. | 2 | Pruned | .. |
| 6th November 1926 | 11 | 1 | 90.9 | 6 | . | . | 9 | .. | .. | 7 | .. | .. | 13 | .. | .. |
| 13th November 1926 | 28 | 2 | 92.9 | 17 | .. | . | 19 | .. | .. | 17 | .. | .. | 18 | .. | .. |
| 20th November 1926 | 49 | 9 | 81.6 | 37 | 7 | 81.1 | 33 | .. | .. | 25 | . | .. | 38 | .. | .. |
| 27th November 1926 | 57 | 8 | 86.0 | 48 | 16 | 66.6 | 43 | .. | .. | 37 | .. | .. | 96 | .. | .. |
| 4th December 1926 | 111 | 50 | 55.0 | 56 | 23 | 38.9 | 58 | 2 | 96.7 | 63 | .. | . | 48 | .. | .. |
| 11th December 1926 | 121 | 94 | 22.3 | 74 | 62 | 16.2 | 58 | 17 | 70.7 | 59 | . | . | 58 | .. | .. |
| 18th December 1926 | 125 | 101 | 19.2 | 46 | 37 | 19.6 | 49 | 44 | 10.2 | 55 | 9 | 83.6 | 74 | .. | .. |
| 25th December 1926 | 122 | 93 | 23.8 | 35 | 31 | 11.4 | 62 | 60 | 3.2 | 66 | 20 | 69.7 | 71 | .. | .. |
| 1st January 1927 | 79 | 51 | 35.4 | 24 | 22 | 8.3 | 41 | 34 | 17.1 | 54 | 38 | 33.3 | 65 | 7 | 89.2 |
| 8th January 1927 | 38 | 28 | 26.3 | 6 | 3 | 50.0 | 26 | 23 | 11.5 | 60 | 55 | 8.3 | 61 | 12 | 80.3 |
| 15th January 1927 | 6 | 6 | . | . | .. | .. | 5 | 5 | . | 41 | 39 | 4.9 | 51 | 38 | 25.5 |
| 22nd January 1927 | .. | .. | .. | .. | .. | .. | 1 | 1 | .. | 31 | 30 | 3.2 | 51 | 51 | .. |
| 29th January 1927 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 15 | 15 | .. | 27 | 26 | 3.7 |
| 5th February 1927 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 4 | 4 | .. | 14 | 14 | .. |
| 12th February 1927 | .. | .. | .. | .. | .. | .. | .. | .. | .. | 1 | 1 | .. | 7 | 7 | .. |

REFERENCES

- Bindra, S. S. (1928). Studies on *Platyedra gossypiella* Saunders (Pink Boll-worm) in the Punjab. *Mem. Dept. Agri. India, Ent. ser.*, **10**, 6.
- Burt, B. C. (1922). Annual Report of the Secretary, Indian Central Cotton Committee, Bombay for 1922.
- Coad, B. R. and Cassidy, T. P. (1920). Cotton boll-weevil control by the use of poison. *U. S. Dept. of Agri. Bull.* 875.
- Desai, B. M. and Naik, K. B. (1925). Ridge cultivation in lower Gujarat. *Bom. Dept. of Agri. Bull.* No. 123.
- Fletcher, T. B. and Misra, C. S. (1921). Cotton boll-worms in India. *Bull.* No. 105, *Imp. Inst. of Agri. Res.*
- Folsom, W. J. (1927). Calcium arsenate as a cause of aphid infestation. *Jour. Eco. Ent.*, **20**, Dec. 1927.
- Gough, L. H. (1919). Proceedings of the Third Entomological Meeting at Pusa, India.
- Husam, M. A. and Mathur, U. B. (1923). Proceedings of the Entomological Meeting held at Pusa, India, in 1923.
- Javeri, T. N. (1921). Proceedings of the Entomological Meeting held at Pusa in 1921.
- Lefroy, H. M. (1906, 1). The Bombay locust. *Mem. Dept. Agri. Ind., Ent. ser.* **1**, No. 1.
- Lefroy, H. M. (1906, 2). An outbreak of cotton pests in the Punjab, 1905. *Bull.* No. 2 of *Agri. Res. Inst., Pusa*.
- Madan Mohan Lal (1913). Preliminary report on cotton boll-worm in the Punjab.
- Osborne, M. R. (1926). Comparative tests with sodium fluosilicate and calcium arsenate for control of the cotton boll-weevil. *Jour. Eco. Ent.*, **19**, No. 4.
- Patel, M. L. (1924). Studies in Gujarat cottons, Part II. *Mem. Dept. Agri. Ind., Bot. ser.*, **12**, No. 5.
- Power, F. B. and Chestnut, V. K. (1925). The odorous constituents of the cotton plant. Emanation of ammonia and trimethylamine from the living plants. *Jour. Amer. Chem. Soc.*, **47**, June 1925.
- Renough, W. (1907). Note on the cotton failure on account of boll-worm in the central and South-West Punjab in 1905 and on the results of the measures taken to prevent the recurrence in 1906. *Bull.* No. 1 of 1907 of *Dept. of Agri., Punjab*.
- Thakar, B. J. and Desai, M. H. (1929). Woolly mites (*Eriophyes gossypii*) on cotton. *Agri. Jour. Ind.*, **24**, Part III.
- Wardle, R. A. (1929). The problem of applied entomology.
- William, C. B. and Kirkpatrick, T. W. (1924). The Multiple temperature incubator. *Ministry of Agri. of Egypt. Tech. Bull.* 38.

PUBLICATIONS OF THE IMPERIAL COUNCIL OF AGRICULTURAL RESEARCH, INDIA

1. Agriculture and Livestock in India.

A bi-monthly Journal of Agriculture and Animal Husbandry for the general reader interested in Agriculture or Live-stock in India or the Tropics. (Established 1931. Published in January, March, May, July, September and November. Prepayable subscription Rs. 6 or 9s. 9d. per annum inclusive of Indian postage. Price per part Rs. 2 or 3s. 6d. inclusive of Indian postage.) Volumes I to V complete are available.

2. The Indian Journal of Agricultural Science.

A bi-monthly Scientific Journal of Agriculture and the allied sciences, mainly devoted to the publication of the results of original research and field experiments. (Established 1931. Published in February, April, June, August, October and December. Prepayable subscription Rs. 15 or 24s. per annum inclusive of Indian postage. Price per part Rs. 3 or 5s. 3d. inclusive of Indian postage.) Volumes I to V complete are available.

3. The Indian Journal of Veterinary Science and Animal Husbandry.

A quarterly Journal for the publication of scientific matter relating to the health nutrition and breeding of live-stock. (Established 1931. Published in March, June, September and December. Prepayable subscription Rs. 6 or 9s. 9d. inclusive of Indian postage. Price per part Rs. 2 or 3s. 6d. inclusive of Indian postage.) Volumes I to V complete are available.

4. Scientific Monographs of the Imperial Council of Agricultural Research.

- | | |
|-----------|---|
| NCAG—SM/1 | No. 1 The Fungi of India, By E. J. Butler, C.I.E., D.Sc., M.B., F.R.S. and G. R. Bisby, Ph.D. (1931). Price Rs. 6-12-0 or 11s (As. 8.) |
| NCAG—SM/2 | No. 2 Life-histories of Indian Microlepidoptera. Second Series: Alucitidae (Pterophoridae, Tortricina and Gelechiadae), By T. Bainbrigge Fletcher, R.N., F.I.S., F.E.S., F.Z.S. (1932) Price Rs. 3-4-0 or 5s. 6d. (As. 6.) |
| NCAG—SM/3 | No. 3. The Open Pan System of White-Sugar Manufacture, By R. C. Srivastava, B.Sc. (1935, 2nd edition). Price Rs. 3-2-0 or 5s. 6d. (As. 9.) |
| NCAG—SM/4 | No. 4. Life-histories of Indian Microlepidoptera: Cosmopterygidae to Neopseustidae, By T. Bainbrigge Fletcher, R.N., F.I.S., F.E.S., F.Z.S. (1933). Price Rs. 4-8-0 or 7s. 6d. (As. 8.) |
| NCAG—SM/5 | No. 5. The Bombay Grasses, By E. J. Blatter, S.J., Ph.D., F.I.S. and C. McCann, F.I.S. Illustrated by R. K. Bhide. Price Rs. 20-12-0 or 32s. 6d. |
| NCAG—SM/6 | No. 6. Helminth Parasites of the Domesticated Animals in India, By G. D. Bhalerao, M.Sc. Price Rs. 7-12-0 or 13s. 3d. |
| NCAG—SM/7 | No. 7. Influence of Manures on the Wilt Disease of <i>Cajanus indicus</i> Spreng. and the isolation of Types Resistant to the Disease, By W. McRae, M.A., D.Sc. (Edin.), F.I.S. and F. J. F. Shaw, D.Sc (Lond.), A.R.C.S., F.I.S. (1933). Price Rs. 2-4-0 or 4s. 3d. (As. 5.) |
| NCAG—SM/8 | No. 8. The Silk Industry of Japan with Notes on Observations in the United States of America, England, France and Italy, By C. C. Ghosh, B.A., F.E.S. (1933). Price Rs. 4 or 6s. 9d. (As. 7.) |
| NCAG—SM/9 | No. 9. Mechanical Cultivation in India. A History of the Large-Scale Experiments carried out by Burma-Shell Oil Storage and Distributing Company of India, Ltd., By C. P. G. Wade. Price Rs. 3-14-0 or 6s. 6d. |

5. Miscellaneous Bulletins of the Imperial Council of Agricultural Research.

- NCAG—MB/1 No. 1. List of Publications on Indian Entomology (1930), By the Imperial Entomologist, Pusa. Price As. 14 or 1s. 6d.
- NCAG—MB/2 No. 2. List of Publications on Indian Entomology (1931), By the Imperial Entomologist, Pusa. Price As. 8 or 10d.
- NCAG—MB/3 No. 3. List of Publications on Indian Entomology (1932), By the Imperial Entomologist, Pusa. Price As. 12 or 1s. 3d.
- NCAG—MB/4 No. 4. Host Plant Index of Indo-Ceylonese Coccidae, by S. Ramachandran, L.Ag. and T. V. Ramakrishna Ayyar, B.A., Ph.D., F.Z.S. Price Re. 1-10 or 2s. 9d.
- NCAG—MB/5 No. 5. List of Publications on Indian Entomology, 1933. Compiled by the Officiating Imperial Entomologist, Pusa. Price As. 9.
- NCAG—MB/6 No. 6. Bee-keeping, By C. C. Ghosh, B.A., F.E.S. (3rd edition). Price Re. 1-14 or 3s. 3d.
- NCAG—MB/7 No. 7. List of Publications on Indian Entomology, 1934. Compiled by the Officiating Imperial Entomologist and the Imperial Entomologist, Pusa. Price Re. 1-2-0 or 2s.
- NCAG—MB/8 No. 8. Selected clinical Articles, by Messrs. G. H. Sharma and R. L. Kaura. Price As. 8 or 10d.
- NCAG—MB/9 No. 9. Statistical Methods and their Application to Agronomy. A Bibliography, by K. K. Gulia Roy. (*In the Press.*)
- NCAG—MB/10 No. 10. Diseases of Sugarcane and Methods for their control, by L. S. Subramanian. (*In the Press.*)
- NCAG—MB/11 No. 11. Tables of Standard Errors of Mandelian Ratio, by Swarn Singh Purwall and P. Krishna Rao. Price As. 12 or 1s. 3d.
- NCAG—MB/12 No. 12. List of Publications on the Botany of Indian Crops, Part II, for the period 1928-32. By R. D. Bose. (*In the Press.*)

6. Annual Report of the Imperial Council of Agricultural Research.

- NCAG—AR/31 Annual Report of the Imperial Council of Agricultural Research for the years 1929-30 and 1930-31. Price As. 12 or 1s. 3d. (Anna 1.)
- NCAG—AR/32 Annual Report of the Imperial Council of Agricultural Research for the year 1931-32. Price As. 6 or 8d. (As. 2.)
- NCAG—AR/33 Annual Report of the Imperial Council of Agricultural Research for the year 1932-33. Price As. 6 or 8d.
- NCAG—AR/34 Annual Report of the Imperial Council of Agricultural Research for the year 1933-34.
- NCAG—AR/35 Annual Report of the Imperial Council of Agricultural Research for the year, 1934-35.

7. Review of Agricultural Operations in India.

- NCAG—R.1/29 Review of Agricultural Operations in India, 1928-29. Price Rs. 3-2 or 5s. 6d. (As. 8.)
- NCAG—R.1/31 Review of Agricultural Operations in India, 1929-31. Price Rs. 5 or 8s. 3d. (As. 10.)
- NCAG—R.1/33 Review of Agricultural Operations in India, 1931-33. Price Rs. 5-12 or 9s. 6d.

8. Proceedings of the Board of Agriculture and Animal Husbandry.

Proceedings of the First Meeting of the Animal Husbandry Wing of the Board of Agriculture and Animal Husbandry held at New Delhi from the 20th to 23rd February 1933, with Appendices. Price Rs. 5-14-0 or 9s. 6d.

A Description of the Imperial Institute of Veterinary Research, Muktesar and its sub-station the Imperial Veterinary Serum Institute, Izatnagar, By F. Ware, F.R.C.V.S. Price Re. 1-4 or 2s.

The Production of Cigarette Tobacco by Flue-curing, by F. J. F. Shaw, D.Sc., A.R.C.S., F.L.S. and Kashi Ram, Imp. Inst. Agr. Res. Pusa Bull. No. 187. Reprinted 1935. Price Re. 1 or 1s. 9d.

A Handbook of Statistics for use in Plant Breeding and Agricultural Problems, by Dr. F. J. F. Shaw, D.Sc., A.R.C.S., F.L.S., I.A.S. (*In the Press.*)

[Copies of the above publications can be had from :—

In India—Manager of Publications, Civil Lines, Delhi.

In Europe and America—From Public Department (Publications Branch), Office of the High Commissioner for India, India House, Aldwych, London, W. C. 2.]

